

12 Blackspot seabream (*Pagellus bogaraveo*)

12.1 Stocks description and management units

The stock structure of blackspot seabream in ICES area is still unknown. Thus, for stock assessment and scientific advice on management purposes ICES considers three different components : a) Subareas 6, 7, and 8; b) Subarea 9, and c) Subarea 10 (Azores region).

The interrelationships of the blackspot seabream from subareas 6, 7, and 8, and the northern part of Division 9.a, and their migratory movements within these areas have been observed by tagging studies (Gueguen, 1974). However, there is no evidence of movement to the southern part of 9.a where a target longline fisheries take places, which extends outside the ICES area.

Genetics studies show that there are no genetic differentiation between populations from different locations within the Azores region (east, central and west group of Islands, and Princessa Alice Bank) but there are genetic differences between Azores (ICES Subdivision 10.a.2) and mainland Portugal, ICES Division 9.a (Stockley et al., 2005). These results, combined with the known distribution of the species by depth, suggest that Subarea 10 component of this stock can effectively be considered as a separate assessment unit. Not genetic structure has been found on the Atlantic continental shelf with small genetic differentiation between the Mediterranean Sea and the Atlantic (Stockley et al., 2005, Pinera et al., 2007).

12.2 Blackspot seabream in Subareas 6, 7 & 8

12.2.1 The fishery

From the 1950s to the 1970s, the blackspot seabream was exploited mainly by French and Spanish bottom offshore trawlers, by artisanal pelagic trawlers in the eastern Bay of Biscay (ICES Divisions 8.a,b), and by Spanish longliners in the Cantabrian Sea (ICES Division 8.c), with smaller contributions from other fisheries (Lorance, 2011). Currently, EU Regulations state that no directed fisheries are permitted under the quota, therefore catches should be only bycatches.

In the period considered (1988–2020), most of the estimated landings from the Subareas 6, 7 and 8 were taken by Spain (70%), followed by France (18%), UK (10%) and Ireland (1%).

The fishery in Subareas 6, 7 and 8 strongly declined in the mid-1970s, and the stock is seriously depleted (Figure 12.2.1a and Table 12.2.1b). Since the 1980s, the species is mainly a bycatch from otter trawl, longline and gillnet fleets and only a few small-scale hand liners have been targeting the species. Since 1988 the landings from Subarea 8 represent 67% and 33% of total accumulated landings are from subareas 6 and 7. At present the blackspot seabream reported catches in these areas are almost all bycatches of longline and otter trawl fleets from France, Ireland and Spain.

12.2.2 Landings trends

Landings data by ICES Subareas reported to the working group are shown in Table 12.2.1a–c. Figure 12.2.1a presents an overview of the historical series of landings in Subareas 6, 7 and 8 since the middle of the last century. Figure 12.2.1b shows, in greater detail, landings of the same subareas since 1988. In 2014, UK (Scotland) reported landings for the first time in 7.j, and Netherlands since 2017 in Subarea 7. This ICES division represents part of the historical species distribution area (Olivier, 1928; Desbrosses, 1932).

For those three subareas combined, landings decreased from 461 t in 1989 to 52 t in 1996, increased again to a peak in 2007 (324 t) and then decreased in 2020 to 91 t in parallel to the reduction of the TAC in following years (from 350 t in 2003 and to 105 t in 2020).

12.2.3 ICES Advice

In 2018, ICES advised that when the precautionary approach is applied, there should be zero catch in each of the years 2019 and 2020.

12.2.4 Management

The EU TAC for subareas 6, 7, and 8 was set for the first time in 2003 and has been reducing since then from 350 t to 105 t in 2020. Landings in 2007, 2010, 2012, 2014, 2015, 2016 and 2018 were slightly above the TAC. A minimum landing size of 35 cm applied from 2010 to 2012 and a minimum conservation reference size of 33 cm applies since 11 May 2017 (commission implementing regulation (EU) 2017/787 of 8 May 2017).

Pagellus bogaraveo TACs and total landings in European countries in Subarea 27.6, 7, and 8 in recent years.

Pagellus bogaraveo		
year	TAC	landings
2003	350	129
2004	350	183
2005	298	158
2006	298	139
2007	298	324
2008	298	159
2009	253	203
2010	215	281
2011	215	177
2012	215	257
2013	196	295
2014	178	256
2015	169	177
2016	160	164
2017	144	126
2018	130	133
2019	117	98
2020	102	91

Under Common Fisheries Policy it is stated that "Recreational fisheries can have a significant impact on fish resources and Member States should, therefore, ensure that they are conducted in a manner that is compatible with the objectives of the CFP" (Regulation (EU) no 1380/2013 of the European Parliament and of the Council). Therefore, a short account of regulations relevant to blackspot seabream in recreational fisheries is given here.

The Irish Specimen Fish Committee recommends that all recreational catches be returned alive, and the SI No. 747 of 2004 forbids commercial catching of blackspot seabream except where it is less than 5% of the total catch. In France, specific regulation for blackspot seabream set in 2019 forbids the landings of individuals smaller than 35 cm and the fishing of this species from 1 of January to 30 of June. Moreover, the French regulation, forbids the catch, landing and sale of this species to the purse seine fleet and established several catch limits by trip or by year to the rest of the fleets (trawlers, gillnetters and liners).

Since 2019 Spain has been established closure areas with the aim to protect the juveniles of this species (MAPA 2019). The regulation bans the Spanish trawling and deep-water long-liners fleets to fish in several areas of the centre and west of Division 8.c from April to September. Spain also established annually a maximum catch per day to the vessels involved in the fishery in subareas 6, 7, 8.

12.2.5 Data available

12.2.5.1 Landings and discards

The Spanish, French and UK extended landing time-series of *P. bogaraveo* in Northeast Atlantic were updated (Figure 12.2.1b). In recent years landings have been dropping in accordance to the continuous reduction of the biannual TAC since 2003.

Historically, discards are considered negligible, and estimates are available since 2014 representing between 0.0 % and 2.7% of the annual catches in all subareas (Table 12.2.2). Discards resulting from low quotas are compulsory as the fishery for the species ceases. In 2015 and 2016, discards in French fisheries may have resulted from legal closures of quota (MEDDE, 2015; MEEM, 2016). As the blackspot seabream is a highly valued species, it is likely that these reported discards are carcasses in bad condition recovered from nets, misidentification of the species in on-board observation and discards related to low quotas. The table 12.2.3, show that since 2017 there were not catches inside the NEAFC Regulatory Area (RA)

Misidentification in on-board observer program may occur as *P. bogaraveo* occurs at low abundance and other similar sparids species may also occur (*P. acarne*, *P. erythrinus*, *P. bellotii* and *Pagrus pagrus*).

12.2.5.2 Length compositions

Length–frequency distribution of commercial landings and discards since 2015, are presented (Figure 12.2.2). Length frequency distribution of discards reported data in InterCatch in 2017 were very scarce, therefore no length distribution for this year is presented. No length–frequency distribution for discards were presented in 2020 as in this year reported discards were 0 .

12.2.5.3 Age compositions

No age data were available to the working group. No age estimations are carried out for this stock.

12.2.5.4 Weight-at-age

Mean size and weight-at-age (Table 12.2.4) derived from Guéguen (1969) and Krug (1998) were used by Lorance (2011) as input data for the yield-per-recruit model used to simulate the effect of fishing mortality on the blackspot seabream stock of Bay of Biscay.

12.2.5.5 Maturity and natural mortality

Natural mortality of 0.2 was estimated by Lorance (2011). M was derived from the presumed longevity in the population according the rule $M = 4.22/t_{max}$, where t is the maximum age in the population derived from data from many populations (Hewitt and Hoenig, 2005).

12.2.5.6 Catch, effort and research vessel data

At the current level of abundance, the blackspot seabream is rarely caught in the northern surveys by French EVHOE IBTS (Divisions 8.f,g,h,j; 8.a,b, and 7.d), Irish IGFS (Divisions 6.a South and 7.b), by Spanish Groundfish Survey in the Porcupine bank (SP-PorcGFS) in Divisions 7.c and 7.k, and is a scarce species in the Northern Spanish Shelf Groundfish Survey (SP-NGFS in Divisions 8c and 9a).

At the Northern Spanish Shelf Groundfish Survey *P. bogaraveo* is a scarce species. In 2019, both biomass (0.11 ± 0.11 Kg·haul⁻¹) and abundance (0.53 ± 0.53 ind·haul⁻¹) increased slightly after the decreasing trend from 2015, but for first time since 2013 no read sea breams were caught in 2020 (**Error! Reference source not found.**12.2.3, 12.2.4 and 12.2.5) (Ruiz-Pico et al 2021). Last information available indicated that specimens caught in 2019 ranged from 22 cm to 29 cm, with a mode in 25-26 cm (Figure 12.2.6) (Fernández-Zapico et al 2020).

In French surveys, similar to the current western IBTS, from early 1980s when the stocks were already low red seabream was still presented in 40–60% of the hauls. This proportion dropped to around zero by 1985 (Lorance, 2011). This observation indicates that the current survey is appropriate to detect and monitor a recovery of the stock if ever it happens. Catch of blackspot seabream in the EVHOE survey have been too rare to allow the calculation of a survey indicator. However, data from the survey are in accordance with a possible recent increase of the stock. In particular, a large catch of more than 1000 individuals occurred in the 2016 survey. Although, one single event is not significant, it is noteworthy that it occurred in the area where on-board observations of the species occur, and fishers report an increasing occurrence. These indications do not allow revising the stock status which should still be considered to lag below any possible reference point. However, they imply that a rebuilding has probably started. A quick appraisal of the level of occurrence that would be expected if the stock rebuilt to past levels can be found from two surveys carried out in the Bay of Biscay only in 1973 and 1976 with the same protocol and gear as the current EVHOE survey but covering only strata of Bay of Biscay shelf up to 200 m (Figure 12.2.7). In 1973 and 1976, blackspot seabream was caught in 25% and 55 % of the hauls respectively (Figure 12.2.8). Since the start of the current survey series in 1987, it has always been caught in less than 5% of the hauls in the same strata, some years not at all. In the same strata, it was caught in one out of more than 60 in each of 2015 and 2016. Therefore, a ten to thirty-fold increase in occurrence might occur to consider that the stock rebuilt to level from the 1960s and 1970s, where catch amounted to 15 000 t/year. The current monitoring with on-board observations and the EVHOE survey is insufficient to monitor this rebuilding accurately, while the stock is still low. The increase occurrence in on-board observations is however consistent with fishers reporting more encounter. If the increase persists, which is likely under the current management, occurrences in on-board observations and the survey might become significant in the next few years.

In the Irish IGFS (Divisions 6.a South and 7.b) is also a very scarce species and since 2010 only few kg in were caught in four years of the series. Also the occurrence along the whole stations in the survey is very low ranging since 2010 from 0% to 4.3% (Table 12.2.5)

12.2.6 Data analyses

Landings since 1988 are well below those recorded in the period from 1960 to 1986 in which landings ranged from 2000 t to up to 13 000 t (Figure 12.2.1a). Catches recorded in the surveys are very scarce and are mainly juveniles smaller than 30 cm.

In 2003 when TACs were set for this species there were conflicts between fishing métiers in this area small artisanal handliners requesting vessels targeting pelagic species, mostly sardine with trawls and seine, to avoid any bycatch of blackspot seabream. The introduction of the TAC and national quota had an impact on fishing practices.

In the same area, fishers report to encounter more frequently the species in recent years. This was investigated using French on-board observations (Figure 12.2.9). The method used consisted in estimating the proportion of fishing operations where the species was caught (landings and discards combined) in French on-board observations to the south of 49°N. The limit at 49°N north was set to include the south of the Celtic Sea to the West of Brittany, where the species was historically abundant. This was made for all bottom trawl types combined, and all bottom nets combined for years 2010 to 2016. Some increasing trend in the proportion of hauls with catch of the species can actually be seen for bottom trawls, although the proportion of positive hauls is still small (Figure 12.2.10).

12.2.7 Biological reference points

WKLIFE has not yet suggested methods to estimate biological reference points for stocks which have only landings data or are bycatch species in other fisheries. Therefore, no attempt was made to propose reference points for this stock.

12.2.8 Exploratory assessment

Ongoing studies carried out as part of the H2020 Pandora and the French National DynRose projects were presented to the group. This include an analysis of the essential habitats of the species and approaches to assess the current biomass.

The study of the habitats modelling applies several Species Distribution Model (SDM) in an Ensemble modelling approach. The study is carried out at the scale of whole species distribution area, including therefore not only the stock in the Celtic Sea and Bay of Biscay but also the area of the two other stock units considered by ICES (in Iberian and Azorean waters) and the Mediterranean western basin. Occurrence data from a number of sources including (1) French on-board observation, carried out in application of the EU data collection framework (DCF), (2) surveys, (3) CPUEs derived from the vessel monitoring system installed on Spanish artisanal vessels in the Strait of Gibraltar using GPRS/GSM (Burgos et al., 2013) and (4) data available from the WEB such as OBIS. Occurrence data were modelled using several physical chemical and biological environment variables including bathymetric, hydrological, seafloor and water data. Preliminary results suggest that only a low fraction of its potential habitat is occupied (realized habitat) by the blackspot seabream in the Bay of Biscay in recent years (Figure 12.2.11).

Approaches to assess the current biomass include acoustics and environmental DNA (eDNA) investigations. Acoustics surveys were carried out to the West of Brittany in 2019 and were presented during last year's meeting (ICES, 2020). In September 2020, a three days eDNA survey was carried out in the same area as the acoustic survey of 2019 (Figure 12.2.12). The results from the two approaches were consistent in terms of spatial distribution of the species. So far none of this method allow to derive a direct quantitative estimate of the biomass in the area surveyed and both have advantages and inconveniences. For acoustics, one drawback is that fishing

operation are needed for identification of echoes and their classification. In the rocky area surveyed, this was done by handlining, which appeared to be selective as more species were identified from eDNA. In particular, with eDNA seabass seemed to occur at a similar abundance as blackspot seabream in the surveyed area, while it was not caught on handlines and the two species may have similar echoes. eDNA has a number of advantages, it covers all species (from microbes to mammals), all habitats (no issue like such as not trawlable grounds and does not depend on behaviour (egg daily vertical migration) and does not need identification fishing. However, as no catches are implied, eDNA provides no information of population composition (size, sex).

12.2.9 Management considerations

In the 2014 advice, ICES recommend the establishment of a recovery plan for the stock. This stock is collapsed, and the advice is to reduce mortality by all means to allow the stock to rebuild, however neither a recovery plan nor scientific studies to support this recommendation have ever been applied in subareas 6, 7 and 8, only a minimum landing size of 35 cm was applied but only for the period from 2010–2012.

Measures should include protection for areas where juveniles occur. Recreational fisheries may be a significant proportion of the mortality of those juveniles owing to their coastal distribution. This was confirmed for the stock in Subarea 10 (Pinho, 2015).

Landings in 2007, 2010, 2012, 2014, 2015, 2016 and 2018 were slightly above the TAC.

12.2.10 References

- Burgos, C., J. Gil, and L. A. del Olmo. 2013. The Spanish blackspot seabream (*Pagellus bogaraveo*) fishery in the Strait of Gibraltar: spatial distribution and fishing effort derived from a small-scale GPRS/GSM based fisheries vessel monitoring system. *Aquatic Living Resources* 26:399–407.
- Desbrosses, P. 1932. La dorade commune (*Pagellus centrodontus* Delaroche) et sa pêche. *Revue des Travaux de l'Office des Pêches Maritimes* 5:167–222.
- Guéguen, J. 1969. Croissance de la dorade, *Pagellus centrodontus* Delaroche. *Revue des Travaux de l'Institut des Pêches Maritimes* 33:251–264.
- Hewitt, D. A. and J. M. Hoenig. 2005. "Comparison of two approaches for estimating natural mortality based on longevity." *Fishery Bulletin* 103: 433–437.
- Krug, H. M., D. Rosa, G. Menezes, and M. Pinho. 1998. Age and growth of some demersal species of the Azores. Page 11 in ICES science conference, ICES CMO:84,11pp.
- Lorance, P. 2011. "History and dynamics of the overexploitation of the blackspot seabream (*Pagellus bogaraveo*) in the Bay of Biscay." *ICES Journal of Marine Science* 68(2): 290–301.
- MAPA (Ministerio de Agricultura, Pesca y Alimentación) 2019. Orden APA/359/2019, de 26 de marzo, pola que se modifica la Orden AAA/661/2016, de 3 de abril, por la que se establecen criterios de desembarque de besugo capturado en aguas de la Unión y aguas internacionales de las zonas VI, VII y VIIdel Consejo Internacional para la Exploración del Mar (CIEM) en lo relativo al establecimiento de vedaen determinadas zonas del caladero Cantábrico Noroeste.
- MEDDE (Ministère de l'écologie, du développement durable et de l'énergie) 2015. Avis no 4 relatif à la fermeture de certains quotas et/ou sous-quotas de pêche pour l'année 2015, JORF n°0242 du 16 octobre 2016.
- MEEM (Ministère de l'Environnement, de l'Energie et de la Mer) 2016. Avis n° 24 relatif à la fermeture de certains quotas et/ou sous-quotas de pêche pour l'année 2016, JORF n°0242 du 16 octobre 2016.
- Olivier, R. 1928. Poissons de chalut, la dorade (*Pagellus centrodontus*). *Revue des Travaux de l'Office des Pêches Maritimes* 1:5–32.

- O. Fernández-Zapico, S. Ruiz-Pico, M. Blanco, I. Preciado, A. Punzón, F. Velasco. 2020. Results on Greater forkbeard (*Phycis blennoides*), Bluemouth (*Helicolenus dactylopterus*), Spanish ling (*Molva macrophthalma*) and Blackspot seabream (*Pagellus bogaraveo*) of the Northern Spanish Shelf Groundfish Survey. Working Document for the ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources Copenhagen, 24 April–1 May 2020. 19 pp.
- S. Ruiz-Pico, O. Fernández-Zapico, M. Blanco, A. Punzón, I. Preciado, J.M. Gonzalez Irusta, F. Velasco. 2021. Results on greater forkbeard (*Phycis blennoides*), Spanish ling (*Molva macrophthalma*), roughsnout grenadier (*Trachyrincus scabrus*), bluemouth (*Helicolenus dactylopterus*) and other scarce deep water species on the Northern Spanish Shelf Groundfish Survey. Working Document for the ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources By correspondence, 22th to 28th April, 2021. 19 pp.
- Pinera, J. A., G. Blanco, E. Vazquez, and J. A. Sanchez. 2007. Genetic diversity of blackspot seabream (*Pagellus bogaraveo*) populations off Spanish Coasts: a preliminary study. *Marine Biology* 151:2153-2158.
- Pinho, M. (2015). "Harvesting juveniles of blackspot seabream (*Pagellus bogaraveo*) in the Azores (Northeast Atlantic): biological implications, management, and life cycle considerations." *ICES Journal of Marine Science* [ICES J. Mar. Sci.].
- Stockley, B., G. Menezes, M. R. Pinho, and A. D. Rogers. 2005. Genetic population structure in the black-spot seabream (*Pagellus bogaraveo* Brunnich, 1768) from the NE Atlantic. *Marine Biology* 146:793-804.

12.2.11 Tables and Figures

Table 12.2.1a. Blackspot seabream in subareas 6 and 7; WG estimates of landings by country.

YEAR	FRANCE*	IRELAND	SPAIN	UK (E & W)	UK (Scot)	CH. ISLANDS*	NETHERLANDS	TOTAL
1988	52	0	47	153		0		252
1989	44	0	69	76		0		189
1990	22	3	73	36		0		134
1991	13	10	30	56		14		123
1992	6	16	18	0		0		40
1993	5	7	10	0		0		22
1994	0	0	9	0		1		10
1995	0	6	5	0		0		11
1996	0	4	24	1		0		29
1997	0	20	0	36				56
1998	0	4	7	6				17
1999	2	8	0	15				25
2000	4	n.a.	3	13				20
2001	2	11	2	37				52
2002	4	0	9	13				25
2003	13	0	7	20				40
2004	33		4	18				55
2005	29		4	7				41
2006	36	0	8	19				63
2007	46	0	27	57				130
2008	39	0	2	22				63
2009	34	1	16	10				61
2010	22	0	40	1				62
2011	21		11	4				37
2012	38		118					156
2013	28		146	4				178
2014	15		35	9	0			60

YEAR	FRANCE*	IRELAND	SPAIN	UK (E & W)	UK (Scot)	CH. ISLANDS*	NETHERLANDS	TOTAL
2015	13	0	21					34
2016	24	0	15	1	0			40
2017	15	1	19	1		0	0	37
2018	17	0	2	1			1	22
2019	19	0	15	1				35
2020	8		13	0				21

*Channel Islands

Table 12.2.1b. Blackspot seabream in Subarea 8; WG estimates of landings by country.

YEAR	FRANCE*	SPAIN	UK (E & W)	TOTAL
1988	37	91	9	137
1989	31	234	7	272
1990	15	280	17	312
1991	10	124	0	134
1992	5	119	0	124
1993	3	172	0	175
1994	0	131	0	131
1995	0	110	0	110
1996	0	23	0	23
1997	18	7	0	25
1998	18	86	0	104
1999	13	84	0	97
2000	11	189	0	200
2001	8	168	0	176
2002	10	111	0	121
2003	6	83	0	89
2004	37	82	8	128
2005	28	90	0	118
2006	20	57	0	77
2007	44	149	1	193
2008	55	40	0	95

YEAR	FRANCE*	SPAIN	UK (E & W))	TOTAL
2009	5	137	0	142
2010	61	157	0	218
2011	19	122	0	141
2012	18	82	0	101
2013	26	91	0	117
2014	36	161	0	196
2015	18	125	0	143
2016	7	117	0	124
2017	3	85	0	89
2018	6	105	0	111
2019	4	59	0	63
2020	4	59		63

Table 12.2.1c Blackspot seabream in Subareas 6, 7 and 8; WG estimates of landings by subarea.

YEAR	6 AND 7	8	TOTAL
1988	252	137	389
1989	189	272	461
1990	134	312	446
1991	123	134	257
1992	40	124	164
1993	22	175	197
1994	10	131	141
1995	11	110	121
1996	29	23	52
1997	56	25	81
1998	17	104	121
1999	25	97	122
2000	20	200	220
2001	52	176	227
2002	25	121	147

YEAR	6 AND 7	8	TOTAL
2003	40	89	129
2004	55	128	183
2005	41	118	158
2006	63	77	139
2007	130	193	324
2008	63	95	159
2009	61	142	203
2010	62	218	281
2011	37	141	177
2012	156	101	257
2013	178	117	295
2014	60	196	256
2015	34	143	177
2016	40	124	164
2017	37	89	126
2018	22	111	133
2019	35	33	98
2020	21	71	91

Table 12.2.2. Blackspot seabream in subareas 6, 7 and 8; WG estimates of discards in subareas 6, 7 and 8 since 2014.

	Discards (t)	Landings (t)	Catches (t)	Discards/Catches (%)
2014	2.40	256	258	0.9
2015	2.33	177	179	1.3
2016	0.91	164	165	0.6
2017	1.17	126	127	0.9
2018	2.3	133	136	1.7
2019	2.7	98	101	2.7
2020	0	91	91	0

Table 12.2.3. Blackspot seabream in Subareas 6, 7 and 8. Working group estimates of landings. Catches inside and outside the NEAFC Regulatory Area (RA) as estimated by ICES for the stock in WGDEEP.

WGDEEP Stock gfb.27.nea	Catch Inside NEAFC RA (t)	Catch Outside NEAFC RA (t)	Total Catches	Proportion of catch inside the NEAFC RA (%)	NEAFC RA areas where caught
2020	0	91	91	0%	
2019	0	98	98	0%	
2018	0	133	133	0%	
2017	0	126	126	0%	

Table 12.2.4 Mean size and weight-at-age of Blackspot seabream in Bay of Biscay. From Lorance (2011), derived from Guéguen (1969b) and Krug (1998).

Age group	Mean size (total length, cm)	Mean weight (g)	Proportion of mature females
0			0
1	11.2	18	0
2	17.6	72	0
3	22.3	149	0
4	26	239	0
5	29.2	342	0
6	31.9	449	0.007
7	34.3	562	0.05
8	36.1	658	0.15
9	37.9	765	0.31
10	39.5	870	0.45
11	40.9	969	0.54
12	42.3	1076	0.62
13	43.7	1190	0.68
14	44.8	1285	0.73
15	45.9	1386	0.77
16	46.7	1462	0.80
17	47.8	1572	0.83
18	49.2	1719	0.86
19	49.9	1796	0.88

Age group	Mean size (total length, cm)	Mean weight (g)	Proportion of mature females
20	50.2	1830	0.89

Table 12.2.5. Occurrence (kg and % of occurrence in the sampled stations) of the Blackspot seabream (*P. bogaraveo*) in Irish IGFS survey time-series (2010–2020).

	kg	% of occurrence in the stations
2010	0.2	0.8%
2011	0	0
2012	0.1	0.6%
2013	0	0
2014	0	00
2015	0	0
2016	2.1	2.4%
2017	8.2	4.3%
2018	0	0
2019	0	0
2020	0	0

Table 12.2.6. References and sources of reconstructed landings data in the Figure 12.2.1a.

France	<p>-Years 1977–1987: Landings of <i>P.bogaraveo</i> (sic?) from the Northeast Atlantic. M. Pinho, pers. com. Source: SGDeep 1995.</p> <p>-Years 1950–1984: Landings of <i>Pagellus</i> sp. ("seabreams") from the Northeast Atlantic. Source: Dardignac (1988), quoted by Castro (1990). SGDeep</p>
Portugal	<p>-Years 1948–1987 Subarea 10: Landings of <i>P.bogaraveo</i> (sic). M.Pinho, pers. com. Source: H. Krug (for 1948–1969) and SGDeep 1995 (for 1970–1987).</p> <p>-Years 1948–1987, Subarea 9: Landings of <i>P.bogaraveo</i> (sic?). M.Pinho, pers. com. Source: H. Krug (for 1948–1969) and SGDeep 1995 (for 1970–1987).</p>
Spain	<p>-Years 1960–1986: Landings of <i>Pagellus</i> sp. ("seabreams") from the Northeast Atlantic. Source: Anuarios de Pesca marítima. Castro (1990). SGDeep 1996. Table 12.2.3.</p> <p>-Years 1983–1987: Landings of <i>P.bogaraveo</i> (sic) from Division 9.a correspond only to southern 9.a (Tarifa and Algeciras ports). Source: Cofradías de Pescadores. (WD Gil, 2004) and Cofradías de Pescadores. (Lucio, 1996).</p> <p>-Years 1985–1987: Landings of <i>Pagellus</i> sp. (mainly <i>P. bogaraveo</i>). Source: SGDeep 1996. Table 12.2.4.</p> <p>-Years 1948–1984: Landings of <i>P.bogaraveo</i> (sic) from "Division 8.c" mainly Division 8.c (eastern) and Division VIIIb (southern) correspond only to the Basque</p>
UK	<p>-Years 1978–1987: Landings of <i>P.bogaraveo</i> (sic?) from the Northeast Atlantic. M .Pinho, pers. com. Source: SGDeep 1995.</p>
All countries	<p>-Years 1979–1985 SGDeep official data</p> <p>-Years 1988–2020 WGDeep official data</p>




Figure 12.2.1a. Blackspot seabream in Subareas 6, 7 and 8. Source of the reconstructed landings of blackspot seabream in the Bay of Biscay.





Figure 12.2.1b. Blackspot seabream landing trends in ICES Subareas 6 and 7 since 1988.






Figure 12.2.2. Length frequencies of the blackspot seabream in commercial catches, landings and discards since 2015, in Subareas 6, 7 and 8. No discards were reported for 2020.




Figure 12.2.3. Occurrence (%) of the Blackspot seabream (*P. bogaraveo*) in Northern Spanish Shelf survey time-series (1990–2020).




Figure 12.2.4. Evolution of Blackspot seabream (*P. bogaraveo*) mean stratified biomass (upper panel) and abundance (lower panel) in Northern Spanish Shelf survey time-series (1983–2020).




Figure 12.2.5. Catches in biomass of Blackspot seabream on the Northern Spanish Shelf bottom-trawl surveys, 2010–2020.




Figure 12.2.6. Mean stratified length distributions of Blackspot seabream (*P. bogaraveo*) in Northern Spanish Shelf surveys (2010–2019), no data before 2009.




Figure 12.2.7. Strata covering the Bay of Biscay shelf, sampled in the current EVHOE survey and in two previous surveys in 1973 and 1976.




Figure 12.2.8. Occurrences of Blackspot seabream in surveys carried out in 1973 and 1976 and in the EVHOE survey in 2015 and 2016.




Figure 12.2.9. Geographical distribution on catch of the Blackspot seabream in French on-board observations 2010–2016 in the Bay of Biscay and southern Celtic Sea, all métiers. (Grey) all haul/sets observed, (Blue crosses) hauls with catch of blackspot seabream, (Green dots) hauls with catch of blackspot seabream <20 cm which species identification may be uncertain.




Figure 12.2.10. Proportion of fishing operations with catch of Blackspot seabream in bottom trawls (left) and bottom net (right) in French fisheries to the south of 49°N (ICES divisions 8.a–d and the southern part of 7.d and 7.h–k).




Figure 12.2.11. Potential habitat of the blackspot seabream in the Mediterranean Sea, Azorean waters and European Atlantic shelf estimated from the ensemble modelling.




Figure 12.2.12. Number of eDNA copies (log scale) of blackspot seabream by location sampled in September 2020.

12.3 Blackspot seabream (*Pagellus bogaraveo*) in Subarea 9 (Atlantic Iberian waters)

12.3.1 The fishery

Pagellus bogaraveo is caught by Spanish and Portuguese fleets in Subarea 27.9. Spanish landings data from this area are available from 1983, Portuguese data from 1988 and Moroccan information from 2001. European landings in Subarea 27.9, most of which are taken with lines, are from Spain (~65%) and Portugal (~35%) 2015–2019.

An update of the available information on the Spanish target fishery, from the southern part of Subarea 27.9, Strait of Gibraltar region, has been provided to the WGDEEP (Gil *et al.*, WD 12 to the 2021 WGDEEP). Currently, less than 40 Spanish vessels are involved in the fishery. The fishing grounds of the Spanish fleet are on both sides of the Strait of Gibraltar and near, i.e. mostly less than 20 nautical miles, from the main ports (Tarifa and Algeciras). It should be noted that not all the catches/landings come exclusively from ICES Subarea 9 although it was considered to belong to the same stock, the fishing grounds encompass areas of different Regional Organizations/Commissions (ICES, GFCM and CECAF). Fishing takes advantage of the fluctuation of the tide at depths from 350 to 700 m with “*voracera*” gear, a mechanized handline. Since 2002 artisanal vessels from Conil port have joined the blackspot seabream fishery. Those vessels operate in other fishing grounds and use longlines. This section of the Spanish fleet is currently composed by about six vessels. Species landings are disaggregated into different commercial categories due to the wide size range of the catch and size-varying prices. Historically these categories have varied but from 1999 onwards have remained the same in all ports.

Since 2001, Moroccan longliners held a fishery in the Strait of Gibraltar area. These are about 102 vessels that are mainly based in Tangier. The average technical characteristics of these vessels are: 20 GRT and 160 HP. Moreover, 435 artisanal vessels (± 15 CV, ≤ 2 GRT and 4–6 m length) also target this species in the Strait of Gibraltar area (COPEMED II, 2015). The WGDEEP considers the account of Moroccan catches appropriate as the fishery operates in the same area as the Spanish fishery and obviously targets the same stock. Landings information until 2019 was also available from GFCM Report on Benchmark sessions for the assessment of blackspot seabream in GSAs 1-3 (2020).

Detailed information from Portuguese fisheries has been updated to the Working Group by Farias and Figueiredo (WD 6 to the 2021 WGDEEP). As well as in other Spanish places in Subarea 27.9, it is admitted that there are no fisheries targeting the blackspot seabream in Portugal mainland although the species can be seasonally targeted: the species is usually caught as bycatch of fisheries targeting other species. In mainland Portugal, most of species landings are as fresh specimens and are derived from the polyvalent fleet, which uses mainly longlines. The main landing ports (~89% of the species mainland Portugal total landings) from North to South are: Matosinhos (Portugal North), Aveiro, Nazaré and Peniche (Portugal Centre) and Sagres (Portugal Algarve).

In the Portuguese area of 27.9.a stock, Peniche is the most important landing port for blackspot seabream (landings between 1999 and 2019 represented nearly 50% of the Portuguese landings of the species. The species is mainly landed between December and March: this seasonal fishery pattern can reflect differences on the species’ availability (coinciding with the spawning season) or differences on skippers’ seasonal fishing grounds preferences (Farias and Figueiredo, WD 7 to the 2020 WGDEEP).

12.3.1.1 Landing trends

Since 1990, the maximum catch was reached in 1993–1994 and 1997 (about 1000 t) whereas the minimum (about 60 t) in 2019 and 2020 (Figure and Table 12.3.1). It should be noted that not all Spanish landings from the Strait of Gibraltar come from ICES Subarea 27.9. Moroccan landings from the Strait of Gibraltar area are supposed to be outside ICES Subarea 27.9: 2020 landings were not available.

12.3.2 Advice

The ICES advice for 2021 and 2022 was “that when the precautionary approach is applied, catches should be no more than 119 tonnes in each of the years 2021 and 2022. All catches are assumed to be landed.”

12.3.3 Management

Since 2003, TAC and Quotas have been applied to the blackspot seabream fishery in Subarea 27.9. The table below shows a summary of *P. bogaraveo* recent years' TACs and European countries landings in this Subarea.

Pagellus bogaraveo TACs and total landings in European countries in Subarea 27.9 in recent years.

P. bogaraveo	2014–2015		2016–2017		2018–2019		2020–2021	
ICES Subarea	TAC	Landings	TAC	Landings	TAC	Landings	TAC	Landings
9	780 – 374	262 –153 (142*)	183 – 174	165 (77*) – 130 (17*)	165 – 149	87 (8*) –56 (4*)	149 – 119	59 (3*) -

*from InterCatch info: landings from adjacent waters of the Strait of Gibraltar (FAO 34.1.11 and FAO 37.1.1).

There is a minimum conservation reference size of 33 cm for this species in the Regions 1–5 (as defined in Article 2 of Regulation (EC) No 850/98) since 11 May 2017 (Commission Implementing Regulation (EU) 2017/787 of 8 May 2017). This size coincides with the previously applied minimum size in the Mediterranean Sea. The European Commission granted the exemption for the Strait of Gibraltar target fishery, which is expressed in the discard plan for certain demersal fisheries in South-Western waters for the period 2019–2021 (Commission Delegated Regulation (EU) 2018/2033).

European landings have always been below the adopted TACs although these have been reduced over the years. However, in the year 2016 (considering other areas such as FAO 34.1.11 and FAO 37.1.1) European countries landings (242 t) are above the 2016 TAC (183 t) for ICES Subarea 27.9 (Figure 12.3.1).

12.3.4 Stock identity

Stock structure of the species in ICES Subarea 27.9 is still unknown. Genetic studies showed a restricted gene flow among the populations located in the Azores (ICES Division 27.10.a.2) and those on the Portuguese continental slope (ICES Division 27.9.a) and Madeira (CECAF FAO Division 34.1.2) (Stockley *et al.*, 2005; Piñera *et al.*, 2007). Recent genetic studies using mitochondrial control region indicated a similar genetic diversity among sampling sites in the NE Atlantic and the Mediterranean, and no differentiation between the Azores and the remaining locations (Robalo *et al.*, 2021).

In the Strait of Gibraltar area tagging surveys (56 days at sea in 2001, 2002, 2004, 2006 and 2008) have been conducted. A total of 4500 fish were tagged, of which 423 recaptures have been reported. The main results indicate the inexistence of significant movements. Although strict movements were noted: feeding grounds are distributed along the entire Strait of Gibraltar and the species seems to remain within this area as a resident population (Gil, 2006). Recaptures of tagged fish have also been reported by the Moroccan fishery.

Farias and Figueiredo (WD 14 to the WGDEEP 2019) presents information on blackspot seabream spatial distribution from Portuguese research surveys, considering the relative frequency of fishing hauls with species catch rates higher than 5 specimens in the 1990-2017 surveys. It is concluded that the species is not evenly distributed along the surveyed area, being more frequently caught at specific grounds, suggesting a patchy distribution. In the northern coast of Portugal, the species is caught down to 100 m deep, whereas preferred habitats are between 200 and 400 m deep in the southwestern coast (Figure 12.3.2). There is no evidence of movements between the northernmost component and the southern part of Subarea 27.9 where Spanish fishery takes place.

12.3.5 Data available

12.3.5.1 Landings and discards

Historical landing data series available to the Working Group are described in Section 12.3.1 and detailed in Table 12.3.1. It should be noted that since 2015 Spanish landings include adjacent areas outside ICES Subarea 27.9 (data are not separated in earlier years). In addition, Morocco landings from the Strait of Gibraltar area are available since 2001 (not in 2020), although fishing is supposed to have taken place outside ICES Subarea 9.

Portuguese and Spanish discard information was available to the Working Group from on-board sampling programme (EU DCF/NP). Given the low levels of discards, the discarded rate is admitted to be nearly zero for most assessment purposes and those that do occur are mainly related to catches of small individuals. Consequently all catches of blackspot seabream in management area 27.9a. are assumed to be landed.

12.3.5.2 Length compositions

Length frequencies of landings are available for the Spanish “*voracera*” blackspot seabream target fishery in the Strait of Gibraltar (1997–2020). Figure 12.3.3 show the updated length distribution data (from Gil *et al.*, WD 12 to the 2021 WGDEEP). The table below shows the mean and median landed size since 1997:

Summary statistics of *Pagellus bogaraveo* landed sizes by year since 1997.

Year	Mean	Std. Dev.	Median	Year	Mean	Std. Dev.	Median
1997	35.98	6.38	35	2009	38.29	6.23	37
1998	34.33	5.07	34	2010	36.03	5.28	35
1999	36.23	5.30	36	2011	36.33	6.36	34
2000	36.79	4.81	36	2012	36.40	5.91	35
2001	37.11	5.45	37	2013	34.80	3.64	34
2002	38.10	5.93	38	2014	37.11	5.14	36

Year	Mean	Std. Dev.	Median	Year	Mean	Std. Dev.	Median
2003	38.35	6.27	38	2015	39.15	5.79	38
2004	36.56	5.69	35	2016	37.47	5.28	37
2005	36.79	6.02	35	2017	37.72	4.37	37
2006	35.87	5.58	35	2018	37.84	4.67	37
2007	37.26	5.95	36	2019	37.27	4.21	37
2008	37.76	6.22	36	2020	37.37	4.30	37

Only one mean value (in 1998) is lower than the 2013 year's mean landing size. However, changes are small and gradual. There seem to be a long-term slight decline, despite the mean length ups and downs over the last decade (Figure 12.3.3).

Length the frequency distribution by fishing segment (polyvalent and trawlers) from 2014 till 2020 landings are presented in Figure 12.3.4 (D 6 to the WGDEEP 2021). Differences in length distribution between the polyvalent the trawl segments indicates that polyvalent fleet catch larger fish than the trawl fleet.

It should be noted that in 2020 sampling effort in both countries were low due to the covid-19 disruption: 4 samples (72 specimens) in the polyvalent, 4 samples (52 specimens) in the trawl and 7 samples (517 specimens) in the target "*voracera*" fishery of the Strait of Gibraltar.

12.3.5.3 Age compositions

No new information was presented to the group.

12.3.5.4 Weight-at-age

No new information was presented to the group.

12.3.5.5 Maturity and natural mortality

No new information was presented to the group.

12.3.5.6 Catch, effort and research vessel data

Figure 12.3.5 and Table 12.3.2 present CPUE information, restricted to the Strait of Gibraltar fishery (Gil *et al.*, WD 12 to the 2021 WGDEEP). Effort, as indicated, from sales sheets is not standardized and is potentially underestimated in some years as the effort unit chosen may be inappropriate while CPUE estimated from VMS analysis shows the same trend.

Farias and Figueiredo (WD 6 to the 2021 WGDEEP) identify two reference fleets landing at Peniche port: a total of 40 fishing vessels (with more than 9 fishing trips per year and more than 6 months with positive landings of the species) were selected for the polyvalent (longliners) while 21 fishing vessels (with more than 9 fishing trips per year and more than 5 months with positive landings of the species) were selected for the trawl fleet. The GLM estimates of the reference fleets' CPUE, considered as landed weight per fishing trip, for the selected model are also presented in the WD. Catch rates derived from longliners are slightly higher than those from trawl – this probably reflects a difference on the species length composition between the two fleets (Figure 12.3.6).

12.3.5.7 Data analyses

The stock identity still unclear and there is not certain linkages between the Strait of Gibraltar populations and the populations in the northern and central area of Subarea 27.9.

The trend is clear in the target fishery of the Strait of Gibraltar. Landings declined significantly until 2013 which may be considered as an indication of a substantial reduction in exploitable biomass. Current CPUE low levels may also be consistent with an overexploited population.

However, the analysis from the Portuguese (Peniche port) reference fleets' CPUE is not in accordance with the clear decreasing trend observed in the Strait of Gibraltar target fishery: long-lines and bottom trawl catch rates from West Portugal coast are relatively stable.

12.3.6 Management considerations

A TAC regime (119 t) was established for 2021 and 2022 for whole Subarea 27.9. Although the advice aims to reduce total catch within the whole fishing area, it should be noted that the current TAC does not limit the whole fishery because it only applies to Subarea 27.9, nevertheless catches in the GFCM area 37.1.1 and CECAF area 34.1.11 should be reported (Council Regulation (EU) 2016/2285). Recent landings are below the corresponding TAC levels but in 2016, European landings (including other areas such as FAO 34.1.11 and FAO 37.1.1) were above the 2016 TAC.

The combination of the minimum size of 33 cm for this species and the landing obligation (EU Regulation 2013/1380) might have an effect on certain fisheries: the exemption from the landing obligation of the target fishery of the Strait of Gibraltar ("*voracera*" gear) does not apply to other blackspot seabream catches in ICES Subarea 27.9.

It should be noted that GFCM started a work plan to establish a management plan for the Strait of Gibraltar target fishery in 2019 (Recommendation GFCM/41/2017/2 on the management of blackspot seabream fisheries in the Alboran Sea, geographical subareas 1 - 3, for a two-year transition period). The 2020 GFCM WGSAD endorsed the advice on the status of blackspot seabream in the Strait of Gibraltar – based on an update of the benchmarked gadget model – whereby the stock was considered in overexploitation and overexploited: F_{current} ($F_{\text{BAR4-16}} = 0.49$) is almost the double of the reference point ($F_{\text{MSY}} = 0.26$) estimated and agreed during the Benchmark process (GFCM, 2021). WGDEEP would like to express its concern on the fact that the population of blackspot seabream in the Strait of Gibraltar is being assessed within two different advisory bodies (ICES and GFCM), who derive scientific advice to managers: coordination between all parties would be welcomed.

As well as in other ICES Subareas (27.6, 27.7, 27.8 and 27.10), measures should include protection for areas where juveniles occur: recreational fisheries may be a significant proportion of the mortality of those juveniles owing to their coastal distribution.

Trends in abundance at the western coast of Portugal may not be properly represented by the trend in the Strait of Gibraltar: the CPUE of the Peniche reference fleets does suggest a different trend than the Strait of Gibraltar "*voracera*" fleet and there is no evidence of movements between the northernmost component and the southern part of Subarea 27.9, where Spanish fishery takes place. In fact, Strait of Gibraltar component might be more related to Mediterranean components. Therefore, it might not be appropriate to infer the stock status in all Division 9a from the Strait of Gibraltar target fishery CPUE.

12.3.7 Tables and Figures

Table 12.3.1. Blackspot seabream (*Pagellus bogaraveo*) in Subarea 27.9: Working Group estimates of landings (in tonnes). Spanish landings from 2012 are official statistics.

Year	Portugal	Spain	Morocco*	Unallocated	TOTAL
1983		101			101
1984		166			166
1985		196			196
1986		225			225
1987		296			296
1988	370	319			689
1989	260	416			676
1990	166	428			594
1991	109	423			532
1992	166	631			797
1993	235	765			1000
1994	150	854			1004
1995	204	625			829
1996	209	769			978
1997	203	808			1011
1998	357	520			877
1999	265	278			543
2000	83	338			421
2001	97	277	17		374 (17*)
2002	111	248	32		259 (32*)
2003	142	329	20		471 (20*)
2004	183	297	30		480 (30*)
2005	129	365	37		494 (37*)
2006	104	440	70		544 (70*)
2007	185	407	85		592 (85*)
2008	158	443	72		601 (72*)
2009	124	594	90		718 (90*)

Year	Portugal	Spain	Morocco*	Unallocated	TOTAL
2010	105	379	142		484 (142*)
2011	74	259	148		333 (148*)
2012	143	60	135	92	295 (135*)
2013	90	91	106		181 (106*)
2014	59	203	131		262 (131*)
2015	66	87 (142**)	224		295 (219*)
2016	70	95 (77**)	161		242 (161*)
2017	69	61 (18**)	190		148 (190*)
2018	58	29 (8**)	76		95 (76*)
2019	36	20 (4**)	119		60 (119*)
2020	43	16 (3**)	N/A		62 (N/A*)

*Morocco landings are available from the Benchmark workshop on blackspot seabream (GFCM 2020)

**Figures in brackets includes blackspot seabream from other areas (FAO 34.1.11. and FAO 37.1.1).

Table 12.3.2. Spanish “*voracera*” blackspot seabream fishery of the Strait of Gibraltar (ICES Subarea 27.9): Estimated CPUE using sales sheets or VMS data as effort unit (adapted from Gil *et al.*, WD 12 to the 2021 WGDEEP).

Year	cpue	VMS cpue
1983	78	
1984	76	
1985	71	
1986	61	
1987	76	
1988	73	
1989	89	
1990	77	
1991	70	
1992	86	
1993	85	
1994	94	
1995	60	
1996	104	

Year	cpue	VMS cpue
1997	77	
1998	61	
1999	55	
2000	45	
2001	56	
2002	47	
2003	53	
2004	47	
2005	68	
2006	70	
2007	51	
2008	52	
2009	67	55
2010	46	38
2011	42	31
2012	35	21
2013	30	14
2014	39	22
2015	49	32
2016	41	27
2017	33	14
2018	18	4
2019	24	8
2020	24	13




Figure 12.3.1. Blackspot seabream in ICES Subarea 27.9 (and adjacent waters): Total European landings (Morocco landings are not included) and EU TACs. Since 2015 landings from Strait of Gibraltar includes other areas (FAO 34.1.11 and FAO 37.1.1).




Figure 12.3.2. Blackspot seabream in ICES Subarea 9: Distribution of *Pagellus bogaraveo* along the Portuguese coast based on Portuguese surveys from the period between 1997-2011 and 2013-2017. The coloured blotches are hauls with *Pagellus bogaraveo* catches over 5 n.h-1. The colour intensity of the blotches reflects species occurrence (from Farias and Figueiredo, WD 14 to the 2019 WGDEEP).




Figure 12.3.3. Spanish “voracera” blackspot seabream fishery of the Strait of Gibraltar: 1983–2020 (from Gil *et al.*, WD 12 to the 2021 WGDEEP). Dashed line (at 33 cm) represents the current minimum landing size for the species in Atlantic NE and Mediterranean European waters. Red dot are the mean value while red line represents the median.




Figure 12.3.4. Peniche (Portugal) landing port: *Pagellus bogaraveo* length frequency distribution by fishing gear (polyvalent and trawl fleet) for the years 2014 to 2020 (from Farias and Figueiredo, WD 6 to the 2021 WGDEEP). Length classes are aggregated by 4 cms range (from 18-22 and 20-24 in polyvalent and trawl fleets, respectively).




Figure 12.3.5. Blackspot seabream in ICES Subarea 27.9: Spanish “voracera” target fishery of the Strait of Gibraltar estimated CPUE, using sales sheets (dashed line: 1983-2020) and VMS data as unit of effort (solid line: 2009-2020) (from Gil *et al.*, WD 12 to the 2021 WGDEEP).




Figure 12.3.6. Blackspot seabream in ICES Subarea 27.9: Standardized annual estimates of CPUE by fleet segment (polyvalent and trawl) from the Peniche’s port reference fleets in 2015 - 2020 (from Farias and Figueiredo, WD 6 to the 2021 WGDEEP).

12.3.8 References

- Farias, I. and I. Figueiredo. 2019. *Pagellus bogaraveo* in Portuguese continental waters (ICES Division 27.9.a). Working Document 14 to the 2019 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Farias, I. and I. Figueiredo. 2020. *Pagellus bogaraveo* in Portuguese continental waters (ICES Division 27.9.a). Working Document 7 to the 2020 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Farias, I. and I. Figueiredo. 2021. *Pagellus bogaraveo* in Portuguese continental waters (ICES Division 27.9.a). Working Document 6 to the 2021 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- GFCM. 2020. Report of the Benchmark sessions for the assessment of blackspot seabream in GSAs 1-3. General Fisheries Commission for the Mediterranean. 1–4 April 2019, 9-14 December 2019 and, 26 March, 8 April, 18 May and 28 May 2020.
- GFCM. 2021 (*in prep.*). Report of the Working Group on Stock Assessment of Demersal Species (WGSAD). General Fisheries Commission for the Mediterranean. 18–30 January 2021.
- Gil, J. 2006. *Biología y pesca del voraz [Pagellus bogaraveo (Brünnich, 1768)] en el Estrecho de Gibraltar*. Tesis Doctoral, Universidad de Cádiz (UCA, Spain).
- Gil, J., L. Rueda, J.J. Acosta, C. Farias and M. Soriano. 2021. The Blackspot seabream Spanish target fishery of the Strait of Gibraltar: updating the available information. Working Document 19 to the 2021 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Piñera, J.A., G. Blanco, E. Vázquez and J.A. Sánchez, J.A. 2007. Genetic diversity of blackspot seabream (*Pagellus bogaraveo*) populations off Spanish Coasts: a preliminary study. *Marine Biology* 151: 2153–2158
- Robalo, J.I., I. Farias, S.M. Francisco, K. Avellaneda, R. Castilho and I. Figueiredo. 2021. Genetic population structure of the Blackspot seabream (*Pagellus bogaraveo*): contribution of mtDNA control region to fisheries management, Mitochondrial DNA Part A, DOI: 10.1080/24701394.2021.1882445
- Stockley, B., G. Menezes, M.R. Pinho and A.D. Rogers. 2005. Genetic population structure in the blackspot sea bream (*Pagellus bogaraveo* Brünnich, 1768) from the NE Atlantic. *Marine Biology* 146: 793–804.

12.4 Blackspot seabream (*Pagellus bogaraveo*) in Division 10.a.2

12.4.1 The fishery

Blackspot seabream (*Pagellus bogaraveo*) has been exploited in the Azores (ICES Division 10.a.2), at least since the XVI century as part of the Azorean demersal fishery. A directed hook exploits the species and line fishery that encompasses two fleet components: the artisanal (handlines) and the longliners (Pinho and Menezes, 2009; Pinho *et al.*, 2014). Important expansion of the fishery to offshore seamounts occurred during the 2000s (Ordinance No. 101/2002). This expansion was particularly held by the longline fleet because of the regional spatial management measures introduced (Santos *et al.*, 2019). The artisanal fleet is composed of small open deck boats (<12 m) that operate in local areas near the coast of the islands using several types of handlines. Longliners are closed deck boats (>12 m) that operate in all areas but during the last years the fishery is only authorized to operate on offshore (>6 nm) banks and seamounts (Pinho *et al.*, 2014; Diogo *et al.*, 2015). The tuna fishery caught, until the end of the nineties, juveniles (age 0) of blackspot seabream as live bait, but in a seasonal and irregular way because these catches depend on tuna abundance and on the occurrence of other preferred bait species like *Trachurus picturatus* (Pinho *et al.*, 2014).

The Azorean demersal fishery is a multispecies and multigear fishery where *P. bogaraveo* is considered the target species. The effect of these characteristics on the dynamics of the target fishery is not well understood given the plasticity of the fishery to the target effect related with variability of abundance and markets (prices of the fish in general along the year).

Landings trends

Historically, landings increased from 400 t at the start of the eighties to approximately 1000 t at the start of the nineties (Figure 13.4.1). This increase was mainly due to the development of new markets, increased fish value, entry of new and modern boats, better professional education of the fisher and introduction of bottom longline gear, permitting the expansion of the exploitable area to deeper waters, banks, and seamounts as well as the expansion of the fishing season (ICES, 2006). Between 1990 and 2009 the annual landings have fluctuated around 1000 t, with a peak in 2005. During the period 2010–2012 the landings decreased significantly to an average of 641 t, which correspond to about 57% of the TAC adopted for that period, maintaining thereafter around this value due to the TAC introduced. Since 2005 a continuous decrease of the landings has been observed. Currently the fishery is highly constrained by management measures. Landings of the last four years (2017, 2018, 2019 and 2020) were: 499t, 445t, 474t and 491 respectively.

12.4.2 ICES Advice

ICES advised that when the precautionary approach is applied, catches should be no more than 610 tonnes in 2021 for area 10. All catches are assumed to be landed. The analysis for ICES advice have not been updated due to a lack of survey and fishery sampling data due to the Covid-19 disruption. The advice given in 2020 for 2021 remains valid and is rolled over for 2022.

12.4.3 Management

Under the European Union Common Fisheries policy, a TAC was introduced in 2003 (EC. Reg. 2340/2002). The recent time-series of TACs and landings from ICES Subarea 10 is given below.

Year	2007	2008	2009	2010	2011	2012	2013	2014
EU TAC	1136	1136	1136	1136	1136	1136	1022	920
Landings	1070	1089	1042	687	624	613	692	663
Year	2015	2016	2017	2018	2019	2020	2021	
EU TAC	678	507	517	517	576	553	*	
Landings	701	515	499	445	474	491		

***TAC for 2021 was not yet defined by the EC.**

Since 2003 deep-water fishing within 100 miles of the Azores baseline is restricted to vessels registered in the Azores under the management of fishing effort of the common fishery policy for deep-water species (EC. Reg. 1954/2003). In 2006 the Regional Azorean Government introduced a quota system by island and vessel. Specific access requirements and conditions applicable to fishing for deep-water stocks were established (EC. Reg. 2347/2002). Fishing with trawl gears (EC. Reg. 1811/2004) and bottom gillnets (EC. Reg. 91/2005) are forbidden in the Azores region.

In 2009, the Regional Government introduce (Ordinance No. 1/2010) new technical measures, including the minimum landing size (30 cm total length), area restrictions by vessel size and gear, and gear restrictions (hook size and maximum number of hooks on the longline gear). The seamount (Condor), located approximately 17 km to the southwest off Faial Island, was closed to fisheries (Ordinance No. 48/2010) to allow a multidisciplinary research (ecological, oceanography and geological). During 2015, 2016 and 2017 additional technical measures were introduced which included limitation of the fishing area for long-liners, update of the minimum landing size to 33 cm (Ordinance No. 120/2016) and introduction of marine protected areas for coastal and oceanic areas (Santos *et al.*, 2019). During 2017 new license limitations were introduced for littoral hook and line fisheries. Since 2018 the quota is managed by quarter, island and vessel. In 2019 some techniques measures have been changed by the Regional Government and European Union, as for example a closed season (Ordinance No. 74/2015) implemented in 2016, to reduce effort during the spawning aggregations (among January 15 and end of February), was revoked by Ordinance No. 63/2019 which allows fishing throughout the year. By the end of 2019 the Council Regulation (EU) 2019/1601 proceeded the reduction of Blackspot seabream fishery possibilities assigned to the European Union in 2020 from 576 t (EU Reg. 2019/124) to 553 t.

12.4.4 Data available

12.4.4.1 Landings and discards

Total annual landings data are available since 1980. However, detailed and precise landing data are available for the assessment since 1990 (WD15 Medeiros-Leal *et al.*, 2021). Landings from Area 10.a.2 are presented in the Table 12.4.1 and Figure 12.4.1.

Information on the discards in the longline fishery has been collected in the Azores by a team of observers on board the longline fleet. During 2018 about 6% (12.7 t) of the total landings were discarded.

12.4.4.2 Length compositions

No new information.

12.4.4.3 Age compositions

No new information

12.4.4.4 Weight-at-age

No new information.

12.4.4.5 Maturity, sex-ratio and natural mortality

No new information

12.4.4.6 Catch, effort and research vessel data

There is no new information.

12.4.5 Data analyses

Catches in recent years are highly constrained by severe management measures. There is no new information available and data analysis (standardized fishery cpue and survey trend) was reported in the WGDEEP report 2020 (ICES, 2020).

Exploratory analysis

The length-based indicators (LBI), total mortality (Z), yield-per-recruit (YPR) and SPICT production model were explored and reported in the WGDEEP report 2020 (ICES, 2020). The results from these methods seem to be all in agreement suggesting that the stock has been explored at or above the MSY level. There is no new information available and data analysis for updating this exploratory analysis.

12.4.6 References

- Diogo, H.; Pereira, J. G.; Higgins, R. M.; Canha, A.; Reis, D. 2015. History, effort distribution and landings in an artisanal bottom longline fishery: An empirical study from the North Atlantic Ocean. *Marine Policy*, 51: 75–85.
- ICES. 2006. Report of the Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources. ICES CM 2006/ACFM:28.
- ICES. 2018. Report of the Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP), ICES CM 2018/ACOM:
- Medeiros-Leal, W.M; Santos, R.V.S; Novoa-Pabon, A.M; Silva, H; Pinho, M.R. 2020. Data from the deep-water fishery of the Azores (ICES subdivision 27.10.a.2). Working Document 17 (WD17). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.
- Medeiros-Leal, W.M; Santos, R.V.S; Novoa-Pabon, A.M; Silva, H; Pinho, M.R. 2021. Updating Survey data from the Azores for deep-water species. Working Document 15 (WD15). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 22 April to 28 April 2021.
- Novoa-Pabon, A. M.; Medeiros-Leal, W.; Pinho, M. R; Santos, R. V. S. 2020. Updated standardized CPUE for blackspot seabream (sbr.27.10) caught by bottom longline fleet in the Azores (ICES Subdivision 27.10.a.2), 1990-2017. Working Document 13 (WD13). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.
- Ogle DH, Wheeler P, Dinno A (2020). *FSA: Fisheries Stock Analysis*. R package version 0.8.30, <https://github.com/droglenc/FSA>.

- Pinho, M.R; Novoa-Pabon, A; Gil, J; Krug, H. 2015. Catch curve analysis for the red black spot seabream (*Pagellus bogaraveo*) stock from the Azores (ICES Xa2). Working Document ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 20 to 27 March 2015.
- Pinho, M.R; Medeiros-Leal, W.M; Sigler, M.F; Santos, R.V.S; Novoa-Pabon, A.M. Menezes, G.M; Silva, H.M. Azorean Demersal Longline Survey Abundance Estimates: Procedures and Variability. in prep.
- Pinho, M. R.; Menezes, G. 2009. Pescaria de demersais dos Açores. Boletim do Núcleo Cultural da Horta 2009:85-102. ISSN 1646-0022.
- Pinho, M. R.; Diogo, H.; Carvalho, J.; Pereira, J. G. 2014. Harvesting juveniles of Red (Blackspot) seabream (*Pagellus bogaraveo*) in the Azores: Biological implications, management and life cycle considerations. *ICES Journal of Marine Science*, 71, 2448–2456. doi: 10.1093/icesjms/fsu089.
- Santos, R. V. S.; Silva, W. M.; Novoa-Pabon, A. M.; Silva, H. M.; Pinho, M. R. 2019. Long-term changes in the diversity, abundance and size composition of deep sea demersal teleosts from the Azores assessed through surveys and commercial landings. *Aquatic Living Resources*, 32, 25. doi: 10.1051/alr/2019022

12.4.7 Tables and Figures

Table 12.4.1. Historical landings of blackspot seabream *Pagellus bogaraveo* from the Azores (ICES Area 10.a.2).

Year	Azores (10.a.2)	Total
1980	415	415
1981	407	407
1982	369	369
1983	520	520
1984	700	700
1985	672	672
1986	730	730
1987	631	631
1988	637	637
1989	924	924
1990	889	889
1991	874	874
1992	1090	1090
1993	830	830
1994	989	989
1995	1115	1115
1996	1052	1052
1997	1012	1012
1998	1119	1119

1999	1222	1222
2000	947	924
2001	1034	1034
2002	1193	1193
2003	1068	1068
2004	1075	1075
2005	1113	1113
2006	958	958
2007	1063	1070
2008	1089	1089
2009	1042	1042
2010	687	687
2011	624	624
2012	613	613
2013	692	692
2014	663	663
2015	701	701
2016	515	515
2017	499	499
2018	445	445
2019	474	474
2020	491	491

Table 12.4.2. Nominal and standardized bottom longline fishery abundance index (scaled cpue to the mean) of the black-spot seabream *Pagellus bogaraveo* in Subarea 10.

YEAR	NOMINAL cpue	STANDARDIZED cpue	Lower CI	Upper CI
1990	0.92	0.97	0.87	1.08
1991	0.92	0.94	0.81	1.07
1992	0.96	0.98	0.78	1.17
1993	0.79	1.01	0.87	1.15
1994	0.97	1.01	0.84	1.18
1995	1.09	1.08	0.92	1.23

YEAR	NOMINAL cpue	STANDARDIZED cpue	Lower CI	Upper CI
1996	1.24	1.5	1.25	1.75
1997	1.63	1.32	1.1	1.53
1998	1.03	1.21	1.06	1.35
1999	1.1	1.3	1.16	1.44
2000	0.82	0.82	0.75	0.9
2001	1.12	0.96	0.84	1.07
2002	1.24	1.02	0.9	1.15
2003	0.98	1	0.91	1.1
2004	1.42	1.08	0.96	1.19
2005	1.71	1.16	1.06	1.27
2006	1.26	0.95	0.86	1.04
2007	1.34	1.22	1.09	1.36
2008	1.21	1.13	1.02	1.24
2009	1.18	0.96	0.88	1.05
2010	0.62	0.72	0.66	0.78
2011	0.59	0.76	0.69	0.82
2012	0.62	0.81	0.74	0.88
2013	0.64	0.91	0.83	0.99
2014	0.67	0.83	0.76	0.90
2015	0.56	0.74	0.68	0.80
2016	0.39	0.61	0.56	0.67
2017	0.48	0.59	0.60	0.57
2018	na	na	na	na
2019	na	na	na	na
2020	na	na	na	na
na – not available				

Table 12.4.3. Survey relative abundance index in number of blackspot seabream *Pagellus bogaraveo* from the Azores (ICES Area 10.a.2).

Year	Lower	Index	Upper
1995	88	107	125
1996	33	41	49
1997	33	46	58
1998	na	na	na
1999	80	112	143
2000	38	52	67
2001	58	68	78
2002	126	138	150
2003	66	86	103
2004	69	94	120
2005	118	143	166
2006	na	na	na
2007	54	79	106
2008	84	101	119
2009	na	na	na
2010	53	67	83
2011	52	70	87
2012	49	58	69
2013	38	47	55
2014	na	na	na
2015	na	na	na
2016	114	137	158
2017	125	155	182
2018	92	114	136
2019	134	166	195
2020	na	na	na

na = Not available.




Figure 12.4.1. Historical landings of blackspot seabream *Pagellus bogaraveo* from the Azores (ICES Area 10.a.2).