

11 Alfonsinos/Golden eye perch (*Beryx* spp.) in all ecoregions

11.1 The fishery

Alfonsinos, *Beryx splendens* and *Beryx decadactylus*, are generally considered as bycatch species in the demersal trawl and longline mixed fisheries targeting other deep-water species. For most of the fisheries, the catches of alfonsinos are reported under a single category, as *Beryx* spp.

The proportions of each species in the catches are not well known. Detailed landings data by species are available only for the Portuguese (Azores) hook and line fishery in Division 10a2, where the landings of *B. decadactylus* averaged 20% of the catches of both species in the last twenty years, and for the Russian trawl fishery that targeted *B. splendens*. Portuguese, Spanish and French trawlers and longliners are the main fleets involved in this fishery.

Landings from a targeted fishery by Russian vessels operating in the NEAFC area (ICES Division 10.b) were available for the period 1993 - 2000 and some minor landings as bycatch in fisheries targeting other species since 2000. Since 2000 Currently, there are no target fisheries occurring in Mid-Atlantic Ridge (NEAFC) (see Section 4).

Currently landings are reported from bycatch fisheries occurring in the NEAFC regulatory area (RA) of ICES Division 10.b from Faroese vessels and in the EEZ of Portugal (Subarea 9), Spain (6, 7, 8 and 9), France (6, 7 and 8), and from a small-scale target fishery based in the Azores operation in Division 10.a (See Table 11.1 c, d and e).

11.1.1 Landings trends

The available landings data for Alfonsinos, (*Beryx* spp.), by ICES subarea/division as officially reported to ICES or to the working group, are presented in Tables 11.1(a–g), 11.2 and 11.3 and Figures 11.1–11.5. Total landings are stabilized since 2005, due to management measures introduced (TAC/quotas and effort regulation), being around 337 t between 2005 and 2020, with high landings during 2012 (605 t). Current catches are 171 t. Faroes reported a landing of 141 t for 2015, 48 t for 2016, 5 t for 2019 from area 10.b. There were no landings reported during 2020.

11.1.2 ICES Advice

ICES advice that when the precautionary approach is applied, landings should be no more than 224 t in each of the years 2021 and 2022. ICES cannot quantify the corresponding catches.

11.2 Management

Fishing with trawl gears is forbidden in the Azores region (EC. Reg. 1568/2005). A box of 100 miles limiting the deep-water fishing to vessels registered in the Azores was created in 2003 under the management of fishing effort of the CFP for deep-water species (EC. Reg. 1954/2003). An EU TAC of 187 t for EC vessels is in force for the period 2021–2022 (see historical developments on the table down).

Technical measures were introduced in the Azores in 1998. During 2009 new measures were adopted, particularly to control the effort of longliners through restrictions on fishing area,

minimum length, gear and effort. These measures were updated during 2015–2019. A network of MPAs was implemented on the Azores with closed access to deep-water fisheries (including Sedlo, D. J. Castro and Formigas seamounts). The seamount (Condor) was closed to the fishery.

NEAFC adopted effort regulations for fisheries targeting deep-water species and in closed areas to protect vulnerable habitats on the RA. (http://neafc.org/managing_fisheries/measures/current).

Regulation	Species	Year	ICES Area	TAC	Landings
Reg 2270/2004	<i>Beryx</i> sp	2005	3, 4, 5, 6, 7, 8, 9, 10, 12	328	422
	<i>Beryx</i> sp	2006	3, 4, 5, 6, 7, 8, 9, 10, 12	328	367
Reg 2015/2006	<i>Beryx</i> sp	2007	3, 4, 5, 6, 7, 8, 9, 10, 12	328	396
	<i>Beryx</i> sp	2008	3, 4, 5, 6, 7, 8, 9, 10, 12	328	405
Reg 1359/2008	<i>Beryx</i> sp	2009	3, 4, 5, 6, 7, 8, 9, 10, 12	328	382
	<i>Beryx</i> sp	2010	3, 4, 5, 6, 7, 8, 9, 10, 12	328	296
Reg 1225/2010	<i>Beryx</i> sp	2011	3, 4, 5, 6, 7, 8, 9, 10, 12	328	331
	<i>Beryx</i> sp	2012	3, 4, 5, 6, 7, 8, 9, 10, 12	328	596
Reg 1262/2012	<i>Beryx</i> sp	2013	3, 4, 5, 6, 7, 8, 9, 10, 12	312	272
	<i>Beryx</i> sp	2014	3, 4, 5, 6, 7, 8, 9, 10, 12	296	282
Reg. 1367/2014	<i>Beryx</i> sp	2015	3, 4, 5, 6, 7, 8, 9, 10, 12	296	224
	<i>Beryx</i> sp	2016	3, 4, 5, 6, 7, 8, 9, 10, 12	296	252
Reg. 2285/2016	<i>Beryx</i> sp	2017	3, 4, 5, 6, 7, 8, 9, 10, 12	280	240
	<i>Beryx</i> sp	2018	3, 4, 5, 6, 7, 8, 9, 10, 12	280	263
Reg. 2025/2018	<i>Beryx</i> sp	2019	3, 4, 5, 6, 7, 8, 9, 10, 12	252	294
	<i>Beryx</i> sp	2020	3, 4, 5, 6, 7, 8, 9, 10, 12	252	171
Reg.	<i>Beryx</i> sp	2021	3, 4, 5, 6, 7, 8, 9, 10, 12	187	
Reg.	<i>Beryx</i> sp	2022	3, 4, 5, 6, 7, 8, 9, 10, 12		

11.3 Stock identity

No new information.

11.4 Data available

11.4.1 Landings and discards

Tables 11.1a–g, describe the alfoncinos landings by subarea and country. In 2014 discards estimates for the Azorean longliners were reported (WD, Pinho, 2014). Annual longline discard estimates by year for the sampled trip vessels with alfoncinos catches during the period 2004–2011

range from 0.8% to 8.6% for *B splendens* and 0.07% to 10.2% for the *B. decadactylus* (Table 11.4). In 2018, discards of combined alfonsoinos from longline fisheries represented about 5.8% (8.6 t) of total landings. These discards are mostly a result of the Azorean management measures such as TAC and minimum length.

11.4.2 Length compositions

No new information.

11.4.3 Age compositions

No new information

11.4.4 Weight-at-age

No new information.

11.4.5 Maturity, sex-ratio, length–weight and natural mortality

No new information.

11.5 Data analyses

No new information

11.6 Comments on the assessment

No assessment was carryout this here.

11.7 Management considerations

The spatial distribution of the two *Beryx* species is closely associated with seamounts. This behaviour as well as their life history make them particularly vulnerable to exploitation been easily overexploited by trawl fishing. It is admitted that both species can only sustain low rates of exploitation.

The population dynamics *Beryx* species are uncertain. Recent age estimates suggest high longevity (>50 years), while other estimates suggest a longevity of ~15 years.

Because of their biology and aggregative behaviour, fisheries on those species should not be allowed to expand above current levels unless it can be demonstrated that such expansion is sustainable. Furthermore, to prevent wiping out entire subpopulations that have not yet been mapped and assessed the exploitation of new seamounts should not be allowed.

11.8 References

ICES. 2018. Report of the Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP), 11- 18 April 2018, ICES Headquarters, Copenhagen. ICES CM 2018/ACOM:14.

Medeiros-Leal, W.M.M; Santos, R.V.S; Novoa-Pabon, A.M; Silva, H; Pinho, M.R. 2020. Updating Survey data from the Azores for deep-water species. Working Document (WD). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.

Medeiros-Leal, W.M.M; Santos, R.V.S; Novoa-Pabon, A.M; Silva, H; Pinho, M.R. 2020. Data from deep-water fishery of the Azores (ICES subdivision 27.10.a.2). Working Document (WD). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.

Santos, R. V. S.; Novoa-Pabon, A. M.; Silva, H. M.; Pinho, M. R. 2019. Can we consider the stocks of alfonsoinos *Beryx splendens* and *Beryx decadactylus* from the Azores a discrete Fishery Management Unit? *Journal of Fish Biology*, 94(6): 993-1000. <https://doi.org/10.1111/jfb.13937>

Santos, R. V. S.; Medeiros-Leal, W., Novoa-Pabon, A. M.; Pinho, M. R. 2020. Updated standardized CPUE for *Beryx decadactylus* caught by bottom longline fleet in the Azores (ICES Subdivision 27.10.a.2), 1990-2017. Working Document (WD). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.

Santos, R. V. S.; Medeiros-Leal, W., Novoa-Pabon, A. M.; Pinho, M. R. 2020. Updated standardized CPUE for *Beryx* spp. (alf.27.nea) caught by bottom longline fleet in the Azores (ICES Subdivision 27.10.a.2), 1990-2017. Working Document (WD). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.

Santos, R. V. S.; Medeiros-Leal, W., Novoa-Pabon, A. M.; Pinho, M. R. 2020. Updated standardized CPUE for *Beryx splendens* caught by bottom longline fleet in the Azores (ICES Subdivision 27.10.a.2), 1990-2017. Working Document (WD). ICES Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP), 24 April to 01 May 2020.

11.9 Tables and Figures

Table 11.1a. Landings (tonnes) of *Beryx* spp. from Subarea 4.

YEAR	FRANCE	TOTAL
1988	0	0
1989	0	0
1990	1	1
1991	0	0
1992	2	2
1993	0	0
1994	0	0
1995	0	0
1996	0	0
1997	0	0
1998	0	0
1999	0	0
2000	0	0

YEAR	FRANCE	TOTAL
2001	0	0
2002	0	0
2003	0	0
2004	0	0
2005	0	0
2006	0	0
2007	0	0
2008	0	0
2009	0	0
2010	0	0
2011	0	0
2012	0	0
2013	0	0
2014	0	0
2015	0	0
2016	0	0
2017	0	0
2018	3	3
2019	0	0
2020*	0	0

*Preliminary.

Table 11.1b. Alfonsinos (*Beryx spp.*) from Division 5.b.

YEAR	FAROES	FRANCE	TOTAL
988			0
1989			0
1990		5	5
1991		0	0
1992		4	4
1993		0	0

YEAR	FAROEES	FRANCE	TOTAL
1994		0	0
1995	1	0	1
1996	0	0	0
1997	0	0	0
1998	0	0	0
1999	0	0	0
2000	0	0	0
2001	0	0	0
2002	0	0	0
2003	0	0	0
2004	0	0	0
2005	0	0	0
2006	0	0	0
2007	0	0	0
2008	0	0	0
2009	0	0	0
2010	0	0	0
2011	0	0	0
2012	0	0	0
2013	0	0	0
2014	0	0	0
2015	0	0	0
2016	0	0	0
2017	0	0	0
2018	0	0	0
2019	0	0.1	0.1
2020*	0	0	0

*Preliminary.

Table 11.1c. Alfonsinos (*Beryx* spp.) from Subareas 6 and 7.

YEAR	FRANCE	E & W	SPAIN	IRELAND	SCOTLAND	TOTAL
1988						0
1989	12					12
1990	8					8
1991						0
1992	3					3
1993	0		1			1
1994	0		5			5
1995	0		3			3
1996	0		178			178
1997	17	4	5			26
1998	10	0	71			81
1999	55	0	20			75
2000	31	2	100			133
2001	51	13	116			180
2002	35	15	45			95
2003	20	5	55	4		84
2004	15	3	46			64
2005	15	0	55	0		70
2006	27	0	51	0		78
2007	17	1	47	0		65
2008	22	0	32	0		54
2009	9	0	0	0	1	10
2010	4	0	0	0	1	5
2011	7	0	33	0	0	40
2012	4	0	337	0	0	341
2013	14	1	33	0	0	77
2014	10	0	38	0	0	49
2015	6	0		6	0	12
2016	5	0.45	13	0	1	20

YEAR	FRANCE	E & W	SPAIN	IRELAND	SCOTLAND	TOTAL
2017	7	0	11	0	0	18
2018	27	0.209	19	0	0	46
2019	57		24	0	0	81
2020*	7	3.1	0	0	0	11

*Preliminary.

Table 11.1d. Alfonsinos (*Beryx spp.*) from Subareas 8 and 9.

YEAR	FRANCE	PORTUGAL	SPAIN	E & W	TOTAL
1988					0
1989					0
1990	1				1
1991					0
1992	1				1
1993	0				0
1994	0		2		2
1995	0	75	7		82
1996	0	43	45		88
1997	69	35	31		135
1998	1	9	258		268
1999	11	29	161		201
2000	7	40	117	4	168
2001	6	43	179	0	228
2002	13	60	151	14	238
2003	10	0	95	0	105
2004	21	53	209	0	283
2005	9	45	141	0	195
2006	8	20	64	3	97
2007	8	45	67	0	120
2008	5	42	54	0	101
2009	1	42	18	0	61

YEAR	FRANCE	PORTUGAL	SPAIN	E & W	TOTAL
2010	12	27	1	0	41
2011	4	21	40	0	65
2012	4	11	27	0	42
2013	5	17	4	0	26
2014	3	18	81	0	102
2015	3	0	59		61
2016	3	1	71	0	76
2017	3	2	67	0	73
2018	6	0	52	0	58
2019	5	10	55		70
2020*	10	11			21

* Preliminary.

Table 11.1e. Alfonsinos (*Beryx* spp.) from Subarea 10.

	10.a	10.b				
YEAR	PORTUGAL	FAROEES	NORWAY	RUSSIA**	E & W	TOTAL
1988	225					225
1989	260					260
1990	338					338
1991	371					371
1992	450					450
1993	533		195			728
1994	644		0	837		1481
1995	529	0	0	200		729
1996	550	0	0	960		1510
1997	379	5	0			384
1998	229	0	0			229
1999	175	0	0	550		725
2000	203	0	0	266	15	484
2001	199	0	0	0	0	199

	10.a	10.b				
2002	243	0	0	0	0	243
2003	172	0	0	0	0	172
2004	139	0	0	0	0	139
2005	157	0	0	0	0	157
2006	192	0	0	0	0	192
2007	211	0	0	0	0	211
2008	250	2	0	0	0	252
2009	311	1	0	0	0	312
2010	240	0	0	5	0	245
2011	226	4	0	5	0	235
2012	213	10	0	0	0	222
2013	168	0	0	0	0	168
2014	131	0	0	0	0	131
2015	151	141	0	0	0	292
2016	156	48	0	0	0	204
2017	149	0	0	0	0	149
2018	159	0	0	0	0	159
2019	138	5				143
2020*	139					139

* Preliminary.

** Not official data from ICES Area 10.b.

Table 11.1f. Alfonsinos (*Beryx* spp.) from Subarea 12.

YEAR	FAROEES	TOTAL
1988		
1989		
1990		
1991		
1992		
1993		
1994		

YEAR	FAROEES	TOTAL
1995	2	2
1996	0	0
1997	0	0
1998	0	0
1999	0	0
2000	0	0
2001	0	0
2002	0	0
2003	0	0
2004	0	0
2005	0	0
2006	0	0
2007	0	0
2008	0	0
2009	0	0
2010	0	0
2011	2	2
2012	0	0
2013	0	0
2014	0	0
2015	0	0
2016	0	0
2017	0	0
2018	0	0
2019	0	0
2020*		

* Preliminary.

Table 11.1g. Landings of Alfonsinos (*Beryx* spp.) from Madeira (Portugal) outside the ICES area.

YEAR	<i>B. splendens</i>	<i>B. decadactylus</i>	TOTAL
1988*			
1989*			
1990*			
1991*			
1992*			
1993*			
1994*			
1995*			
1996*			
1997*			
1998*			
1999*			
2000*			
2001*			
2002*			
2003*			
2004*			
2005*			
2006*			
2007*			
2008	290	342	632
2009	88	16	104
2010	355	17	372
2011	79	137	216
2012	228	51	279
2013	38	11	49
2014	140	26	166
2015	63	12	75
2016	58	20	78

YEAR	<i>B. splendens</i>	<i>B. decadactylus</i>	TOTAL
2017	41	78	119
2018	234	83	317
2019	90	146	236
2020	12	11	23

* No information.

Table 11.1h. Reported landings for the alfonsinos, (*Beryx* spp.), by ICES subarea/division.

YEAR	4	5.b	6+7	8+9	10.a	10.b	12	TOTAL
1988			0	0	225	0		225
1989			12	0	260	0		272
1990	1	5	8	1	338	0		353
1991			0	0	371	0		371
1992	2	4	3	1	450	0		460
1993			1	0	533	195		729
1994			5	2	644	837		1488
1995		1	3	82	529	200	2	817
1996			178	88	550	960	0	1776
1997			26	135	379	5	0	545
1998			81	268	229	0	0	579
1999			75	201	175	550	0	1001
2000			133	168	203	281	0	785
2001			180	228	199	0	0	607
2002			95	238	243	0	0	577
2003			84	105	172	0	0	361
2004			64	283	139	0	0	485
2005			70	195	157	0	0	422
2006			78	97	192	0	0	367
2007			65	120	211	0	0	396
2008	0	0	54	101	250	2	0	407
2009	0	0	10	61	311	1	0	383

YEAR	4	5.b	6+7	8+9	10.a	10.b	12	TOTAL
2010	0	0	5	41	240	5	0	291
2011	0	0	40	65	226	9	2	342
2012	0	0	341	42	213	10	0	605
2013	0	0	77	26	168	0	0	272
2014	0	0	49	102	131	0	0	282
2015	0	0	12	61	151	141	0	365
2016	0	0	20	76	156	48	0	300
2017	0	0	18	73	149	0	0	240
2018	0	0	46	58	159		0	263
2019	0	0	81	70	143		0	294
2020*	0	0	11	21	139	0	0	171

*Preliminary.

Table 11.3. Reported landings of *Beryx splendens* and *B. decadactylus* in the Azores (ICES Division 10a2).

YEAR	<i>B. Splendens</i>	<i>B. Decadactylus</i>	TOTAL
1988	122	103	225
1989	113	147	260
1990	137	201	338
1991	203	168	371
1992	274	176	450
1993	316	217	533
1994	410	234	644
1995	335	194	529
1996	379	171	550
1997	268	111	379
1998	161	68	229
1999	119	56	175
2000	168	35	203
2001	182	17	199
2002	223	20	243

YEAR	<i>B. Splendens</i>	<i>B. Decadactylus</i>	TOTAL
2003	150	22	172
2004	110	29	139
2005	134	23	157
2006	152	40	192
2007	165	46	211
2008	187	63	250
2009	243	68	311
2010	189	51	240
2011	179	47	226
2012	175	37	213
2013	140	28	168
2014	109	22	131
2015	120	31	151
2016	127	29	156
2017	119	30	149
2018	107	50	157
2019	92	46	138
2020*	67	72	139

*Preliminary.

Table 11.4. Annual percentage of *Beryx* spp. discarded by year in the Azores (ICES Division 10a2) from the sampled trip vessels that caught and discard alfonsinos.

SPECIES	2004	2005	2006	2007	2008	2009	2010	2011
<i>Beryx splendens</i>	1,79	1,87	1,55	1,02	1,19	8,64	4,69	0,76
<i>Beryx decadactylus</i>	0,37	0,07	1,31	0,14	0,57	10,18	2,36	0,95




Figure 11.1. Catches of alfonsinos by French, Irish, UK (England and Wales and Scotland) and Icelandic vessels, 2006.




Figure 11.2. Catches of alfonsinos by French, Irish, UK (England and Wales and Scotland) and Icelandic vessels, 2007.




Figure 11.3. Catches of alfonsinos by Azores vessels, 2008–2011 (ICES, 10a2).




Figure 11.4. Reported landings for the alfonsinos, (*Beryx* spp), by ICES subarea/division.