

## Stock Annex: Herring (*Clupea harengus*) in subdivisions 30 and 31 (Gulf of Bothnia)

---

Stock specific documentation of standard assessment procedures used by ICES.

|                        |  |
|------------------------|--|
| <b>Stock</b>           | Herring ( <i>Clupea harengus</i> ) in Subdivisions 30 and 31 (Gulf of Bothnia) |
| <b>Working Group</b>   | Baltic Fisheries Assessment Working Group (WGBFAS)                             |
| <b>Created</b>         | February 2017  |
| <b>Last updated</b>    | WGBFAS, April 2017   |
| <b>Last updated by</b> | Stock Coordinator Jukka Pönni  |

---

### A. General

#### A.1. Stock definition

Management units of Atlantic herring, *Clupea harengus*, have traditionally been based on geographic separation in meristic or morphological characteristics or as a result of stocks with different migration and spawning behaviour (ICES, 2002). The amount of genetic differentiation in Atlantic herring is generally small (Larsson, 2008), and spatial structure and morphological characters have not consistently coincided with genetic differences (Ryman *et al.* 1984; Jørgensen *et al.*, 2008). Although previous studies did not find genetic differences between Atlantic herring (Ryman *et al.*, 1984 and Koskineniemi and Parmanne, 1991), recent genetic research however, supports the need of spatially separate management units due to transoceanic genetic divergence between the Baltic and North Sea herring populations (Bekkevold *et al.*, 2005; Jørgensen *et al.*, 2005) as well as within the Baltic Sea between spatially distinct populations (Bekkevold *et al.* 2005; Jørgensen *et al.*, 2005; 2008; Teacher *et al.*, 2013). Moreover, Jørgensen *et al.* (2005; 2008) found significant associations between genetic differentiation (using microsatellite loci), sea surface temperature and salinity. The variable results that are found depending on the method used to assess spatial and temporal variation of biological characteristics within the current management units (e.g. Rahikainen and Stephenson, 2004) highlight the complexity in the stock structure of herring and calls for future multivariate approaches to stock identification of herring in the Baltic Sea (Waldman, 1999). Recent full genome studies of North Sea and Baltic Sea herring indicate a metapopulation structure with reproductive mixing combined with strong environmental selection (Barrio *et al.*, 2016).

The Gulf of Bothnia area, i.e. subdivisions 30 (Bothnian Sea) and 31(Bothnian Bay) is inhabited by the Gulf of Bothnia herring stock. The stock boundary is set at 60°30'N in the south (till 2016 regarded separately as Bothnian Sea and Bothnian Bay stocks). This area is also the Management Unit III of the International Baltic Sea Fishery Commission (IBSFC), which has included subdivisions 30 and 31 since 2005 (until the end of 2004, SD 29N was also included).

According to tagging studies, there are two spring-spawning coastal herring populations in the Bothnian Sea (Subdivision 30) (Hannerz, 1955; 1956; Otterlind, 1957; 1962; 1976; Sjöblom, 1961; Parmanne and Sjöblom, 1982; 1986), one is distributed along the west coast and the second along the east coast. Similarly, the migration pattern on both

sides of the Bothnian Bay is the same and a mirror image of each other (Aro, 1989). Different spawning groups have been presented in the spring-spawning herring (Ehnholm, 1951).

The autumn-spawning stock is very sparse (Sjöblom, 1966; Sjöblom, 1978). The separation into the coastal and open sea components is not applicable to these populations, although different spawning groups have been proposed (Ehnholm, 1951).

The spring-spawning coastal herring on the Swedish and Finnish coasts spawn in May–July in coastal areas starting in the shallowest areas and shifting to deeper waters following rising water temperatures (Neuman, 1982). Spawning occurs along the entire coastline (Aro, 1989), but some areas are more preferred (e.g. Bergström, 2012). The migration of the adult population from the open sea to the spawning grounds on the Swedish coast takes place in late autumn and early winter (Aro, 1989), which is also indicated by the concentration of large herring in the main spawning areas in autumn (Bergström, 2012). The new spawners, as well as immature fish, overwinter inside the Archipelago Sea and near the coast, and new spawners join the spawning stock during the spawning time. The feeding migration starts soon after spawning. The main feeding areas are the slopes of the Bothnian Sea Basin and the outer Archipelago Sea, and in the north the slopes of the Bothnian Bay Basin and archipelago, as well. The herring in the western side of the Bothnian Sea seem to be more bound to the coast than the southern Baltic stocks (Otterlind, 1976). The feeding migration occurs mainly along the coast and from the Bothnian Sea, there seems to be some migration to the Åland Sea and the Quark (Otterlind, 1957; 1962). The Åland Sea is a transition area with the Central Baltic stock. The proportion of the eastward migration to the Finnish coast is very low.

The spring-spawning coastal herring on the Finnish coast also spawn in May–July along the whole coastline, from the northern side of the Archipelago Sea up to the Quark. Some of the younger mature age groups also overwinter and feed near the coastline and in the Archipelago Sea (Sjöblom, 1961). The migration of the adult population to the spawning grounds from the feeding and overwintering areas in the open sea, the Archipelago Sea and in the Quark Archipelago, occurs during winter, when the first spawning shoals are near the coast. After spawning, the feeding migration to the open sea near the slopes of the Bothnian Sea Basin and to the outer Archipelago Sea occurs quite rapidly. Herring is located in the feeding grounds usually during July–December. The feeding migration extends to southern parts of the Bothnian Sea and inside the Archipelago Sea, the Quark and the Bothnian Bay. There is also some exchange between the Finnish and Swedish coasts (Parmanne and Sjöblom, 1982; 1986). The spring-spawning coastal herring in the east coast of the Bothnian Sea has a clear philipatric behaviour as shown by tagging experiments where about 95% of the recaptures were obtained within 150 km from the tagging place (Parmanne and Sjöblom, 1986).

The biological parameters such as mean weights, annual weight increase, condition and maturity of herring show similarities in the two basins suggesting they are the same stock (WKBALT (2017)). However recruitment patterns and the dynamics in the occurrence of large year classes differed indicating they may be separate stocks. Differences in recruitment and year-class strength could however be due to differences in climate conditions in the two basins, for example SD 31 having longer ice cover periods and harsher winters. Genetics and tagging studies are recommended to further clarify if these stocks are similar. The vast majority of the combined stock in the Gulf of Bothnia inhabits the Bothnian Sea area.

Herring in SDs 30 and 31 have been assessed separately historically, but were combined into one assessment unit at WKBALT (2017). The main arguments for combining these previously separate assessment units:

- i) There is currently no strong biological evidence to conclude on whether to separate or combine SDs 30 and 31 in the stock assessment, based on biology alone.
- ii) Data availability (lack of survey in SD 31) does not support a good quality assessment for SD 31 and this is unlikely to be possible to improve in future.
- iii) There is no concern for overexploitation of the smaller stock component in SD 31 when merged together with a larger component in SD 30. This is because of natural conditions (ice and bottom features: difficulties in trawling) restrict fisheries in SD 31, and there is generally low economic interest in herring fisheries in SD 31.

## A.2. Fishery

Since the mid-1970s the fishery has been dominated by pelagic and deep midwater trawl and trapnet fisheries. In 2016, the Finnish fishery accounted for 83% of the total catch, and trawl fisheries accounted for 96% of the Finnish catches. The trawling vessels in SD 31 are smaller than in SD 30, and this has not changed with time, probably because of the shallow and stony waters of SD 31, where it is difficult or even impossible for large trawlers to operate. The remaining Finnish catches (3%) were taken mainly by trapnets, which target the spawning component of the stock in May–June, and 0,1% were taken with coastal gillnets. The Swedish part of the fishery has been much smaller (e.g. 4% of the total catches in 2011) and of a different gear composition. However, since 2011 the Swedish gear composition has changed and the share of Swedish catches has increased, having been 10% in 2013 and 11% in 2014, 12% in 2015 and 17% in 2016. In 2016, 97% of the Swedish catch came from trawls (83% from pelagic trawling and 14% from deep midwater trawling), 3% from gillnets, and <1% from other passive gears. Catches increased in the late 1990s because of an increased efficiency in the fishery and again since 2012 to record high levels due to stock growth and TAC increase. The fishery is regulated by a TAC.

In the herring fishery, pelagic and deep midwater trawls overlap. Generally the same vessels carry out pelagic and deep midwater trawling and use the same gear for both. The pelagic trawl fleet exploits the younger part of the stock and the deep midwater trawling is more directed towards the adult part of the stock. In autumn and early winter before the sea is covered with ice, pelagic pair trawling is used for industrial purposes. Many pelagic trawling vessels operate between the Bothnian Sea (SD 30) and the Åland Sea and northern Baltic proper (SD 29), depending on fishing possibilities and ice cover during winter. Deep-water trawling takes exclusively Baltic herring, whereas pelagic trawling catches sprat, as well. The trawlers operate in different fishing grounds depending on the fishing possibilities and ice cover.

Sprat occurs in SD 30 in late autumn, winter and early spring but the sprat bycatches in herring fisheries are usually low. Sticklebacks are also caught in small, but growing numbers. Sprat and stickleback in the herring trawl fisheries are used for fodder along with the herring.

In the trapnet fishery for Baltic herring, a variety of trapnet types are used. This fishery is conducted near the coast and inside the archipelago and is mainly targeting the spawning component of Baltic herring stock in spring and early summer (May–June, in the Bothnian Bay also July).

The herring discarding rates in Swedish fisheries was estimated to be 6–12% of Swedish herring catches in 2008–2010. Further analyses on discarding in the Swedish herring fisheries in SDs 30 and 31 are needed to estimate the herring discarding rates in earlier years.

Discarding rates in the Finnish fisheries are negligible (estimated to be a few tons annually) and very small also in the Swedish fisheries and have therefore not been taken into account in assessments.

### SA.3. Ecosystem aspects

The increasing trend in stock size of the Bothnian Sea herring observed since the 1990s has been driven by good recruitment. The biomass dynamics of recruits (age 1) have been linked to increased abundance of *Bosmina* sp. (Lindegren *et al.*, 2011), which is an important food source for both young of the year herring <5 cm in the northern Baltic Proper (Arrhenius, 1996), and for herring in the Bothnian Sea (Flinkman *et al.*, 1992). Recruit abundance models also show that the number of recruits (age 1) is largely explained by *Bosmina* sp. abundance and sea temperature during the main growing season (July–September) for the young-of-the-year, while the best model did not identify an effect of the spawning stock size in recruitment (Gårdmark, Working Document 1 for WKPELA; ICES, 2012).

Among abiotic factors, climate change can be seen in increasing temperatures in the measurements conducted since 1980 (Kuosa *et al.*, 2017). Increasing temperatures have probably increased the production in the ecosystem and improved the feeding conditions of herring larvae. Several especially abundant year classes of herring have developed in very warm years, which supports the effect of temperature. In addition to increase in temperature, the contents of dissolved inorganic phosphorus and dissolved inorganic nitrogen (winter values) have increased since 1980 (Kuosa *et al.*, 2017), both possibly having additional affects besides the increase in temperature, regardless of the reason to increased nutrient levels. The increase of especially phosphorus is similar to the trends observed in herring abundance.

The Gulf of Bothnia herring is the prey of several predators, including cod (*Gadus morhua*) at least periodically in the southern areas, salmon (*Salmo salar*), coastal piscivores (pike, perch, pike-perch), grey seals, ringed seals, and birds such as cormorants (*Phalacrocorax carbo*). Grey seals can be regarded as the main predator for several reasons: cod and salmon are less abundant and herring makes up to about 70% of the biomass in the diet of an average grey seal (Lundström *et al.*, 2010; Gårdmark *et al.*, 2012). The proportion of herring in the food of cormorants is below 35% (J. Salmi, pers.comm.), and there are only a small number of ringed seals permanently in the Bothnian Sea (Anonymous, 2007).

Grey seals predate selectively on larger individuals of herring, and mean length of prey was about 18 cm (Gårdmark *et al.*, 2012). While grey seals have not been found to be a major driver of herring population dynamics (Lindegren *et al.*, 2011), Östman *et al.* (2014) found that the size-selective removal of herring by grey seals may have decreased average weight-at-age of herring cohorts over the latest decades. The variation in length-specific body growth of cohorts was also explained by zooplankton resources (*Eurytemora* sp.), intraspecific density, and cohort-specific fishing mortality (Östman *et al.*, 2014). In the most recent years, a gradual increase in mean weights-at-age has been observed.

On the basis of the acoustic surveys, the total mortality of herring is low in the Gulf of Bothnia stock. Thus, the natural mortality was found to be low as well, and the mortality by grey seal predation has not been separately included in the current stock assessment.

## **B. Data**

### **B.1. Commercial catch**

Finnish commercial herring catch statistics is based on catch notifications submitted by fishermen at set intervals. The application of the Act (1139/94) on implementing the Common Fisheries Policy of the European Union obliges all commercial fishermen to submit catch notifications.

The discards are negligible in both countries' commercial fisheries and therefore not sampled either. Also, the information of discards from fishermen's reports are not used in assessment.

The fishing data of vessels  $\geq 10$  metres long are entered in the EU fishing logbook. The data entered are the dates of fishing by fishing trip, the size of the catch by species, the fishing (statistical) rectangle, the gear and number of gears used in fishing, and the trawling time in hours. A fisherman is obliged to keep an up-to-date logbook on board his vessel. The logbook must be returned to the regional authorities within 48 hours of the catch being landed.

With the exception of salmon catches, the Finnish fishing data of vessels  $\leq 10$  metres long are entered monthly in a coastal fishery form. The data entered are the size of the catch by species by the statistical rectangle, the type and number of gears used in fishing, and the number of fishing days. The forms must be returned to the regional authorities by the fifth day of the following month. All logbooks and most of the other catch notification forms are checked by national authorities.

The proportion of the Baltic herring catch, landed in Finland for the food and processing industry in relation to the total catch of that species, is estimated with the aid of the fish purchasing register that is maintained by the Ministry of Agriculture and Forestry.

Because all the main fisheries (pelagic trawling, deep midwater trawling and trapnets) have different exploitation patterns, their catches are also sampled separately. The sampling in the Gulf of Bothnia herring fishery is performed according to EU DCF requirements, covering 12 strata (three fleets and four year-quarters).

Since the study projects funded by DG XIV (International Baltic Sea Sampling Programs I & II) in 1998–2001, a length stratified subsampling scheme has been applied to estimate age compositions of Finnish catches of Baltic herring. This sampling scheme is designed to be compatible with international databases and uses standardized methodologies in data processing. Baltic herring samples are collected mainly in fishing harbours and, if necessary, also on board commercial fishing vessels. In the sampling scheme the annual life cycle of Baltic herring and the presence of the ice coverage during the winter in the Gulf of Bothnia have been taken into account. Because of icing conditions, the three fishing gears are not in use year-round (e.g. trapnet fishery is usually conducted only in spawning time during quarters 2 and 3). The sampling effort is roughly based on the proportions of catches in different fisheries. Moreover, the sampling intensity in general is locally adjusted during the year according to temporal and regional changes in fisheries. The seasonal herring fishing intensity is predominantly

dependent on the TAC, which may cause fishing restrictions in certain fisheries and/or seasons and may therefore change the sampling intensity from the original plan. A minimum coverage target is at least one sample by fishery per month (or three samples by fishery per year-quarter). The sampling strategy is to have age-length samples from all major gears in each quarter.

The Finnish and Swedish input files are uploaded to ICES InterCatch database. The data can also be found in the national laboratories and with the stock co-ordinator. The national data have been aggregated to international data in InterCatch.

**Table 1. Description of the types of data available per country.**

| Country | KIND OF DATA            |                                 |                                   |                                    |                             |
|---------|-------------------------|---------------------------------|-----------------------------------|------------------------------------|-----------------------------|
|         | Caton (catch in weight) | Canum (catch-at-age in numbers) | Weca (weight-at-age in the catch) | Matprop (proportion mature by age) | Length composition in catch |
| Finland | x                       | x                               | x                                 | x                                  | x                           |
| Sweden  | x                       | x                               | x                                 |                                    | x                           |

## B.2. Biological sampling

The age and the individual weight data are obtained from both Finnish and Swedish landings from all year quarters, as well as from the catch samples in acoustic surveys in 3rd and 4th quarter. The annual weights-at-age are weighted by the year-quarterly catch-numbers for WECA. The maturity ogives are based on the proportions of mature individuals of each age group before spawning time in the Finnish sampling data, and are updated every year.

### B.2.1 Calculation of catch-at-age

In Finland the calculation of catch-at-age is based on year-quarterly performed length-stratified random sampling of individual fish (at minimum 10 aged individual fish from all prevailing 0,5 cm length classes) and length samples of at least 300 specimens per sample from different commercial fisheries per quarter. The average number of individual samples is 1101 from commercial fisheries and 2473 from surveys in SD 30 and 587 from commercial fisheries in SD 31 annually, and the average number of length measurements is 18 537 and 5987 respectively.

The quarterly collected length distributions (from length sampling) are converted into age distributions with year quarterly prepared age-length keys, ALKs, which are derived from the sampling of individuals.

The quarterly catches from the main herring fisheries (OTM + PTM carried out in mid-water and deep midwater and trapnets, FPN + FYK) are divided by the mean weight of the herring from length samples of respective fisheries in order to get the total catch number of fish for all strata (all fisheries, four quarters). The total catch numbers from each fishery and year quarter are then multiplied by the proportions of the age classes in the age distributions and summed up to get the annual catch-at-age.

In Sweden the length samples of at least 300 specimens per sample from two (main) commercial fisheries [bottom trawls (XTB) and gillnets (GNS)] in SD 30 and only from gillnets in SD 31 are collected quarterly each year. The catches of pelagic trawl (OTM and PTM) fisheries are not sampled. Length-stratified random sampling of individual

fish (approximately 20 aged individual fish from all prevailing 0,5 cm length classes per quarter) is performed only for gillnet fisheries. In SD 30 the average total number of annual length measurements is 5600 from bottom trawls and 2300 from gillnet fisheries, and the average total number of sampled fish individuals is 490, and in SD 31 the average total number of annual length measurements is 1700, and the average number of sampled fish individuals is 450.

The length distributions (from length sampling) are converted into age distributions with quarterly prepared age-length keys (ALKs). For that purpose, additionally Finnish ALK and mean weight-at-length data from trawl fisheries are borrowed.

The calculation of total annual catch-at-age follows the same procedure as in Finland.

### **B.2.2 Calculation of mean weight**

The mean weights-at-age are derived from the individual data collected from commercial catches all year-round as well as from the individual data of acoustic survey trawl samples during September–October (2600 individuals annually), and averaged over year quarters. The annual mean weights-at-age for assessment are derived by weighting the year-quarterly mean weights by the year-quarterly catch numbers.

### **B.2.3 Maturity**

The maturities are defined from the individual data that is collected all the year round from commercial catches with other so called “stock related variables” as length, weight and age, and from the trawl samples of the acoustic survey. The data for the maturity ogive used in assessments are collected from samples before spawning (i.e. January to March in SD 30 and March to May in SD 31) because the idea is to get the proportion of spawners by age from the whole population, before the spawning part separates itself from non-spawners by approaching the coastline to spawning areas.

The share of mature fish in each age group is calculated from annual data and the annual number of the individual samples for maturity definitions that are used for the maturity ogives has been on average (2010–2015) 283 in SD 30 and 212 in SD 31.

The maturity scale (Table 6.2) in use is the modified European standard 9-stage scale and the same scale is used both in Finland and Sweden. The stages II–VIII (VIII-A and VIII-B) are considered mature, while stage I and IX are counted as “non-mature” although stage nine (abnormal) is usually mature, but not accounted to take part to spawning.

The maturities defined during a Swedish acoustic survey in 4th quarter and the maturities derived from Finnish 1st quarter sampling of commercial catches have showed very small differences.

In WKPELA in 2012 benchmark the sensitivity of the annual changing proportions of spawners in age groups was tested (by several types of averages over time<sup>1)</sup> and although there are clearly visible annual changes in mostly 2-year-olds, there was only negligible impact to e.g. estimates of SSB. It was concluded then that it was still better to have the latest real information on maturity-at-age than assume something else.

<sup>1)</sup> Four new combinations of maturity ogives were introduced to XSA (maturity ogive with 3- and 5 year running averages for the whole time-series, constant maturity ogive for the whole time-series as an average of the whole time-series and two different averages over the time-series according to periods before and after the alleged regime shift (1973–1988 and 1989–2010)). Resulting estimates of SSB were compared to the annually updated maturity ogive in SPALY run, and the differences were found to be negligible with the exception of year 2010 only.

The reason for the “instability” was found to be the high interannual variation in the maturation of 2-year olds in the whole time-series and especially in 2010. The maturity calculations from raw data were examined carefully, and no mistakes were revealed.

Table 2. Maturity scale in use in Finland and Sweden.

| Std Eur. Scale | Maturity stage (code) | Maturity stage                |
|----------------|-----------------------|-------------------------------|
| Ia             | I                     | Immature, juvenile            |
| Ib             | II                    | Immature, early development   |
| IIa            | III                   | Maturing, early stage         |
| IIb            | IV                    | Maturing, late stage          |
| IIIa           | V                     | Spawning, prepared            |
| IIIb           | VI                    | Spawning, running             |
| IV             | VII                   | Spent                         |
| Va             | VIII A                | Regeneration, regressing      |
| Vb             | VIII B                | Regenerating/skipped spawning |
| VI             | IX                    | Abnormal                      |

From 2002 to 2006, Finnish samples were age determined with two different methods in parallel, using traditionally whole otoliths and as a new method, neutral red stained slices of cut otoliths. The effects of the age-determination method were presented at the WGBFAS meeting in 2006 (Raitaniemi and Pönä, 2006). The method affects the age distribution as well as the proportion of mature fish at-age. Especially in old age groups (from age 5 or 6 on), determination from cut otoliths generally results in an older assessed age compared to whole otoliths. In the comparison, the numbers-at-age in the total catch differed about 2% on average, but ranged from 0.4% to 52% depending on year and age. On average the proportion of four- to eight-year-olds in the catch was 11% lower, and the proportion of ages 9+ was 32% higher, when using neutral red stained slices compared to whole otoliths.

According to Peltonen *et al.* (2002), the agreement between the determinations of different age readers was better with the cut otoliths technique than with whole otoliths. A combination of age data from Finnish cut otoliths (representing 98% of the catch) and Swedish whole otoliths (representing 2% of the catch) was used between 2002–2006. The slicing method was calibrated between Finland and Sweden in 2007, and it has been applied also to Swedish catches as well as Bothnian Sea surveys since 2007. Since age determination using cut otoliths is considered to be more accurate (Raitaniemi and Pönä, 2006), this method is used as the standard method for ageing all the samples, and the time-series including ages from whole otoliths from 1980–2001 and cut otoliths from 2002 onwards is used in the assessments of this stock.

### B.3. Surveys

Annual hydroacoustic surveys have been conducted in SD 30 in October from 2007 until 2010 with Swedish RV Argos. In 2011 and in 2012 the survey was performed with



the Danish RV Dana and since 2013 with Finnish RV Aranda. This survey is coordinated by ICES within the Baltic International Acoustic Surveys (BIAS). The annual survey indices are collected and calculated with standardised methods within the international coordination of ICES WGBIFS and stored in international databases. The actual calculations have been performed in years 2007–2012 in the Swedish marine research institute (Havsfiskelaboratoriet) by Niklas Larson and from 2013 onwards in the Natural Resources Institute Finland by Juha Lilja.

The acoustic survey has been considered as a reliable tuning fleet and has been included into the SD 30 assessment in 2013 after an independent review process (ICES, 2015a).

The acoustic estimates are used as abundance indices (tuning fleet) for the assessment. In the acoustic tuning fleet, age groups 1–14 (true ages) are applied (compared with former acoustic fleet the age group one was added, since it showed similar consistency between ages as older ages (Figure 1)).




Figure 1. Consistency between consecutive age classes in acoustic tuning fleet.

The coverage of the acoustic transects and trawl samples has mostly been good (Figure 2). In 2012 the coverage was only half of the “normal” because of a sudden 50% reduction in funding. In 2014 there were problems with the fishing gear, which reduced the trawl hauls, but the spatial acoustic coverage was not affected that much. In 2015 a storm damaged the ship so that the most northern part of the area had to be skipped due to lack of time after fixing the damage in harbour.

The 2012 50% reduction in the survey effort, as well as the 2014 and 2015 results were, however, considered acceptable for the index by the survey expert working group, ICES WGBIFS (ICES, 2013; ICES, 2015; ICES, 2016).

The survey is based on IBAS (International Baltic Acoustic Survey) manual (ICES, 2016) with the aim of 60 Nm of acoustic transect and two trawl hauls per statistical rectangle. In the catch sampling length distributions at least 300 fish are measured in 0,5 cm length classes and ten individuals from all prevailing length class are aged per rectangle, comprising normally of about 20 000 length measurements and 2600 age readings annually.




Figure 2. Spatial coverage of acoustic surveys conducted in Bothnian Sea in years 2007–2015.

#### B.4. Commercial cpue

Trapnet fishing that takes place during the spawning period is used as a tuning fleet in the stock assessment. The main commercial fleets, i.e. pelagic trawling (single- and pair-trawling) and deep midwater trawling are not used.

The reason for not using commercial trawl fishery fleets in the stock assessment is the change in trawls. There has been an increase in trawl gear size and changes in gear characteristics since the early 1980s. In the past, reported fishing effort data (trawling hours) may not have been accurate.

The trapnet tuning fleet was revised in 2012. Before 2012, the total number of trapnets set in SD 30 was used as effort for the herring catch caught. Throughout the 1980s, the total number of trapnets decreased, but since 1990s it has remained more or less stable. Due to reduction in effort in years 1974–1989, the revised dataset starts at 1990.

The revised dataset is considered geographically representative and only includes trapnets that are used for catching herring in spawning time (April–June). The 1990–2006 dataset used in the assessment includes only those areas (statistical rectangles) that have unbroken time-series for the catches and the corresponding effort (number of herring trapnets). The cpue taking into account the difference in catchability between sites is used as an index for stock size (numbers-at-age) in the assessment.

In 2015, the trapnet index was, however, not considered sufficiently reliable by the working group, as it is introducing increasing uncertainties to the assessment.

After a review process (ICES, 2015a) it was decided that the trapnet tuning series should be truncated and the last years of data (2007 onwards; overlap period between the acoustic and the trapnet tuning series) were not included in the assessment (ICES, 2015a). In the benchmark of 2017 and assessment of 2017, this practice was continued, excluding trapnet data from the years 2007 onwards in the tuning.

The trapnet abundance indices standardization model was changed from previously used GLM to the multilayer perceptron (MLP) algorithm (WD1). The statistical model performances of MLP and GLM models were roughly comparable. However, in very high abundance years the MLP-based model fit against observed data was better than that of GLM. Hence, the MLP model was updated with additional age groups (3–14, previously 3–9). The statistical performance of the MLP model with age groups 3–14 was better than using age groups 3–9. Therefore (and altogether), the working group used MLP-based cpue estimates of age groups 3–14 in 1990–2006 in stock assessment of the combined Gulf of Bothnia stock.

### C. Assessment: data and method

Model used: stochastic state–space model SAM (ICES, 2009).

The model was run using web interface that can be viewed at <https://www.stockassessment.org>.

Details concerning input data and model configuration can also be found on this webpage (username: guest, password: guest).

The model configurations are as follows:

**# Min Age (should not be modified unless data are modified accordingly)**

1

**# Max Age (should not be modified unless data are modified accordingly)**

10

**# Max Age considered a plus group (0=No, 1=Yes)**

1

**# The following matrix describes the coupling**

**# of fishing mortality STATES**

**# Rows represent fleets.**

**# Columns represent ages.**

|   |   |   |   |   |   |   |   |   |   |
|---|---|---|---|---|---|---|---|---|---|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 9 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

**# Use correlated random walks for the fishing mortalities**

**# ( 0 = independent, 1 = correlation estimated)**

0

**# Coupling of catchability PARAMETERS**

```

0 0 0 0 0 0 0 0 0 0
1 2 2 2 2 2 2 2 2 0
0 0 3 3 3 3 3 3 3 0
# Coupling of power law model EXPONENTS (if used)
0 0 0 0 0 0 0 0 0 0
0 0 0 0 0 0 0 0 0 0
0 0 0 0 0 0 0 0 0 0
# Coupling of fishing mortality RW VARIANCES
1 1 2 2 2 2 2 2 2 2
0 0 0 0 0 0 0 0 0 0
0 0 0 0 0 0 0 0 0 0
# Coupling of log N RW VARIANCES
1 2 3 3 3 3 3 3 3 3
# Coupling of OBSERVATION VARIANCES
1 1 1 1 1 1 1 1 2 2
3 4 4 4 4 5 5 5 5 0
0 0 6 6 6 7 7 7 7 0
# Stock-recruitment model code (0=RW, 1=Ricker, 2=BH, ... more in time)
0
# Years in which catch data are to be scaled by an estimated parameter
0
# first the number of years
# Then the actual years
# Then the model config lines years cols ages
# Define  $F_{bar}$  range
3 7

```

## Input data types and characteristics:

| TYPE    | NAME  | YEAR RANGE            | AGE RANGE | VARIABLE FROM YEAR TO YEAR<br>Yes/No                       |
|---------|---|-----------------------|-----------|--|
| Caton   | Catch in tonnes                                 | 1980–last data year   |           | Yes  |
| Canum   | Catch-at-age in numbers                         | 1980 – last data year | 1–10+     | Yes  |
| Weca    | Weight-at-age in the commercial catch           | 1980–last data year   | 1–10+     | Yes  |
| West    | Weight-at-age in the stock                      | 1980–last data year   | 1–10+     | Yes - assumed to be the same as weight-at-age in the catch |
| Mprop   | Proportion of natural mortality before spawning | 1980–last data year   | 1–10+     | No - set to 0.33 for all ages in all years                 |
| Fprop   | Proportion of fishing mortality before spawning | 1980–last data year   | 1–10+     | No - set to 0.15 for all ages in all years                 |
| Matprop | Proportion mature at-age                        | 1980–last data year   | 1–10+     | Yes  |
| Natmor  | Natural mortality                               | 1980–last data year   | 1–10+     | No - set to 0.15 for all ages in all years                 |

## Tuning data:

| TYPE           | NAME                  | YEAR RANGE          | AGE RANGE |
|----------------|-----------------------|---------------------|-----------|
| Tuning fleet 1 | Finnish trapnet fleet | 1990–2006           | 3–9       |
| Tuning fleet 2 | Acoustic              | 2007–last data year | 1–9       |

**D. Short-term projection**

The model has been run using web interface that can be viewed at <https://www.stock-assessment.org>. (username: guest, password: guest)

The short-term forecast is based on the SAM short-term forecast module. The short-term prediction carried out by the SAM model is simulation based, and accounts for uncertainty in the final year estimates. From the assessment model it takes the final estimates of fishing mortality and stock numbers, and their estimation variances and covariances. These quantities are then simulated forward in time for a number of specified scenarios (e.g. scaling of fishing mortality levels). The uncertainties are propagated forward in time, and the process variation (as estimated from the historic period) is added. Medians are used as main summary statistic. It is important to note that taking uncertainty into account does not merely supply confidence intervals on the final future catch estimates, but can also affect the estimates themselves as the errors accumulate in the non-linear projections.

The settings for the short-term projection are as follows:

Initial stock size: Final year estimates from SAM.

Maturity: Assumed to be equal to the average of maturity ogives across the last three years

F and M before spawning: The proportion of total annual natural mortality before spawning is assumed to be 33% and proportion of F before spawning 15% of the annual fishing mortality. Natural mortality is set to 0.15.

Weight-at-age in the stock and the catch: These are assumed to be equal to the average of mean weights-at-age across the last three years

Exploitation pattern: The average selectivity pattern, scaled to F in the last three years

Stock–recruitment model used: Recruitment are based on resampling from the observed distribution in all years

### **G. Biological reference points**

The two stocks were merged and thus, reference points were re-estimated at WKBALT (2017). The Eqsim based reference point analysis used the newest (1980–2015) assessment results from the SAM assessment (WKBALT 2017). The stock–recruitment fit using the three models (Ricker, B&H and segmented regression) weighted by the default "Buckland" method available in EqSim gave a "straight" line for all models (Figure 2).






Figure 2. The stock–recruitment fit using the three models (Beverton–Holt, Ricker and segmented regression) weighted by the default "Buckland" method gave a "straight" line for all models.

Thus, a segmented regression model was used with a breakpoint set arbitrarily at the average observed SSB (i.e.  $B_{lim} = 405\,980$  t) as dictated by ICES guidelines for reference point estimation. However, this will result in a rather large value of  $B_{trigger}$  and  $B_{pa}$ . Thus, the ICES reference points' guidance was modified in this case. The first step was to estimate  $F_{MSY}$  using a hockey-stick S–R relationship with  $B_{lim}$  at the average SSB and without  $MSY$   $B_{trigger}$ , but with assessment and advice error (i.e. using the default values). Once the  $F_{MSY}$  was estimated, the simulations were run again with the same hockey-stick S–R relationship and  $B_{lim}$  to estimate  $MSY$   $B_{trigger}$  defined as the 5th percentile of the SSB at  $F_{MSY}$ . Successively,  $B_{pa}$  was set as  $MSY$   $B_{trigger}$  and a new value of  $B_{lim}$  was estimated as  $B_{pa}$  divided by  $\exp(1.645 \times 0.2)$ . After  $B_{lim}$ ,  $B_{pa}$  and  $MSY$   $B_{trigger}$  were all defined, the ICES procedure for setting the reference points was used to estimate the remaining reference points. The S–R relationship used for these runs was a hockey stick with the breakpoint set at the new  $B_{lim}$ . The number of samples used to fit the S–R relationship and the number of runs used in all EqSim simulations were 1000 and 200, respectively. Autocorrelation of recruitment was used in all EqSim simulations.  $F_{pa}$  was estimated using the ICES standard procedure ( $F_{pa} = F_{lim} \times \exp(-1.645 \times \sigma)$ ). Sigma was estimated as the uncertainty associated to the  $F$  in last assessment year (i.e. 2015;  $\sigma = 0.223$ ).

The proposed summary table of the combined Gulf of Bothnia stock reference points is:

| STOCK   |           |
|---|-----------|
| Reference point   | Value     |
| F <sub>P,05</sub> (5% risk to B <sub>lim</sub> ) with MSY B <sub>trigger</sub>    | 0.21      |
| F <sub>P,05</sub> (5% risk to B <sub>lim</sub> ) without MSY B <sub>trigger</sub> | 0.180     |
| F <sub>MSY</sub>  | 0.21      |
| F <sub>MSY</sub> lower  | 0.151     |
| F <sub>MSY</sub> upper  | 0.25      |
| F <sub>pa</sub>   | 0.20      |
| F <sub>lim</sub>  | 0.29      |
| F <sub>MSY</sub> upper precautionary  | 0.20      |
| F <sub>MSY</sub> range with MSY B <sub>trigger</sub>                              | 0.15-0.21 |
| F <sub>MSY</sub> range without MSY B <sub>trigger</sub>                           | 0.15-0.18 |
| MSY B <sub>trigger</sub>  | 283180 t  |
| B <sub>pa</sub>   | 283180 t  |
| B <sub>lim</sub>  | 202272 t  |

#### H. Other issues

Concerning reference points, it was brought up that in the process of combining the stocks, the time-series from SD 30 was unfortunately truncated from 1973 to 2015 to match the time-series of SD 31 that was only years 1980 to 2015. It was further suggested that in future Benchmark efforts should be made to fill in the missing data by some assumptions or using models that can handle more fragmented type of data.

## I. References

- Anonymous. 2007. Management Plan for the Finnish Seal Populations in the Baltic Sea. Finnish Ministry for Agriculture and Forestry, 4b/2007. 96 pp.
- Aro, E. 1989. A review of fish migration patterns in the Baltic. Rapp. P.-v. Réun. Cons. Int. Explor. Mer., 190: 72–96.
- Arrhenius, F. 1996. Diet composition and food selectivity of 0-group herring (*Clupea harengus* L.) and sprat (*Sprattus sprattus* (L.)) in the northern Baltic Sea. ICES Journal of Marine Science, 53: 701–712.
- Barrio *et al.* 2016. <http://dx.doi.org/10.7554/eLife.12081>
- Bekkevold, D, Andre, C, Dahlgren, T.G., Clausen, L.A.W., Torstensen, E., Mosegaard, H. *et al.* 2005. Environmental correlates of population differentiation in Atlantic herring. *Evolution*, 59: 2656–2668.
- Bergström, U. 2012. Information on herring reproduction areas and important fishing areas from PLAN BOTHNIA. Working document for WKPELA 2012.
- Ehnholm, G. 1951. Studier over strömming I Östra Kvarken. Havsforskningsinstitutets skrift., No. 149: 1–94.
- Flinkman, J., I. Vuorinen, and E. Aro. 1992. Planktivorous Baltic herring (*Clupea harengus*) prey selectively on reproducing copepods and cladocerans. *Canadian Journal of Fisheries and Aquatic Sciences*, 49: 73–77.
- Gårdmark, A., Östman, Ö., Nielsen, A., Lundström, K., Karlsson, O., Pönni, J., and Aho, T. In preparation. Does predation by grey seals (*Halichoerus grypus*) affect Bothnian Sea herring stock estimates? In revision for ICES Journal of Marine Science.
- Hannerz, L. 1955. Strömmingsmärkningen i Bottenhavet 1954. *Ostkusten*, 1: 14–15.
- Hannerz, L. 1956. Preliminary results of the herring investigations in the Bothnian Sea 1954. *Annls. biol.*, Copenhagen, 11: 156–158.
- Hiby, L., Lundberg, T., Karlsson, O., Watkins, J., Jüssi, M., Jüssi, J., and Helander, B. 2007. Estimates of the size of the Baltic grey seal population based on photo-identification data. *NAM-MCO Scientific Publications*. 6: 163–175.
- ICES. 2001. Report of the study group of herring assessment units of the Baltic. ICES ACFM/2001:10.
- ICES. 2002. Report of the study group of herring assessment units of the Baltic. ICES CM 2002/H:04.
- ICES. 2009. Report of the Working Group on Methods of Fish Stock Assessment (WGMG), 20–29 October 2009, Nantes, France. ICES CM 2009/RMC:12. 85 pp.
- ICES. 2012. Report of the Benchmark Workshop on Pelagic Stocks (WKPELA 2012), 13–17 February 2012, Copenhagen, Denmark. ICES CM 2012/ACOM:47. 525 pp.
- ICES. 2013. Report of the Inter-Benchmark Protocol for Herring in Subdivision 30 (IBP Her30), 11–15 March 2013, by correspondence. ICES CM 2013/ACOM:60. 94 pp.
- ICES. 2013b. Report of the Baltic International Fish Survey Working Group (WGBIFS), 21–25 March 2013, Tartu, Estonia. ICES CM 2013/SSGESST:
- ICES. 2015a. Report of the Baltic Fisheries Assessment Working Group (WGBFAS), 14–21 April 2015, ICES Headquarters, Copenhagen, Denmark. ICES CM 2015/ACOM:10.
- ICES. 2015b. EU request to ICES to provide FMSY ranges for selected North Sea and Baltic Sea stocks. ICES Special Request Advice.
- ICES. 2015c. Report of the Baltic International Fish Survey Working Group (WGBIFS), 23–27 March 2015, Öregrund, Sweden. ICES CM 2015/SSGESST:

- ICES. 2016. Report of the Baltic International Fish Survey Working Group (WGBIFS), 30 March–4 April 2016, Rostock, Germany. ICES CM 2016/SSGIEOM:07. 121 pp.
- ICES. 2017. Report of the Benchmark Workshop on Baltic Stocks (WKBALT), 7–10 February 2017, Copenhagen, Denmark. ICES CM 2017/ACOM:30. 108 pp.
- Jørgensen, H.B.H., Hansen, M.M., Bekkevold, D., Ruzzante, D.E., and Loeschcke, V. 2005. Marine landscapes and population genetic structure of herring (*Clupea harengus* L.) in the Baltic Sea. *Mol Ecol*, 14: 3219–3234.
- Jørgensen, H.B.H., Pertoldi, C., Hansen, M.M., Ruzzante, D.E., Loeschcke, V. 2008. Genetic and environmental correlates of morphological variation in a marine fish: the case of Baltic Sea herring (*Clupea harengus*). *Can J Fish Aquat Sci*, 65: 389–400.
- Koskineniemi, J and Parmanne. 1991. The genetic population structure of the Baltic herring in the northern Baltic Sea. ICES C.M. 1991/J:33, 18 pp.
- Kuosa, H., Fleming-Lehtinen, V., Lehtinen, S., Lehtiniemi, M., Nygård, H., Raateoja, M., Raitaniemi, J., Tuimala, J. and Uusitalo, L. 2017. A retrospective view of the development of the Gulf of Bothnia ecosystem. *Journal of Marine Systems* 167: 78–72.
- Larsson, L.C. 2008. Disentangling small genetic differences in large Atlantic herring populations: comparing genetic markers and statistical power. Doctoral dissertation. Stockholm: Department of Zoology, Stockholm University, 2008. ISBN 978-91-7155-780-3.
- Lindgren, M., Östman, Ö., Gårdmark, A. 2011. Interacting bottom-up and top-down effects shape the population dynamics of herring. *Ecology*, 92(7): 1407–1413.
- Lundström, K., Hjerne, O., Lunneryd, S.-G., and Karlsson, O. 2010. Understanding the diet composition of marine mammals: grey seals (*Halichoerus grypus*) in the Baltic Sea. *ICES Journal of Marine Science*, 67: 1230–1239.
- Neuman, E. 1982. Species composition and seasonal migrations of the coastal fish fauna in the southern Bothnian Sea. In *Coastal Research in the Gulf of Bothnia*. pp. 317–351. (Eds. by K. Müller). Dr W. Junk Publishers, The Hague-Boston-London 1982.
- Otterlind, G. 1957. Från strömmingsmärkningarna på ostkusten. *Ostkusten*, 3: 24–30.
- Otterlind, G. 1962. Sillens/strömmingens vandringsvanor vid svenska syd- och ostkusten. *Ostkusten*, 1: 15–21.
- Otterlind, G. 1976. Fish stocks and fish migration in the Baltic Sea environment. *Ambio Spec. Rep.*, 4: 89–101.
- Parmanne R. 1988. Herring growth in the northern Baltic Sea in 1974–1987. ICES C.M. 1988/J:19. (Mimeograph.)
- Parmanne R. 1990. Growth, morphological variation and migrations of herring (*Clupea harengus*) in the northern Baltic Sea. *Finnish Fish. Res.*, 10, 48 p.
- Parmanne, R., and Sjöblom, V. 1982. Recaptures of Baltic herring tagged off the coast of Finland in 1975–1981. ICES C.M. 1982/J:19, 12 pp.
- Parmanne, R., and Sjöblom, V. 1986. Recaptures of Baltic herring tagged off the coast of Finland in 1982–1985. ICES C.M. 1986/J:28, 12 pp.
- Peltonen, H., Raitaniemi, J., Parmanne, R., Eklund, J., Nyberg, K., and Halling, F. 2002. Age determination of Baltic herring from whole otoliths and from neutral red stained otolith cross sections. *ICES Journal of Marine Science*, 59: 323–332.
- Rahikainen, M. and Kuikka, S. 2002. Fleet dynamics of herring trawlers—change in gear size and implications for interpretation of catch per unit effort. *Can. J. Fish. Aquat. Sci.*, 59: 531–541.
- Rahikainen, M., and Stephenson, R. L. 2004. Consequences of growth variation in northern Baltic herring for assessment and management. *ICES Journal of Marine Science*, 61:338–350.

- Ryman, N., Lagercrantz, U., Andersson, L., Chakraborty, R., and Rosenberg R. 1984. Lack of correspondence between genetic and morphologic variability pattern in Atlantic herring (*Clupea harengus*). *Heredity*, 53 (3): 687–704.
- Sjöblom, V. 1961. Wanderungen des Strömlings (*Clupea harengus* L.) in einigen Schären- und Hochseegebieten der nördlichen Ostsee. *Ann. Zool. Soc. Zool. Bot. Fenn. Vanamo*, 23(1): 1–193.
- Sjöblom, V. 1966. Rannikko syyskutuinen silakka vuonna 1964 ja sen runsauteen vaikuttavista tekijöistä. (Summary: Autumn spawning Baltic herring (*Clupea harengus* L.) on the Finnish coast in 1964 and factors affecting fluctuations in its abundance). *Suomen Kalatalous* 25: 1–23.
- Sjöblom, V. 1978. The effect of climatic variations on fishing and fish populations. *Fennia*, 150: 33–37.
- Teacher, A.G.F., Andre, C., Jonsson, P.R. and Merilä, J. 2013. Oceanographic connectivity and environmental correlates of genetic structuring in Atlantic herring in the Baltic Sea. *Evolutionary Applications* 6: 549–567.
- Östman, Ö., Karlsson, O., Pönni, J., Kaljuste, O., Aho, T. and Gårdmark, A. 2014. Relative contributions of evolutionary and ecological dynamics to body size and life-history changes of herring (*Clupea harengus*) in the Bothnian Sea. *Evolutionary Ecology Research*, 2014, 16: 417–433.
- Authors of this Stock Annex: Pekka Joutela, Zeynep Pekcan, Jukka Pönni, Jari Raitaniemi on behalf of WGBFAS 2017.