


New Trends in Important Diseases Affecting the Culture of Fish and Molluscs in the ICES Area 2002- 2015

ICES COOPERATIVE RESEARCH REPORT

RAPPORT
DES RECHERCHES
COLLECTIVES



New Trends in Important Diseases Affecting the Culture of Fish and Molluscs in the ICES Area 2002 – 2015

Editors

Neil Ruane • Ryan Carnegie



ICES
CIEM

International Council for
the Exploration of the Sea

Conseil International pour
l'Exploration de la Mer

International Council for the Exploration of the Sea Conseil International pour l'Exploration de la Mer

H. C. Andersens Boulevard 44–46
DK-1553 Copenhagen V
Denmark
Telephone (+45) 33 38 67 00
Telefax (+45) 33 93 42 15
www.ices.dk
info@ices.dk

Recommended format for purposes of citation:

Alfjorden, A., Areskog, M., Bruno, D., Carnegie, R., Cheslett, D., Feist, S., Ford, S., *et al.* 2017. New Trends in Important Diseases Affecting the Culture of Fish and Molluscs in the ICES Area 2002 – 2015. ICES Cooperative Research Report No. 337. 50 pp.
<https://doi.org/10.17895/ices.pub.2800>

Series Editor: Emory D. Anderson

The material in this report may be reused for non-commercial purposes using the recommended citation. ICES may only grant usage rights of information, data, images, graphs, etc. of which it has ownership. For other third-party material cited in this report, you must contact the original copyright holder for permission. For citation of datasets or use of data to be included in other databases, please refer to the latest ICES data policy on the ICES website. All extracts must be acknowledged. For other reproduction requests please contact the General Secretary.

This document is a report of an Expert Group under the auspices of the International Council for the Exploration of the Sea and does not necessarily represent the view of the Council.

Cover image: © OCEANA/Carlos Suárez

ISBN 978-87-7482-201-1

ISSN 2707-7144

© 2016 International Council for the Exploration of the Sea

Contents

1	Background	1
2	Viral Diseases of Farmed Fish	2
2.1	Infectious salmon anaemia	2
2.1.1	Description of Agent	2
2.1.2	Geographical Distribution and Temporal Trends	2
2.1.3	Short Description of Clinical Signs	2
2.1.4	Control/Preventative Measures	2
2.1.5	Other Host Species	3
2.2	Viral Haemorrhagic Septicaemia	3
2.2.1	Description of Agent	3
2.2.2	Geographic Distribution and Temporal Trends	3
2.2.3	Short Description of Clinical Signs	3
2.2.4	Control/Preventative Measures	4
2.2.5	Other Host Species	4
2.3	Pancreas Disease	4
2.3.1	Description of Agent	4
2.3.2	Geographical Distribution and Temporal Trends	4
2.3.3	Short Description of Clinical Signs	5
2.3.4	Control/preventative measures	5
2.3.5	Other Host Species	5
2.4	Infectious Pancreatic Necrosis	5
2.4.1	Description of Agent	5
2.4.2	Geographical Distribution and Temporal Trends	5
2.4.3	Short Description of Clinical Signs	6
2.4.4	Control/Preventative Measures	6
2.4.5	Other Host Species	6
2.5	Viral Nervous Necrosis/Viral Encephalopathy and Retinopathy	6
2.5.1	Description of Agent	6
2.5.2	Geographical Distribution and Temporal Trends	7
2.5.3	Short Description of Clinical Signs	7
2.5.4	Control/Preventative Measures	7
2.5.5	Other Host Species	7
2.6	Heart and Skeletal Muscle Inflammation	7
2.6.1	Description of agent	7
2.6.2	Geographical Distribution and Temporal Trends	7
2.6.3	Short Description of Clinical Signs	8
2.6.4	Control/preventative measures	8
2.6.5	Host Species	8
2.7	Cardiomyopathy Syndrome	8
2.7.1	Description of agent	8
2.7.2	Geographical Distribution and Temporal Trends	8

2.7.3	Short description of clinical signs	8
2.7.4	Control/Preventative Measures.....	8
2.7.5	Host Species	9
3	Bacterial Diseases of Farmed Fish	10
3.1	Francisellosis	10
3.1.1	Description of Agent.....	10
3.1.2	Geographical Distribution and Temporal Trends	10
3.1.3	Short description of clinical signs	10
3.1.4	Control/Preventative Measures.....	10
3.1.5	Host species.....	10
3.2	Rainbow Trout Fry Syndrome/Bacterial Coldwater Disease.....	10
3.2.1	Description of Agent.....	10
3.2.2	Geographical Distribution and Temporal Trends	11
3.2.3	Short description of clinical signs	11
3.2.4	Control/Preventative Measures.....	11
3.2.5	Host species.....	11
3.3	Enteric Redmouth Disease	12
3.3.1	Description of Agent.....	12
3.3.2	Geographical Distribution and Temporal Trends	12
3.3.3	Short description of clinical signs	12
3.3.4	Control/Preventative Measures.....	12
3.3.5	Host species.....	13
3.4	Red Spot Disease/Pseudomoniasis.....	13
3.4.1	Description of Agent.....	13
3.4.2	Geographical Distribution and Temporal Trends	13
3.4.3	Short description of clinical signs	13
3.4.4	Control/Preventative Measures.....	13
3.4.5	Host species.....	13
4	Parasitic Diseases of Farmed Fish.....	15
4.1	Amoebic Gill Disease	15
4.1.1	Description of Agent.....	15
4.1.2	Short Description of Clinical Signs	15
4.1.3	Geographical Distribution and Temporal Trends	15
4.1.4	Control/Preventative Measures.....	15
4.1.5	Host Species	15
	Finfish References.....	16
5	Viral Diseases in Farmed Molluscs	26
5.1	Ostreid herpesvirus 1 in bivalves	26
5.1.1	Description of Agent.....	26
5.1.2	Geographical Distribution and Temporal Trends	26
5.1.3	Short Description of Clinical Signs	26
5.1.4	Control/Preventative Measures.....	26

5.1.5	Host Species	27
6	Bacterial Diseases of Farmed Molluscs	28
6.1	<i>Vibrio</i> sp. infecting marine molluscs.....	28
6.1.1	<i>Vibrio splendidus</i> infecting Pacific oysters	28
6.1.2	<i>Vibrio aestuarianus</i> infecting oysters	28
6.1.3	<i>Vibrio harveyi</i> in abalone.....	29
6.2	<i>Nocardia crassostreae</i> in oysters	29
6.2.1	Description of Agent.....	30
6.2.2	Geographical Distribution and Temporal Trends	30
6.2.3	Short Description of Clinical Signs	30
6.2.4	Control/Preventative Measures.....	30
6.2.5	Host species.....	30
6.3	<i>Candidatus xenohalictis californiensis</i> in abalone.....	30
6.3.1	Description of Agent.....	30
6.3.2	Geographical Distribution and Temporal Trends	30
6.3.3	Short Description of Clinical Signs	31
6.3.4	Control/Preventative Measures.....	31
6.3.5	Host species.....	31
7	Diseases of Mollusc: Parasitic Diseases	32
7.1	<i>Bonamia exitiosa</i>	32
7.1.1	Description of Agent.....	32
7.1.2	Geographical Distribution and Temporal Trends	32
7.1.3	Short Description of Clinical Signs	32
7.1.4	Control/Preventative Measures.....	32
7.1.5	Host species.....	33
7.2	<i>Bonamia ostreae</i>	33
7.2.1	Description of Agent.....	33
7.2.2	Geographical Distribution and Temporal Trends	33
7.2.3	Short Description of Clinical Signs	33
7.2.4	Control/Preventative Measures.....	33
7.2.5	Host species.....	34
7.3	Marteiliosis	34
7.3.1	Description of Agent.....	34
7.3.2	Geographical Distribution and Temporal Trends	34
7.3.3	Short Description of Clinical Signs	34
7.3.4	Control/Preventative Measures.....	35
7.3.5	Host Species	35
7.4	Haplosporidiosis.....	35
7.4.1	Description of Agent.....	35
7.4.2	Geographical Distribution and Temporal Trends	35
7.4.3	Short Description of Clinical Signs	36
7.4.4	Control/Preventative Measures.....	36
7.4.5	Host Species	36

7.5	Perkinsosis	37
7.5.1	Description of Agent	37
7.5.2	Geographical Distribution and Temporal Trends	37
7.5.3	Short Description of Clinical Signs	37
7.5.4	Control/Preventative Measures	38
7.5.5	Host Species	38
	Molluscan References.....	39
	Author contact information.....	49

1 Background

The ICES Working Group on Pathology and Diseases of Marine Organisms (WGPDMO) provides annual reviews of national reports on the disease status of wild and farmed fish and molluscs in the ICES area. In 2004, the group published a first report collating this information from 1998-2002. This second report aims to provide an update on the status of the major diseases described in the original report and also to provide an overview of new diseases which have emerged since the previous report was published.

2 Viral Diseases of Farmed Fish

2.1 Infectious salmon anaemia

2.1.1 Description of Agent

Infectious salmon anaemia virus (ISAV), the causal agent of infectious salmon anaemia (ISA) is an enveloped virus consisting of eight single-stranded RNA segments. It is classified as the type species of the genus *Isavirus* within the family *Orthomyxoviridae*. Differentiation of ISAV isolates is based on sequencing a highly polymorphic region (HPR) of the haemagglutinin esterase (HE) gene (Kibenge *et al.*, 2007). All clinical isolates have deletions (HPRΔ) in this region while isolates which do not have deletions in the HPR, (HPR0), are avirulent. It has been shown that HPR0 ISAV is widespread in farmed Atlantic salmon, *Salmo salar*, (McBeath *et al.*, 2009; Christiansen *et al.*, 2011; Godoy *et al.*, 2013).

2.1.2 Geographical Distribution and Temporal Trends

Clinical ISA has now been reported in Norway, Scotland, Faroe Islands, eastern Canada, the eastern USA and Chile. ISA occurs each year in Norway; the number of cases varies annually with 15 cases reported in 2015 (Lyngstad *et al.*, 2008; Hjeltne *et al.*, 2016). Between 2001 and 2004 an ISA epidemic occurred in the Faroe Islands almost wiping out the Atlantic salmon industry. An ISA contingency plan was introduced in 2005 with the result that no clinical outbreaks of ISA have occurred since then. As part of the plan, all Atlantic salmon going to sea are vaccinated against ISA and a large scale monitoring programme involving monthly samples from all sites is in place. In Scotland, an ISA outbreak occurred in the Shetland Isles in 2009 (Murray *et al.*, 2010). The disease was eradicated with the result that Scotland is once again declared free of ISA. ISA outbreaks occurred in New Brunswick, Canada and Maine, USA between 1998 and 2004 (Gustafson *et al.*, 2007), but there have been no outbreaks at these locations since 2007. However, outbreaks were reported in Nova Scotia and Newfoundland, Canada in 2012 (OIE, 2012). The disease has been reported in farmed Atlantic salmon in Chile since 2007 (Godoy *et al.*, 2008; 2013).

2.1.3 Short Description of Clinical Signs

Infected fish are lethargic, congregate in the upper water level, gasp at the surface, go off feed and hang motionless at the sides of the cage. Affected fish may exhibit exophthalmia, ocular haemorrhage, distended abdomen and/or skin haemorrhage. Internal pathology may include dark, pale or yellow liver, ascites, pale gill and heart, enlarged spleen, petechial haemorrhage in visceral fat and a darkened foregut. Low haematocrit values (< 10) are a typical finding. Histological findings include multifocal haemorrhagic hepatic necroses that may become confluent to give the changes a “zonal” appearance, leaving areas around large veins intact (late stage of disease development). Focal congestion and dilatation of hepatic sinusoids, sometimes with distribution as described for necroses (early stage), and rupture of the sinusoidal endothelium with the presence of erythrocytes within the space of Disse (early sign) are also observed

2.1.4 Control/Preventative Measures

According to the OIE International Aquatic Animal Health Code and EU Directive 2006/88/EC (detection of HPR0 ISAV is not reportable in the EU), ISA is a notifiable disease which means that specific eradication protocols have to be implemented in areas which have been declared disease-free. In Canada ISA is reportable under the Health of Animals Act. ISA control plans are based on active surveillance, vaccination

and eradication of infected cages. These measures are supplemented with increased biosecurity, disease free certification and a greater traceability. The virus is known to be susceptible to a range of common disinfectants (Smail *et al.*, 2004) and to ultraviolet irradiation (Øye and Rimstad, 2001).

2.1.5 Other Host Species

Clinical ISA has only been reported in farmed Atlantic salmon, however the virus has been found in marine farmed rainbow trout, *Oncorhynchus mykiss*, without visible signs of disease. Virus can be detected in wild salmonids only and to date there is no evidence that the virus is present in non-salmonid fish (Raynard *et al.*, 2001; Plarre *et al.*, 2005).

2.2 Viral Haemorrhagic Septicaemia

2.2.1 Description of Agent

The causative agent of viral haemorrhagic septicaemia (VHS) is a rhabdovirus of the genus *Novirhabdovirus* within the family *Rhabdoviridae*. It is a single stranded, enveloped, RNA virus. Sequencing of the VHSV genome has revealed four major genotypes (Snow *et al.*, 2004; Einer-Jensen *et al.*, 2005):

Genotype I: Danish freshwater isolates (I_(unclassified)); predominantly continental Europe (Ia); northern Europe marine (Ib); continental Europe (Ic); Scandinavia – Baltic Sea and freshwater (Id); Black Sea region (Ie) (Cieslak *et al.*, 2016).

Genotype II: Baltic Sea marine isolates.

Genotype III: North Atlantic marine isolates.

Genotype IV: North American Pacific coast (IVa), Great Lakes (IVb), North American Atlantic coast (IVc), (Cieslak *et al.*, 2016).

2.2.2 Geographic Distribution and Temporal Trends

VHSV has been isolated throughout the Northern Hemisphere from a wide range of marine and freshwater species (Studer and Janies, 2011). VHS (genotype Ia) was reported in a rainbow trout farm in England in 2006 (Stone *et al.*, 2008) and an outbreak of genotype III VHS occurred in wrasse species (ballan, *Labrus bergylta*, corkwing, *Symphodus melops*, cuckoo, *Labrus mixtus*, goldsinny, *Ctenolabrus rupestris* and rock cook, *Centrolabrus exoletus*) held in a marine hatchery in the Shetland Isles, Scotland (Munro *et al.*, 2015). In 2007, the virus (genotype III) was detected in three marine rainbow trout sites in Norway (Dale *et al.*, 2009). There were two sites positive in 2008, one in 2009 and the country has remained free since. In Finland, VHS (genotype Id) continued to spread after its first isolation in 2000, with 24 farms reported infected by 2004 (Raja-Halli *et al.*, 2006). Denmark has been declared free of VHS after a long term eradication programme, with no clinical outbreaks since 2009 (Bang Jensen *et al.*, 2014). In N. America the majority of isolates belong to genotype IVa (Garver *et al.*, 2013). In 2005, VHS (genotype IVb) occurred for the first time in the Great Lakes Basin, North America, affecting a number of different species in Lake St Claire and Lake Ontario (Elsayed *et al.*, 2006).

2.2.3 Short Description of Clinical Signs

The clinical signs of classical VHS in freshwater salmonids are dark skin and pale gills, petechiae in the gills and in the skin, and haemorrhages in the orbits and exophthalmia.

Widespread petechiae developing to haemorrhages are observed in the peritoneal surfaces, in the swimbladder, in the skeletal muscles and in the meninges. The liver is pale with haemorrhages, and the spleen is often enlarged and reddish. High mortality is observed in the acute phase of the disease.

The clinical signs vary among species of marine fish. In Pacific salmonids, VHSV was isolated from fish without clinical signs. In Pacific cod (*Gadus macrocephalus*) and herring (*Clupea harengus pallasii*), infection appeared to be associated with skin lesions, while other Pacific fish species from which VHSV was isolated did not display any gross clinical signs. In European waters, the clinical signs in farmed salmonids and turbot, *Scophthalmus maximus*, were almost identical: classical signs were observed in freshwater salmonids, whereas VHSV in most other fish species was isolated from specimens showing no clinical signs.

2.2.4 Control/Preventative Measures

According to the OIE Aquatic Animal Health Code and EU Directive 2006/88/EC, VHS is a notifiable disease which means that, should the disease occur, specific eradication protocols have to be implemented in areas which have been declared disease-free. Strict biosecurity protocols are essential for remaining disease free. Øye and Rimstad (2001) demonstrated that VHSV is sensitive to ultraviolet irradiation.

2.2.5 Other Host Species

VHS virus has been isolated from 48 different marine and freshwater fish species in the Northern hemisphere (Skall *et al.*, 2005). The updated annex IV of EU Directive 2006/88/EC lists the following species as susceptible to VHSV, herring (*Clupea* sp.), whitefish (*Coregonus* sp.), pike (*Esox lucius*), haddock (*Gadus aeglefinus*), Pacific cod, Atlantic cod (*G. morhua*), Pacific salmon (*Oncorhynchus* sp.), rainbow trout, rockling (*Onos mustelus*), brown trout (*Salmo trutta*), turbot, sprat (*Sprattus sprattus*) and grayling (*Thymallus thymallus*). It is likely that wrasse will be added to this list in the near future (Munro *et al.*, 2015).

2.3 Pancreas Disease

2.3.1 Description of Agent

Pancreas disease (PD) is caused by the salmonid alphavirus (SAV), a member of the genus *Alphavirus* of the family *Togaviridae* (McLoughlin and Graham, 2007). Based on sequence analysis, six subtypes of SAV have been reported. In general, SAV1, 4, 5 and 6 have been associated with PD in Ireland or Scotland and SAV3 causes PD in Norwegian farmed Atlantic salmon (Fringuelli *et al.*, 2008). SAV2 has been associated with sleeping disease in freshwater rainbow trout, however it has now been isolated from diseased Atlantic salmon in Scotland and Norway (Graham *et al.*, 2012).

2.3.2 Geographical Distribution and Temporal Trends

Pancreas disease has emerged as a significant disease of farmed Atlantic salmon in Ireland, Scotland and Norway (McLoughlin and Graham, 2007). The disease is endemic in Ireland (Rodger and Mitchell, 2007), and has increased significantly in Scotland as well (Lester *et al.*, 2011). PD is a listed disease in Norway and an endemic zone was established for SAV3 cases in 2007 (Jansen *et al.*, 2010). A second zone was established for the newly identified SAV2 in 2012 (Hjortaas *et al.*, 2013). In 2015, there were 137 cases of PD registered in Norway (Hjeltnes *et al.*, 2016). There have been no reports of PD in Chile or North America. Sleeping disease of freshwater reared rainbow trout has

been reported in Scotland, England, France, Spain, Italy (Graham *et al.*, 2007b), Germany (Bergmann *et al.*, 2008) and Croatia (Vardić Smrzlić *et al.*, 2013).

2.3.3 Short Description of Clinical Signs

Early clinical signs include a cessation of feeding, lethargy and the observation of yellow faecal casts in the water column. A small percentage of survivors typically fail to thrive and become runts. The changes most commonly found in clinically diseased fish were severe loss of exocrine pancreatic tissue, cardiomyocytic necrosis and heart inflammation, inflammation of the red skeletal muscle and degeneration of white skeletal muscle. Muscle lesions are typical of hyaline degeneration with swollen fragmented eosinophilic sarcoplasm, central migration of myocytic nuclei and subsequent invasion of the sarcoplasm by phagocytic macrophages (McLoughlin *et al.*, 2002; Taksdal *et al.*, 2007).

2.3.4 Control/preventative measures

Pancreas disease was listed by the OIE in 2014 and is a list 3 disease in Norway, but is not a notifiable disease under EU Directive 2006/88/EC. A commercial vaccine is available and is widely used by the industry (Bang Jensen *et al.*, 2012). Regular screening of fish is recommended using serological (Graham *et al.*, 2003) or molecular techniques (Hodneland and Endresen, 2006) as an early indicator of infection. Strict biosecurity protocols are essential and the salmonid alphavirus is known to be sensitive to a range of common disinfectants (Graham *et al.*, 2007a). There are also indications that selective breeding for resistance to SAV infection is possible although research is still ongoing (Norris *et al.*, 2008).

2.3.5 Other Host Species

SAV has only been isolated from farmed Atlantic salmon and rainbow trout. The virus has been detected by PCR in wild flatfish; (common dab *Limanda limanda*, long rough dab *Hippoglossoides platessoides*, and plaice *Pleuronectes platessa*) off the coasts of Scotland (Snow *et al.*, 2010) and in common dab and plaice in Ireland (McCleary *et al.*, 2014). Bruno *et al.* (2014) demonstrated that SAV5 isolated from common dab, off the coast of Scotland could be grown on salmonid cell lines.

2.4 Infectious Pancreatic Necrosis

2.4.1 Description of Agent

Infectious pancreatic necrosis (IPN) is caused by a double-stranded, non-enveloped, RNA virus of the genus *Aquabirnavirus*. Aquabirnaviruses infect a wide range of fish species, aquatic molluscs and crustaceans. Seven genogroups of the IPN virus have been described (Blake *et al.*, 2001; Nishizawa *et al.*, 2005).

2.4.2 Geographical Distribution and Temporal Trends

Historically IPN has been a major problem for freshwater salmonid aquaculture, however with the expansion of the industry it has emerged in the last twenty years as a significant disease of marine farmed Atlantic salmon. In farmed Atlantic salmon, the disease has been reported in Scotland (Bain *et al.*, 2008), Ireland (Ruane *et al.*, 2009; 2015) and Norway (Hjeltnes *et al.*, 2016) where genogroup 5 isolates are the dominant form. The number of cases in Norway is decreasing, with 30 cases reported in 2015 (Hjeltnes *et al.*, 2016) and in Ireland there have been no clinical cases of IPN since 2012 (ICES, 2016). In contrast, IPN has spread from the coastal region of Finland to inland areas since 2012. Both genogroups 2 and 5 are found in Finland, however only genogroup 2

isolates occur in the inland areas (Eriksson-Kallio *et al.*, 2016). IPN has been present in Chile for many years where most isolations are genogroup 5, but genogroup 1 has also been detected (Calleja *et al.*, 2012). Genogroup 1 isolates also dominate in Mexico (Barrera-Mejía *et al.*, 2011). Although clinical IPN has not been recorded in Australia or New Zealand, an aquatic birnavirus which clusters with genogroup 5 isolates was reported in Tasmania (Davies *et al.*, 2010).

2.4.3 Short Description of Clinical Signs

Diseased fish usually appear dark with abdominal distension, particularly in fry. Swimming behaviour is lethargic: a spinning movement (whirling around the longitudinal axis) is characteristic. At necropsy, petechiae in the perivisceral adipose tissue are the most consistent lesions. Fry often show pronounced ascites, whereas post-smolts usually have little or no ascitic fluid and a remarkably dry body muscle. A whitish liver is most commonly found in fry. In the acute stage, extensive necrosis of the exocrine pancreas is the most prominent lesion at histopathological examination. In these lesions, individual acinar cells are seen at different stages of degeneration and cell death. Also, foci with necrotic remnants of exocrine pancreatic cells appearing as an amorphous eosinophilic mass are often found. In liver tissue, similar foci of necrotic eosinophilic hepatocytes may be found. Usually there are small necrotic foci in the pyloric intestinal epithelium and cell debris in the lumen of the pyloric caeca. Some individuals develop a chronic disease with fibroplasias of pancreatic tissue and emaciation (Roberts and Pearson, 2005).

2.4.4 Control/Preventative Measures

The disease has now become so widespread that it is no longer listed by the OIE and very few countries perform screening for the virus. The inland area of Finland has an IPN-free status for genogroup 5 IPNV under EC Decision 2010/221/EU. It is common practice to immunize smolts against IPN before going to sea as a control measure in combination with selective breeding and stricter biosecurity measures. Compared with other fish pathogenic viruses, the IPN virus is more resistant to inactivation (Smail *et al.*, 2003, 1993; Øye and Rimstad, 2001). It is however sensitive to a number of commonly used chlorine and iodine based disinfectants which can be used to inactivate the virus on equipment and the surface of eggs. In recent years the number of cases in Norway have reduced which is believed to be due to the use of more resistant strains of Atlantic salmon (Moen *et al.*, 2009).

2.4.5 Other Host Species

Aquabirnaviruses have been isolated from a variety of wild and farmed teleost fish, molluscs and crustacea in freshwater, estuarine and marine environments (Hill and Way, 1995; Ahne *et al.*, 2003).

2.5 Viral Nervous Necrosis/Viral Encephalopathy and Retinopathy

2.5.1 Description of Agent

Viral nervous necrosis (VNN) is caused by a piscine nodavirus, of the genus *Betanodavirus* from the *Nodaviridae* family which are small, non-enveloped RNA viruses.

2.5.2 Geographical Distribution and Temporal Trends

The disease has been commonly found in the Mediterranean region. Over the last decade nodavirus infections have occurred in marine fish farming in Norway and Scotland, most notably in turbot (Johansen *et al.*, 2004) and Atlantic cod (Hellberg *et al.*, 2010). It also occurs on the Atlantic coast of North America (Johnson *et al.*, 2002).

2.5.3 Short Description of Clinical Signs

Clinical signs are associated with lesions in the brain and retina and include failure to control movement and swim bladder function. Sight and colouration are also affected, however the most significant outcome is mortality, particularly among larvae (Munday *et al.*, 2002). Mortality has also occurred among harvest size sea bass, groupers and Atlantic halibut and may be related to elevated water temperature.

2.5.4 Control/Preventative Measures

VNN is listed by the OIE and is a list 3 disease in Norway, but is not a notifiable disease under EU Directive 2006/88/EC. In the absence of vaccines or treatment, biosecurity is the most effective way to limit the introduction and spread of nodavirus within populations of cultured finfish.

2.5.5 Other Host Species

VNN is associated with mortalities in larvae and juveniles of several marine species worldwide, including haddock *Melanogrammus aeglefinus*, Atlantic halibut *Hippoglossus hippoglossus*, Atlantic cod, turbot, Gilthead sea bream *Sparus aurata* and sea bass *Dicentrarchus labrax*.

2.6 Heart and Skeletal Muscle Inflammation

2.6.1 Description of agent

Heart and skeletal muscle inflammation (HSMI) is believed to be a viral disease although the causative agent has not yet been conclusively identified. Studies have suggested that a novel reovirus, termed piscine reovirus (PRV) may be the causative agent (Palacios *et al.*, 2010). Direct localisation of the virus in heart tissues of HSMI affected fish has indicated an association of the virus with HSMI. The virus has also been detected in healthy farmed and wild fish sometimes at elevated titres similar to those measured in fish diagnosed with HSMI (Garseth *et al.*, 2013). The development of HSMI may involve environmental co-factors which influence the pathogenicity of an infectious agent (Løvoll *et al.*, 2012).

2.6.2 Geographical Distribution and Temporal Trends

HSMI was first reported in farmed Atlantic salmon in Norway in 1999 (Kongtorp *et al.*, 2004a). Outbreaks of HSMI have been officially recorded in farmed salmon in Norway since 2004 and the disease is now widespread in Norwegian salmon aquaculture with 135 cases recorded in 2015 (Hjeltnes *et al.*, 2016). The disease has also been reported in Scotland (Ferguson *et al.*, 2005) and in Ireland (ICES, 2016). The virus was detected, by qPCR, in wild Atlantic salmon broodstock and progeny for the first time in Denmark in 2014 (ICES, 2016). In 2014, a new disease showing HSMI-like clinical symptoms was reported in freshwater farmed rainbow trout in Norway (Olsen *et al.*, 2015). Outside Europe, HSMI has been described in farmed Atlantic salmon and coho salmon, *O. kisutch*, in Chile (Godoy *et al.*, 2016). The virus has been detected in farmed Atlantic salmon in the Pacific north-west of America and Canada (Siah *et al.*, 2015).

2.6.3 Short Description of Clinical Signs

The disease is most frequently reported in spring and early summer, approximately 6 months after sea transfer. Morbidity can be high and mortality up to 20%. Gross signs include pallor and loose texture of the heart, and pericardial haemorrhage with ascites. Haematocrit tends to be normal. Microscopic lesions include myocardial necrosis and a severe ventricular myocarditis of both compact and spongy layers with a predominantly monocytic infiltrate. Coincident epicarditis is common. Lesions occasionally observed include myocarditis of red skeletal muscle, focal necrosis of liver, and oedema and congestion in several organs (Kongtorp *et al.*, 2004b).

2.6.4 Control/preventative measures

None available. The disease is not listed by the OIE nor under EU Directive 2006/88/EC. In 2014 the disease was delisted in Norway (Hjeltnes, 2016).

2.6.5 Host Species

HSMI has only been diagnosed in farmed Atlantic salmon. The virus has been detected in wild Atlantic salmon and sea trout, *Salmo trutta*, (Garseth *et al.*, 2013) and in a range of marine fish (Wiik-Nielsen *et al.*, 2012a) by qPCR in Norway. The virus has also been detected in pacific salmon, namely coho salmon in Chile (Godoy *et al.*, 2016) and coho and chinook salmon, *O. tshawytscha*, on the Pacific coast of America and Canada (Siah *et al.*, 2015).

2.7 Cardiomyopathy Syndrome

2.7.1 Description of agent

A novel virus, the piscine myocarditis virus (PMCV) has been proposed as the causative agent of cardiomyopathy syndrome (CMS). The virus has been detected in fish with clinical CMS and has also been localised in the infected tissues (Haugland *et al.*, 2011). Infection trials have also supported the hypothesis that CMS has a viral aetiology (Bruno and Noguera, 2009; Fritsvold *et al.*, 2009). The virus may be present in farmed salmon for long periods without any signs of clinical disease (Wiik-Nielsen *et al.*, 2012b).

2.7.2 Geographical Distribution and Temporal Trends

CMS was first diagnosed in Norway in the 1980's and later in Scotland and the Faroe Islands (Brun *et al.*, 2003). A single case of CMS was reported in Ireland with very low mortalities in 2012 (Rodger *et al.*, 2014). The number of cases reported in Norway was 105 in 2015 (Hjeltnes *et al.*, 2016).

2.7.3 Short description of clinical signs

Gross signs include haemopericardium (cardiac tamponade) resulting from rupture of the atrium or sinus venosus. Microscopic lesions include myocardial degeneration and coagulative necrosis associated with the spongy layer of the ventricle and atrium. An infiltrate comprised of lymphocytes and macrophages can be observed.

2.7.4 Control/Preventative Measures

None available. The disease is not listed by the OIE nor under EU Directive 2006/88/EC.

2.7.5 Host Species

CMS has only been diagnosed in farmed Atlantic salmon although the virus has been detected by PCR in wild Norwegian Atlantic salmon (Garseth *et al.*, 2012). In Norway, a range of wild marine fish species have been screened for PMCV by PCR and only samples from the Atlantic argentine, *Argentina silus* were positive (Böckerman *et al.*, 2011).

3 Bacterial Diseases of Farmed Fish

3.1 Francisellosis

3.1.1 Description of Agent

Bacteria belonging to the genus *Francisella* are Gram-negative, non-motile, aerobic, facultative intracellular organisms belonging to the γ -proteobacteria. Most, if not all strains isolated from teleost fish belong to either *F. noatunensis* subsp. *orientalis* in warm water fish species or *F. noatunensis* subsp. *noatunensis* in coldwater fish species (Colquhoun and Duodu, 2011). Within the ICES area, francisellosis, caused by *F. noatunensis* subsp. *noatunensis*, has emerged as a major disease in farmed Atlantic cod (Ottem *et al.*, 2009).

3.1.2 Geographical Distribution and Temporal Trends

Francisellosis was diagnosed for the first time in farmed Atlantic cod in 2004 in Norway (Olsen *et al.*, 2006), however the bacteria (*F. noatunensis* subsp. *noatunensis*) have also been detected in archived samples of wild-caught cod in the North Sea (Zerihun *et al.*, 2011). The bacteria have also been detected by real-time PCR in wild cod along the Norwegian coast line (Ottem *et al.*, 2008). There was one reported case of francisellosis in Norway in 2014 and none in 2015, down from a peak of 14 in 2008 (Hjeltnes *et al.*, 2016). In 2009, the disease occurred in Ireland in a stock of wild-caught cod held in captivity as broodstock (Ruane *et al.*, 2015).

3.1.3 Short description of clinical signs

Francisellosis is principally a chronic systemic granulomatous inflammatory disease with varying degrees of mortality. The gross signs observed among most host species are similar and include disseminated white nodules of various sizes in liver, spleen and kidney although most tissues and organs may be affected. Splenomegaly and serosanguinous ascites has been observed in cod. Microscopic lesions include widespread chronic granulomatous inflammation in all organs, associated with variable numbers of Gram-negative bacteria (Birkbeck *et al.*, 2011).

3.1.4 Control/Preventative Measures

None available. The disease is not listed by the OIE nor under EU Directive 2006/88/EC, but is a list 3 disease in Norway.

3.1.5 Host species

F. noatunensis subsp. *noatunensis* has primarily been reported infecting Atlantic cod, however there is a single report of francisellosis in Atlantic salmon in Chile (Birkbeck *et al.*, 2007).

3.2 Rainbow Trout Fry Syndrome/Bacterial Coldwater Disease

3.2.1 Description of Agent

Flavobacterium psychrophilum is the causative agent of Rainbow Trout Fry Syndrome and Bacterial Coldwater Disease. *F. psychrophilum* is a Gram-negative, slender, flexible rod that displays gliding motility. The bacterium requires specific media for isolation and growth, e.g. tryptone yeast extract salts agar (Holt *et al.*, 1993). The name “psychrophilum” refers to the low growth optimum of this bacterium, which is around

15°C *in vitro* (Holt *et al.* 1993) with no growth above 25°C. Other features of this bacterium are its high proteolytic activity as well as its lack of growth in media with > 1% salt under laboratory conditions (Dalsgaard and Madsen, 2000).

3.2.2 Geographical Distribution and Temporal Trends

F. psychrophilum was originally isolated from diseased juvenile coho salmon in the USA in the 1940s (Borg, 1948, 1960), and was first found outside North America during the 1980s (see Nematollahi *et al.*, 2003), where it appeared in Europe, South America, Japan, Korea and Australia. It has for many years been considered an obligate freshwater pathogen due to its sensitivity to salt, but there have been reports of disease outbreaks with *F. psychrophilum* in rainbow trout reared in brackish water in Finland (ICES, 2001) and spawning bream, *Abramis brama*, on the Baltic coast of Sweden (ICES, 2013). In 2008, *F. psychrophilum* was isolated for the first time in farmed Atlantic salmon in Norway, causing septicaemia and mortalities (Nilsen *et al.*, 2011a). The disease has regularly caused high mortalities among rainbow trout farmed in the Scandinavian countries such as Norway (Nilsen *et al.*, 2011b). Baltic salmon brood fish have been shown to be carriers of *F. psychrophilum* during their spawning migration (Ekman *et al.* 1999), and laboratory experiments with water microcosms have shown that the bacterium is able to survive salinities up to 0.6‰ (Madetoja *et al.*, 2003). In all, this means that *F. psychrophilum* is able to adapt to more saline environments than originally expected (Nilsen *et al.*, 2011b).

3.2.3 Short description of clinical signs

Disease caused by *F. psychrophilum* results in septicaemia and/or skin lesions and fin rot (Nematollahi *et al.*, 2003). In salmonid fry, the primary sign will be high mortalities due to septicaemia, up to 90 % in an affected population, whereas larger fish display skin lesions with lower mortalities (Borg, 1960; Holt *et al.*, 1993; Nematollahi *et al.*, 2003).

3.2.4 Control/Preventative Measures

F. psychrophilum disease outbreaks can be treated with antibiotics such as oxytetracycline, amoxicillin, oxolinic acid and florfenicol (Barnes and Brown, 2011), however resistance to antibiotic treatments is becoming an issue (Bruun *et al.*, 2003). There are no registered vaccines against the disease at present, therefore prophylactic measures like good management procedures (including egg disinfection) as outlined by Madsen and Dalsgaard (2008) are recommended. Phage therapy as a potential method for controlling *F. psychrophilum* infections is also being investigated (Madsen *et al.*, 2013).

3.2.5 Host species

Juvenile rainbow trout and coho salmon are particularly susceptible (Nematollahi *et al.*, 2003). However, *F. psychrophilum* infections have been reported in a wide range of both anadromous and non-anadromous salmonids of various sizes. In addition, *F. psychrophilum* has either caused disease or been detected in Japanese eel *Anguilla japonica*, European eel *Anguilla anguilla*, common carp *Cyprinus carpio*, crucian carp *Carassius carassius*, tench *Tinca tinca*, ayu *Plecoglossus altivelis*, pale chub *Zaco platypus*, perch *Perca fluviatilis* and roach *Rutilus rutilus* (Barnes and Brown, 2011).

3.3 Enteric Redmouth Disease

3.3.1 Description of Agent

The aetiological agent of enteric redmouth disease (ERM) is *Yersinia ruckeri*, a gram negative, slightly curved, rod shaped bacteria. Most of the bacteria are motile due to the presence of flagellae (Barnes, 2011). Strains of *Y. ruckeri* can be classified on the basis of biotype, serotype and outer-membrane protein (OMP) type (Tobback *et al.*, 2007) however a combination of these is useful for discriminating between strains (Davies, 1991; Wheeler *et al.*, 2009). The most common *Y. ruckeri* serovar is O1, with the most virulent being the O1a serotype Hagerman strain (Davies, 1991). Serovar O1 *Y. ruckeri* grow well on tryptone soya agar with or without 5% blood. After a 48 h incubation at 20-25°C, round, raised, shiny off-white colonies of 2-3 mm in diameter develop (Austin and Austin, 2007). Over the last decade, a new biotype 2 strain has emerged causing mortalities in Europe (Wheeler *et al.*, 2009). There were 34 confirmed sites affected by ERM in Norway in 2015 (Hjeltnes *et al.*, 2016).

3.3.2 Geographical Distribution and Temporal Trends

ERM was first described in the USA in 1955 and became widespread in farmed salmonid species throughout the USA and Canada (Furones *et al.*, 1993). The disease is now present throughout Europe, Australia, Chile and South Africa (Barnes, 2011). The most probable route of spread has been by horizontal transmission through the import of live fish and ova. In recent years, Biotype 2 strains have emerged causing significant losses in rainbow trout facilities in the UK (Austin *et al.*, 2003), Spain (Fouz *et al.*, 2006), USA (Arias *et al.*, 2007) and Finland (ICES, 2012).

3.3.3 Short description of clinical signs

ERM can affect fish of all ages, but is most acute in juveniles, while in larger fish the disease appears as a more chronic condition. The characteristic haemorrhages around the oral cavity led to the name 'redmouth' disease, although this is often not apparent in many cases of disease (Tobback *et al.*, 2007). The disease is mainly seen as a generalised haemorrhagic septicaemia with anorexia, darkening of the skin, haemorrhages in the skin, gills and at the base of the fins and the lateral line. Bleeding is also common in internal organs; swimbladder, liver, spleen, pancreas and visceral fat in connection with the pyloric caecae. Kidney and spleen are swollen and darkened in colour (Roberts, 2012; Barnes, 2011). McArdle (2014) recently reported that heart pathology was a consistent finding in outbreaks in Ireland over a number of years and may explain the behavioural changes of diseased fish.

3.3.4 Control/Preventative Measures

ERM has successfully been controlled by the use of a monovalent vaccine based on a Hagerman type strain, biotype 1. Following reports of vaccine failures, connected to new strains classified as biotype 2, new vaccines contain antigens from both the Hagerman type strain and Biotype 2 (Deshmuk *et al.*, 2012). This vaccine offers good protection when administered by intra-peritoneal injection supplemented with an additional booster by immersion (Chettri *et al.*, 2013). Antibiotics also play a role in controlling infection as *Y. ruckeri* is sensitive to a range of antibiotics (e.g. oxytetracycline, oxolinic acid and potentiated sulphonamides), however there are increasing reports of antibiotic resistance (Tobback *et al.*, 2007).

3.3.5 Host species

ERM is primarily a disease infecting salmonid fish species, particularly rainbow trout. The agent has also been isolated from a wide range of non-salmonid fish species, though not all exhibited pathology related to infection with *Y. ruckeri* (Furones *et al.*, 1993).

3.4 Red Spot Disease/Pseudomoniasis

3.4.1 Description of Agent

Pseudomona anguilliseptica, a Gram-negative motile rod, is an opportunistic pathogen for a variety of fish species cultured in marine and brackish waters worldwide. It is the aetiological agent of 'red spot disease' also known as 'Sekiten-byo' disease of eels. The bacteria can be isolated from blood, kidney, liver and spleen samples on a nutrient agar, supplemented with 10% blood. The bacteria are quite slow growing and incubation at 20-25°C should continue for at least 7 days, when round, raised, shiny, pale-grey colonies, less than 1 mm in diameter develop (Wakabashi and Egusa, 1972). Serologically, two major groups have been established on the basis of their O-antigens with serotype O2 including the majority of eel isolates and serotype O1 including isolates from all other fish species (Lopez-Romalde *et al.*, 2003).

3.4.2 Geographical Distribution and Temporal Trends

P. anguilliseptica has been reported from Japan and Taiwan (Wakabayashi and Egusa, 1972; Nakai *et al.*, 1985), Kuwait (Al-Marzouk, 1999) and in a number of European countries such as Scotland, Finland, France, Spain and The Netherlands (Stewart *et al.*, 1983; Wiklund and Bylund, 1990; Berthe *et al.*, 1995; Domenech *et al.*, 1999; Haenan and Davidse, 2001). In Finland, *P. anguilliseptica* has become a significant cause of disease outbreaks in rainbow trout reared in brackish water during extremely warm summers when water temperatures rise above 20°C and rearing conditions are poor (ICES, 2012).

3.4.3 Short description of clinical signs

Diseased fish have petechial haemorrhage of the skin, peritoneum and liver. Kidney can have liquefactive necrosis (Wakabashi and Egusa, 1972; Ellis *et al.*, 1983; Wiklund and Bylund, 1990). Diseased rainbow trout in Finnish brackish water farms appear dark in colour, with a sluggish appearance and large hyperaemic and oedematic areas are observed on the sides of the fish.

3.4.4 Control/Preventative Measures

In Finland *P. anguilliseptica* infections in rainbow trout have successfully been treated with Sulfa-Trimetoprim (30 mg/Kg for 7 days) in combination with improve husbandry on the farms. There is no commercial vaccine available in Europe.

3.4.5 Host species

P. anguilliseptica was first described as red spot disease of Japanese eels (Wakabashi and Egusa, 1972). Since then, the pathogen has been recorded in a range of fish species including European eel, (Stewart *et al.*, 1983), rainbow trout, sea trout, white fish *Coregonus sp.*, Baltic herring, *Clupea harengus membras* (Wiklund and Lönnström, 1994; Lönnström *et al.*, 1994), sea bass, *Dicentrarchus labrax* (Berthe *et al.*, 1995), Gilthead sea bream, *Sparus aurata* (Domenech *et al.*, 1999), orange-spotted grouper, *Epinephelus coioides* (Al-Marzouk, 1999), Atlantic cod, (Ferguson *et al.*, 2004) and turbot (Magi *et al.*,

2009). *P. anguilliseptica* has also been isolated from lumpsucker (*Cyclopterus lumpus*) used as cleaner fish on Atlantic salmon farms in Norway (Hjeltnes *et al.*, 2016).

4 Parasitic Diseases of Farmed Fish

4.1 Amoebic Gill Disease

4.1.1 Description of Agent

AGD was initially ascribed to *Neoparamoeba pemaquidensis*, based on morphological (Dyková *et al.*, 2000) and molecular characterisation (Wong *et al.*, 2004). The isolation of *N. branchiphila* from AGD-affected fish (Dyková *et al.*, 2005) meant that the disease may have a mixed aetiology. However, issues still remained with the development of a reproducible experimental challenge model using both species (Morrison *et al.*, 2004; Vincent *et al.*, 2007). Using molecular techniques, Young *et al.* (2008) showed that *N. perurans* was the aetiological agent of AGD which was subsequently confirmed by laboratory trials and fulfilment of Koch's Postulates (Crosbie *et al.*, 2012).

4.1.2 Short Description of Clinical Signs

AGD is characterised by multifocal lesions that appear as pale gill tissue, or white mucoid spots and plaques. The main histological feature of the disease is prominent epithelial hyperplasia resulting in a complete lamellar fusion. Large mucous cells are often situated on the surface of the hyperplastic epithelium and between the lamellae, with significant leucocyte infiltration (Mitchell and Rodger, 2011).

4.1.3 Geographical Distribution and Temporal Trends

The disease has long been associated with farmed Atlantic salmon in Tasmania (Munday *et al.*, 1993) and has also been reported in farmed Atlantic salmon in Ireland (Palmer *et al.*, 1997), Scotland, Norway (Steinum *et al.*, 2008) and Chile (Bustos *et al.*, 2011). The disease has also been reported in the Mediterranean (Dyková *et al.*, 2000; Munday *et al.*, 2001) and in South Africa (Mouton *et al.*, 2014). The disease re-emerged in Ireland and Scotland in 2012 and in Norway and the Faroe Islands in 2014 (Rodger, 2014; Oldham *et al.*, 2016). AGD has also been reported in Canada (ICES, 2016).

4.1.4 Control/Preventative Measures

The recommended treatment for AGD is a 2 – 3 h freshwater bath (Clark *et al.*, 2003). Treatment with hydrogen peroxide has also shown to be effective (Adams *et al.*, 2012). Regular screening of the gills through gill scoring (Taylor *et al.*, 2009) and non-lethal sampling for molecular diagnostics (Downes *et al.*, 2017) have proven useful for disease monitoring. AGD is not a notifiable disease.

4.1.5 Host Species

Amoebic Gill Disease (AGD) primarily affects salmonids and was first described in marine reared coho salmon (*S. kisutch*) in Washington and California, USA (Kent *et al.*, 1988) and in Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*) in Tasmania (Munday *et al.*, 1993). However, the disease has also been reported in turbot, *Psetta maxima* and sea bass *Dicentrarchus labrax* (Munday *et al.*, 2001) and has recently been described in ballan wrasse, *Labrus bergylta* (Karlsbakk *et al.*, 2013) and lumpsucker (Hjeltnes *et al.*, 2016).

Finfish References

- Adams, M.B., Crosbie, P.B.B., Nowak, B.F. 2012. Preliminary success using hydrogen peroxide to treat Atlantic salmon, *Salmo salar* L., affected with experimentally induced amoebic gill disease (AGD). *Journal of Fish Diseases* 35:839-848.
- Ahne, W., Blake, S., Essbauer, S. and Nicholson, B.L. 2003. Characterization of aquabirnaviruses from flounder *Pseudopleuronectes americanus* and mummichog *Fundulus heteroclitus* in the Chesapeake Bay, Virginia, USA. *Diseases of Aquatic Organisms* 56: 201-206.
- Al-Marzouk, A.E. 1999. Association of *Pseudomonas anguilliseptica* with mortalities in cultures of marine orange-spotted grouper, *Epinephelus coioides*, in Kuwait. *Fish Pathology* 34: 167-168.
- Arias, C.R., Olivares-Fuster, I. and Hayden, K. 2007. First report of *Yersinia ruckeri* biotype 2 in the USA. *Journal of Aquatic Animal Health* 19:35-40.
- Austin, B. and Austin, D.A. (eds.) 2007. *Bacterial fish pathogens: diseases of farmed and wild fish* 4th Ed. Praxis Publishing, Chichester, UK. 553pp.
- Austin, D.A., Robertson, P.A.W. and Austin, B. 2003. Recovery of a new biogroup of *Yersinia ruckeri* from diseased rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Systematic and Applied Microbiology* 26: 127-131.
- Bain, N., Gregory, A. and Raynard, R. 2008. Genetic analysis of infectious pancreatic necrosis virus from Scotland. *Journal of Fish Diseases* 31: 37-47.
- Bang Jensen, B., Kristoffersen, A.B., Myr, C. and Brun, E. 2012. Cohort study of effect of vaccination on pancreas disease in Norwegian salmon aquaculture. *Diseases of Aquatic Organisms* 102: 23-31.
- Bang Jensen, B., Ersbøll, A.K., Korsholm, H., Skall, H.K. and Olesen, N.J. 2014. Spatio-temporal risk factors for viral haemorrhagic septicaemia (VHS) in Danish aquaculture. *Diseases of Aquatic Organisms* 109: 87-97.
- Barnes, A.C. 2011. Enteric redmouth disease (ERM). In *Fish diseases and disorders Vol. 3: Viral, bacterial and fungal infections* 2nd Ed. (eds. Woo, P.T.K. and Bruno, D.W.). CAB International, Wallingford, UK. pp 484-511.
- Barnes, M.E. and Brown, M.L. 2011. A review of *Flavobacterium psychrophilum* biology, clinical signs, and bacterial cold water disease prevention and treatment. *The Open Fish Science Journal* 4: 40-48.
- Barrera-Mejía, M., Martínez, S., Ortega, C. and Ulloa-Arvizu, R. 2011. Genotyping of infectious pancreatic necrosis virus isolates from Mexico State. *Journal of Aquatic Animal health* 23: 200-206.
- Bergmann, S.M., Fichtner, D., Riebe, R. and Castric, J. 2008. First isolation and identification of sleeping disease virus (SDV) in Germany. *Bulletin of the EAAP* 28: 148-156.
- Berthe, F.C.J., Nichel, C. and Bernardet, J.F. 1995. Identification of *Pseudomonas anguilliseptica* isolated from several species in France. *Diseases of Aquatic Organisms* 21: 151-155.
- Birkbeck, T.H., Feist, S.W. and Verner-Jeffreys, D.W. 2011. *Francisella* infections in fish and shellfish. *Journal of Fish Diseases* 34: 173-187.
- Birkbeck, T.H., Bordevik, M., Frøystad, M.K. and Baklien, Å. 2007. Identification of *Francisella* sp. from Atlantic salmon, *salmo salar* L., in Chile. *Journal of Fish Diseases* 30: 505-507.
- Blake, S., Ma, J.Y., Caporale, D.A., Jairath, S. and Nicholson, B.L. 2001. Phylogenetic relationships of aquatic birnaviruses based on deduced amino acid sequences of genome segment A cDNA. *Diseases of Aquatic Organisms* 45: 89-102.
- Böckerman, I., Wiik-Nielsen, C.R., Sindre, H., Johansen, R. And Tengs, T. 2011. Prevalence of piscine myocarditis virus (PMCV) in marine fish species. *Journal of Fish Diseases* 34: 955-957.

- Borg, A.F. 1948. Studies on myxobacteria associated with diseases in salmonid fishes. PhD thesis. University of Washington, Seattle.
- Borg, A.F. 1960. Studies on myxobacteria associated with diseases in salmonid fishes. American Association for the Advancement of Science, Wildlife Disease no. 8, 85 p, Washington D.C.
- Brun, E., Poppe, T., Skrudland, A. and Jarp, J. 2003. Cardiomyopathy syndrome in farmed Atlantic salmon *Salmo salar*: occurrence and direct financial losses for Norwegian aquaculture. *Diseases of Aquatic Organisms* 56: 2411-247.
- Bruno, D.W. and Noguera, P.A. 2009. Comparative experimental transmission of cardiomyopathy syndrome (CMS) in Atlantic salmon *Salmo salar*. *Diseases of Aquatic Organisms* 87: 235-242.
- Bruno, D.W., Noguera, P.A., Black, J., Murray, W., Macqueen, D.J. and Matejusova, I. 2014. Identification of a wild reservoir of salmonid alphavirus in common dab *Limanda limanda*, with emphasis on virus culture and sequencing. *Aquaculture Environment Interactions* 5: 89-98.
- Bruun, M.S., Madsen, L. and Dalsgaard, I. 2003. Efficiency of oxytetracycline treatment in rainbow trout experimentally infected with *Flavobacterium psychrophilum* strains having different in vitro antibiotic susceptibilities. *Aquaculture* 215: 11-20.
- Bustos, P.A., Young, N.D., Rozas, M.A., Bohle, H.M., Ildefonso, R.S., Morrison, R.N., Nowak, B.F. 2011. Amoebic gill disease (AGD) in Atlantic salmon (*Salmo salar*) farmed in Chile. *Aquaculture* 310:281-288.
- Calleja, F., Godoy, M.G., Cárcamo, J.G., Bandín, I., Yáñez, A.J., Dopazo, C.P., Kibenge, F.S. and Avendaño-Herrera, R. 2012. Use of reverse transcription-real time polymerase chain reaction (real time RT-PCR) assays with Universal Probe Library (UPL) probes for the detection and genotyping of infectious pancreatic necrosis virus strains isolated in Chile. *Journal of Virological Methods* 183: 80-85.
- Chettri, J.K., Deshmuk, S., Holten-Andersen, L., Jafaar, R.M., Dalsgaard, I. and Buchmann, K. 2013. Comparative evaluation of administration methods for a vaccine protecting rainbow trout against *Yersinia ruckeri* O1 biotype 2 infections. *Veterinary Immunology and Immunopathology* 154: 42-47.
- Christiansen, D.H., Østergaard, P.S., Snow, M., Dale, O.B. and Falk, K. 2011. A low-pathogenic variant of infectious salmon anaemia virus (ISAV-HPR0) is highly prevalent and causes a non-clinical transient infection in farmed Atlantic salmon (*Salmo salar* L.) in the Faroe Islands. *Journal of General Virology* 92: 909-918.
- Cieslak, M., Mikkelsen, S.S., Skall, H.F., Baud, M., Diserens, N., Engelsma, M.Y., Haenen, O.L.M., Mousakhani, S., Panzarin, V., Wahli, T., Olesen, N.J. and Schutze, H. 2016. Phylogeny of the viral hemorrhagic septicaemia virus in European aquaculture. *PLoS ONE* 11(10): e0164475.
- Clark, G., Powell, M., Nowak, B. 2003. Effects of commercial freshwater bathing on reinfection of Atlantic salmon, *Salmo salar*, with Amoebic Gill Disease. *Aquaculture* 219:135-142.
- Colquhoun, D.J. and Duodu, S. 2011. *Francisella* infections in farmed and wild aquatic organisms. *Veterinary Research* 42:47.
- Crosbie, P.B.B., Bridle, A.R., Cadoret, K., Nowak, B.F. 2012. *In vitro* cultured *Neoparamoeba perurans* causes amoebic gill disease in Atlantic salmon and fulfils Koch's postulates. *International Journal for Parasitology* 42:511-515.
- Dale, O.B., Ørpetveit, I., Lyngstad, T.M., Kahns, S., Skall, H.F., Olesen, N.J. and Dannevig, B.H. 2009. Outbreak of viral haemorrhagic septicaemia (VHS) in seawater farmed rainbow trout in Norway caused by VHS virus Genotype III. *Diseases of Aquatic Organisms* 85: 93-103.
- Dalsgaard, I. and Madsen, L. 2000. Bacterial pathogens in rainbow trout, *Oncorhynchus mykiss* (Walbaum), reared at Danish freshwater farms. *Journal of Fish Diseases* 23: 199-209.

- Davies, K.R., McColl, K.A., Wang, L.F., Yu, M., Williams, L.M. and Crane, M.St.J. 2010. Molecular characterisation of Australasian isolates of aquatic birnaviruses. *Diseases of Aquatic Organisms* 93: 1-15.
- Davies, R.L. 1991 Clonal analysis of *Yersinia ruckeri* based on biotypes, serotypes and outer membrane protein-types. *Journal of Fish Diseases* 14: 221-228.
- Deshmuk, S., Raida, M.K., Chettri, J.K., Kania, P.W. and Buchmann, K. 2012. Comparative protection of two commercial vaccines against *Yersinia ruckeri* serotype O1 and biotype 2 in rainbow trout (*Onchorhynchus mykiss*). *Veterinary Immunology and Immunopathology* 145: 379-385.
- Domenech, A., Fernandez-Garayzabal, J.F., Garcia, J.A., Cutuli, M.T., Blanco, M., Gibello, A., Moreno, M.A. and Dominguez, L. 1999. Association of *Pseudomonas anguilliseptica* infection with 'winter disease' in sea bream, *Sparus aurata* L. *Journal of Fish Diseases* 22:69-71.
- Downes, J.K., Rigby, M.L., Taylor, R.S., Maynard, B.T., MacCarthy, E., O Connor, I., Marcos-Lopez, M., Rodger, H.D., Collins, E., Ruane, N.M. and Cook, M.T. 2017. Evaluation of non-destructive molecular diagnostics for the detection of *Neoparamoeba perurans*. *Frontiers in Marine Science* 4: 61.
- Dyková, I., Figueras, A., Peric, Z. 2000. *Neoparamoeba* Page, 1987: light and electron microscopic observations on six strains of different origin. *Diseases of Aquatic Organisms* 43:217-223.
- Dyková, I., Nowak, B.F., Crosbie, P.B.B., Fiala, I., Pecková, H., Adams, M.B., Macháčková, B., Dvořáková, H. 2005. *Neoparamoeba branchiphila* n. sp., and related species of the genus *Neoparamoeba* Page, 1987: morphological and molecular characterization of selected strains. *Journal of Fish Diseases* 28:49-64.
- Einer-Jensen, K., Winton, J. And Lorenzen, N. 2005. Genotyping of the fish rhabdovirus, viral haemorrhagic septicaemia virus, by restriction fragment length polymorphisms. *Veterinary Microbiology* 106: 167-178.
- Ekman, E., Börjeson, H. and Johansson, N. 1999. *Flavobacterium psychophilum* in Baltic salmon *Salmo salar* brood fish and their offspring. *Diseases of Aquatic Organisms* 37: 159-163.
- Ellis, A.E., Dear, G. and Stewart, D.J. 1983. Histopathology of Sekiten-byo caused by *Pseudomonas anguilliseptica* in the European eel, *Anguilla Anguilla* L., in Scotland. *Journal of Fish Diseases* 6: 77-79.
- Elsayed, E., Faisal, M., Thomas, M., Whelan, G., Batts, W. and Winton, J. 2006. Isolation of viral haemorrhagic septicaemia virus from muskellunge, *Esox masquinongy* (Mitchill), in Lake St. Claire, Michigan, USA reveals a new sub-lineage of the North America genotype. *Journal of Fish Diseases* 29: 611-619.
- Eriksson-Kallio, A.M., Holopainen, R., Viljamaa-Dirks, S., Vennerström, P., Kuukka-Anttila, H., Koski, P. and Gadd, T. 2016. Infectious pancreatic necrosis virus (IPNV) strain with genetic properties associated with low pathogenicity at Finnish fish farms. *Diseases of Aquatic Organisms* 118: 21-30.
- Ferguson, H.W., Collins, R.O., Moore, M., Coles, M. and MacPhee, D.D. 2004. *Pseudomonas anguilliseptica* infection in farmed cod, *Gadus morhua* L. *Journal of Fish Diseases* 27: 249-253.
- Ferguson, H.W., Kongtorp, R.T., Taksdal, T., Graham, D. and Falk, K. 2005. An outbreak of disease resembling heart and skeletal muscle inflammation in Scottish farmed salmon, *Salmo salar* L., with observations on myocardial regeneration. *Journal of Fish Diseases* 28: 119-123.
- Fouz, B., Zarza, C. and Amaro, C. 2006. First description of non-motile *Yersinia ruckeri* serovar I strains causing disease in rainbow trout, *Oncorhynchus mykiss* (Walbaum), cultured in Spain. *Journal of Fish Diseases* 29: 339-346.
- Fringuelli, E., Rowley, H.M., Wilson, J.C., Hunter, R., Rodger, H. and Graham, D.A. 2008. Phylogenetic analyses and molecular epidemiology of European salmonid alphaviruses (SAV) based on partial E2 and nsP3 gene nucleotide sequences. *Journal of Fish Diseases* 31: 811-823.

- Fritsvold, C., Kongtorp, R.T., Taskdal, T., Ørpetveit, I., Heum, M. and Poppe, T.T. 2009. Experimental transmission of cardiomyopathy syndrome (CMS) in Atlantic salmon *Salmo salar*. *Diseases of Aquatic Organisms* 87: 225-234.
- Furones, M.D., Rodgers, C.J. and Munn, C.B. 1993. *Yersinia ruckeri*, the causal agent of enteric redmouth disease (ERM) in fish. *Annual Review of Fish Diseases* 3: 105-125.
- Garseth, A.H., Biering, E. and Tengs, T. 2012. Piscine myocarditis virus (PMCV) in wild Atlantic salmon *Salmo salar*. *Diseases of Aquatic Organisms* 102: 157-161.
- Garseth, A.H., Fritsvold, C., Opheim, M., Skjerve, E. and Biering, E. 2013. Piscine reovirus (PRV) in wild Atlantic salmon, *Salmo salar* L., and sea-trout, *Salmo trutta* L., in Norway. *Journal of Fish Diseases* 36: 483-493.
- Garver, K.A., Traxler, G.S., Hawley, L.M., Richard, J., Ross, J.P. and Lovy, J. 2013. Molecular epidemiology of viral haemorrhagic septicaemia virus (VHSV) in British Columbia, Canada, reveals transmission from wild to farmed fish. *Diseases of Aquatic Animals* 104: 93-104.
- Godoy, M.G., Aedo, A., Kibenge, M.J.T., Groman, D.B., Yason, C.V., Grothusen, H., Lisperguer, A., Calbucura M., Avendaño, F., Imilán, M., Jarpa, M. and Kibenge, F.S.B. 2008. First detection, isolation and molecular characterisation of infectious salmon anaemia virus associated with clinical disease in farmed Atlantic salmon (*Salmo salar*) in Chile. *BMC Veterinary Research* 4: 28.
- Godoy, M.G., Kibenge, M.J.T., Suarez, R., Lazo, E., Heisinger, A., Aguinaga, J., Bravo, D., Mendoza, J., Llegues, K.O., Avendaño-Herrera, R., Vera, C., Mardones, F. and Kibenge, F.S.B. 2013. Infectious salmon anaemia virus (ISAV) in Chilean Atlantic salmon (*Salmo salar*) aquaculture: emergence of low pathogenic ISAV-HPR0 and re-emergence of virulent ISAV-HPRΔ: HPR3 and HPR14. *Virology Journal* 10: 344.
- Godoy, M.G., Kibenge, M.J.T., Wang, Y., Suarez, R., Leiva, C., Vallejos, F. and Kibenge, F.S.B. 2016. First description of clinical presentation of piscine orthoreovirus (PRV) infections in salmonid aquaculture in Chile and identification of a second genotype (Genotype II) of PRV. *Virology Journal* 13:98.
- Graham, D.A., Jewhurst, V.A., Rowley, H.M., McLoughlin, M.F. and Todd, D. 2003. A rapid immunoperoxidase-based virus neutralization assay for salmonid alphavirus used for a serological survey in Northern Ireland. *Journal of Fish Diseases* 26: 407-413.
- Graham, D.A., Cherry, K., Wilson, C.J. and Rowley, H.M. 2007a. Susceptibility of salmonid alphavirus to a range of chemical disinfectants. *Journal of Fish Diseases* 30: 269-277.
- Graham, D.A., Rowley, H.M., Fringuelli, E., Bovo, G., Manfrin, A., McLoughlin, M.F., Zarza, C., Khalili, M. and Todd, D. 2007b. First laboratory confirmation of salmonid alphavirus infection in Italy and Spain. *Journal of Fish Diseases* 30: 569-572.
- Graham, D.A., Fringuelli, E., Rowley, H.M., Cockerill, D., Cox, D.I., Turnbull, T., Rodger, H., Morris, D. and McLoughlin, M.F. 2012. Geographical distribution of salmonid alphavirus subtypes in marine farmed Atlantic salmon, *Salmo salar* L., in Scotland and Ireland. *Journal of Fish Diseases* 35: 755-765.
- Gustafson, L.L., Ellis, S.K., Beattie, M.J., Chang, B.D., Dickey, D.A., Robinson, T.L., Marengi, F.P., Moffett, P.J. and Page, F.H. 2007. Hydrographics and the timing of infectious salmon anemia outbreaks among Atlantic salmon (*Salmo salar* L.) farms in the Quoddy region of Maine, USA and New Brunswick, Canada. *Preventive Veterinary Medicine* 78: 35-56.
- Haenan, O.L.M. and Davidse, A. 2001. First isolation and pathogenicity studies with *Pseudomonas anguilliseptica* from diseased European eel *Anguilla anguilla* in The Netherlands. *Aquaculture* 196: 27-36.
- Haugland, Ø., Mikalsen, A.B., Nilsen, P., Lindmo, K., Thu, B.J., Eliassen, T.M., Roos, N., Rode, M. and Evensen, Ø. 2011. Cardiomyopathy syndrome of Atlantic salmon (*Salmo salar* L.) is caused by a double-stranded RNA virus of the *Totiviridae* family. *Journal of Virology* 85: 5275-5286.

- Hellberg, H., Kvellestad, A., Dannevig, B., Bornø, G., Modahl, I., Haldorsen, R.N., Vik-Mo, F., Ottesen, K., Saetre, E.M. and Sindre, H. 2010. Outbreaks of viral nervous necrosis in juvenile and adult farmed Atlantic cod, *Gadus morhua* L., in Norway. *Journal of Fish Diseases* 33: 75-81.
- Hill, B.J. and Way, K. 1995. Serological classification of infectious pancreatic necrosis (IPN) virus and other aquatic birnaviruses. *Annual Review of Fish Diseases* 5: 55-77.
- Hjeltnes, B., Walde, C.S., Bang Jensen, B. and Haukaas, A. 2016. Fish health report 2015. Oslo, Norwegian Veterinary Institute, 76 pp.
- Hjortaas, M.J., Skjelstad, H.R., Taksdal, T., Olsen, A.B., Johansen, R., Bang-Jensen, B., Ørpetveit, I. And Sindre, H. 2013. The first detections of subtype 2-related salmonid alphavirus (SAV2) in Atlantic salmon, *Salmo salar* L., in Norway. *Journal of Fish Diseases* 36, 71-74.
- Hodneland, K. and Endresen, C. 2006. Sensitive and specific detection of *Salmonid alphavirus* using real-time PCR (TaqMan®). *Journal of Virological Methods* 131: 184-192.
- Holt, R.A., Rohovec, J.S. and Fryer, J.L. 1993. Bacterial coldwater disease. In *Bacterial diseases of fish* (eds. Inglis V, Roberts RJ and Bromage NR). Blackwell Scientific Publications, Oxford. pp 3-23.
- ICES. 2001. Report of the Working Group on Pathology and Diseases of Marine Organisms. ICES CM 2001/F:02.
- ICES. 2012. Report of the Working Group on Pathology and Diseases of Marine Organisms. ICES CM 2012/SSGHIE:03. 68pp.
- ICES. 2016. Interim report of the working group on pathology and diseases of marine organisms (WGPDMO). ICES CM 2016/SSGEPI:07. 18 pp.
- Jansen, M.D., Wasmuth, M.A., Olsen, A.B., Gjerset, B., Modahl, I., Breck, O., Haldorsen, R.N., Hjelmeland, R. and Taksdal, T. 2010. Pancreas disease (PD) in sea-reared Atlantic salmon, *Salmo salar* L., in Norway; a prospective, longitudinal study of disease development and agreement between diagnostic test results. *Journal of Fish Diseases* 33: 723-736.
- Johansen, R., Sommerset, I., Tørud, B., Korsnes, K., Hjortaas, M.J., Nilsen, F., Nerland, A.H. and Dannevig, B.H. 2004. Characterisation of nodavirus and viral encephalopathy and retinopathy in farmed turbot, *Scophthalmus maximus* (L.). *Journal of Fish Diseases* 27: 591-601.
- Johnson, S.C., Sperker, S.A., Leggiadro, C.T., Groman, D.B., Griffiths, S.G., Ritchie, R.J., Cook, M.D. and Cusack, R.R. 2002. Identification and characterization of a piscine neuropathy and nodavirus from juvenile Atlantic cod from the Atlantic coast of North America. *Journal of Aquatic Animal Health* 14: 124-133.
- Karlsbakk, E., Olsen, A.B., Einen, A.C.B., Mo, T.A., Fiksdal, I.U., Aase, H., Kalgraff, C., Skår, S.Å., Hansen, H. 2013. Amoebic gill disease due to *Paramoeba perurans* in ballan wrasse (*Labrus bergylta*). *Aquaculture* 412-413: 41-44.
- Kent, M.L., Sawyer, T.K., Hedrick, R.P. 1988. *Paramoeba pemaquidensis* (Sarcomastigophora: Paramoebidae) infestation of the gill of coho salmon *Oncorhynchus kisutch* reared in sea water. *Diseases of Aquatic Organisms* 5:163-169.
- Kibenge, F.S.B., Kibenge, M.J.T., Wang, Y., Qian, B., Hariharan, S. and McGeachy, S. 2007. Mapping of putative virulence motifs on infectious salmon anaemia virus surface glycoprotein genes. *Journal of General Virology* 88: 3100-3111.
- Kongtorp, R.T., Kjerstad, A., Taksdal, T., Guttvik, A. and Falk, K. 2004a. Heart and skeletal muscle inflammation in Atlantic salmon, *Salmo salar* L.: a new infectious disease. *Journal of Fish Diseases* 27: 351-358.
- Kongtorp, R.T., Taksdal, T. and Lyngøy, A. 2004b. Pathology of heart and skeletal muscle inflammation (HSMI) in farmed Atlantic salmon *Salmo salar*. *Diseases of Aquatic Organisms* 59: 217-224.

- Lester, K., Black, J. and Bruno, D.W. 2011. Prevalence of salmonid alphavirus in Scottish fish farms from 2006 to 2007. *Bulletin of the European Association of Fish Pathologists* 31: 199-204.
- Lönnström, L., Wiklund, T. and Bylund, G. 1994. *Pseudomonas anguilliseptica* isolated from Baltic herring (*Clupea harengus membras*) with eye lesions. *Diseases of Aquatic Organisms* 18: 143-147.
- López-Romalde, S., Magariños, B., Ravelo, C., Toranzo, A.E. and Romalde, J.L. 2003. Existence of two O-serotypes in the fish pathogen *Pseudomonas anguilliseptica*. *Veterinary Microbiology* 94, 325-333.
- Løvoll, M., Alarcon, M., Bang Jensen, B., Taksdal, T., Kristoffersen, A.B. and Tengs, T. 2012 Quantification of piscine reovirus (PRV) at different stages of Atlantic salmon *Salmo salar* production. *Diseases of Aquatic Organisms* 99: 7-12.
- Lyngstad, T.M., Jansen, P.A., Sindre, H., Jonassen, C.M., Hjortaas, M.J., Johnsen, S. and Brun, E. 2008. Epidemiological investigation of infectious salmon anaemia (ISA) outbreaks in Norway 2003-2005. *Preventative Veterinary Medicine* 84: 213-227.
- Madetoja, J., Nystedt, S. and Wiklund, T. 2003. Survival and virulence of *Flavobacterium psychrophilum* in water microcosms. *FEMS Microbiology Ecology* 43: 217-223.
- Madsen, L. and Dalsgaard, I. 2008. Water recirculation and good management; potential methods to avoid disease outbreaks with *Flavobacterium psychrophilum*. *Journal of Fish Diseases* 31: 799-810.
- Madsen, L., Bertelsen, S.K., Dalsgaard, I. and Middelboe, M. 2013. Dispersal and survival of *Flavobacterium psychrophilum* phages *in vivo* in rainbow trout and *in vitro* under laboratory conditions: implications for their use in phage therapy. *Applied and Environmental Microbiology* 79: 4853-4861.
- Magi, G.E., Lopez-Romalde, S., Magariños, G.E., Lamas, J., Toranzo, A.E. and Romalde, L. 2009. Experimental *Pseudomonas anguilliseptica* infection in turbot *Psetta maxima* (L.): a histopathological and immunohistochemical study. *European Journal of Histochemistry* 53:e9.
- McArdle, J. 2014. Enteric redmouth disease in rainbow trout: an affair of the heart? *Veterinary Record* 174: 386.
- McBeath, A.J.A., Bain, N. and Snow, M. 2009. Surveillance for infectious salmon anaemia virus HPR0 in marine Atlantic salmon farms across Scotland. *Diseases of Aquatic Organisms* 87: 161-169.
- McCleary, S., Giltrap, M., Henshilwood, K. and Ruane, N.M. 2014. Detection of salmonid alphavirus RNA in Celtic and Irish Sea flatfish. *Diseases of Aquatic Organisms* 109: 1-7.
- McLoughlin, M.F. and Graham, D.A. 2007. Alphavirus infections in salmonids – a review. *Journal of Fish Diseases* 30: 511-531.
- McLoughlin, M.F., Nelson, R.N., McCormick, J.I., Rowley, H.M. and Bryson, D.B. 2002. Clinical and histopathological features of naturally occurring pancreas disease in farmed Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases* 25: 33-43.
- Mitchell, S.O., Rodger, H.D. 2011. A review of infectious gill disease in marine salmonid fish. *Journal of Fish Diseases* 34:411-432.
- Moen, T., Baranski, M., Sonesson, A.K., Kjøglum, S. 2009. Confirmation and fine-mapping of a major QTL for resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*): population-level associations between markers and trait. *BMC Genomics* 10: 368.
- Morrison, R.N., Crosbie, P.B.B., Nowak, B.F. 2004. The induction of laboratory-based amoebic gill disease revisited. *Journal of Fish Diseases* 27:445-449.
- Mouton, A., Crosbie, P., Cadoret, K., Nowak, B. 2014. First record of amoebic gill disease caused by *Neoparamoeba perurans* in South Africa. *Journal of Fish Diseases* 37:407-409.
- Munday, B.L., Lange, K., Foster, C., Lester, R., Handler, J. 1993. Amoebic gill disease in sea-caged salmonids in Tasmanian waters. *Tasmanian Fisheries Research* 28:14-19.

- Munday, B.L., Zilberg, D., Findlay, V. 2001. Gill disease of marine fish caused by infection with *Neoparamoeba pemaquidensis*. *Journal of Fish Diseases* 24:497-507.
- Munday, B.L., Kwang, J. and Moody, N. 2002. Betanodavirus infections of teleost fish – a review. *Journal of Fish Diseases* 25: 127-142.
- Munro, E.S., McIntosh, R.E., Weir, S.J., Noguera, P.A., Sandilands, J.M., Matejusova, I., Mayes, A.S. and Smith, R. 2015. A mortality event in wrasse species (Labridae) associated with the presence of viral haemorrhagic septicaemia virus. *Journal of Fish Diseases* 38: 335-341.
- Murray, A.G., Munro, L.A., Wallace, I.S., Berx, B., Pendrey, D., Fraser, D. and Raynard, R.S. 2010. Epidemiological investigation into the re-emergence and control of an outbreak of infectious salmon anaemia in the Shetland Islands, Scotland. *Diseases of Aquatic Organisms* 91: 189-200.
- Nakai, T., Muroga, K., Chung, H.-Y. and Kou, G.-H. 1985. A serological study on *Pseudomonas anguilliseptica* isolated from diseased eels in Taiwan. *Fish Pathology* 19: 256-261.
- Nematollahi, A., Decostere, A., Pasmans, F. and Haesebrouck, F. 2003. *Flavobacterium psychrophilum* infections in salmonid fish. *Journal of Fish Diseases* 26: 563-574.
- Nilsen, H., Johansen, R., Colquhoun, D.C., Kaada, I., Bottolfsen, K., Vågnes, Ø., and Olsen, A.B. 2011a. *Flavobacterium psychrophilum* associated with septicaemia and necrotic myositis in Atlantic salmon, *Salmo salar*: a case report. *Diseases of Aquatic Organisms* 97: 37-46.
- Nilsen, H., Olsen, A.B., Vågnes, Ø., Hellberg, H., Bottolfsen, K., Skjelstad, H. and Colquhoun, D.J. 2011b. Systemic *Flavobacterium psychrophilum* infection in rainbow trout, *Oncorhynchus mykiss* (Walbaum), farmed in fresh and brackish water in Norway. *Journal of Fish Diseases* 34: 403-408.
- Nishizawa, T., Kinoshita, S. and Yoshimizu, M. 2005. An approach for genogrouping of Japanese isolates of aquabirnaviruses in a new genogroup, VII, based on the VP2/NS junction region. *Journal of General Virology* 86: 1973-1978.
- Norris, A., Foyle, L. and Ratcliff, J. 2008. Heritability of mortality in response to a natural pancreas disease (SPDV) challenge in Atlantic salmon, *Salmo salar* L., post-smolts on a West of Ireland sea site. *Journal of Fish Diseases* 31: 913-920.
- OIE. 2012. <http://www.oie.int/en/international-standard-setting/specialists-commissions-groups/aquatic-animal-commission-reports/>
- Oldham, T., Rodger, H. And Nowak, B.F. 2016. Incidence and distribution of amoebic gill disease (AGD) – an epidemiological review. *Aquaculture* 457: 35-42.
- Olsen, A.B., Mikalsen, J., Rose, M., Alfjorden, A., Hoel, E., Straum-Lie, K., Haldorsen, R. and Colquhoun, D.J. 2006. A novel systemic granulomatous inflammatory disease in farmed Atlantic cod, *Gadus morhua* L., associated with a bacterium belonging to the genus *Francisella*. *Journal of Fish Diseases* 29: 307-311.
- Olsen, A.B., Hjortaas, M., Tengs, T., Hellberg, H. and Johansen, R. 2015. First description of a new disease in rainbow trout (*Oncorhynchus mykiss* (Walbaum)) similar to heart and skeletal muscle inflammation (HSMI) and detection of a gene sequence related to piscine orthoreovirus (PRV). *PLoS ONE* 10(7): e0131638.
- Ottem, K.F., Nylund, A., Karlsbakk, E., Friis-Møller, A. and Kamaishi, T. 2009. Elevation of *Francisella philomiragia* subsp. *noatunensis* Mikalsen et al. (2007) to *Francisella noatunensis* comb. nov. [syn. *Francisella piscida* Ottem et al. (2008) syn. nov.] and characterisation of *Francisella noatunensis* subsp. *orientalis* subsp. nov., two important fish pathogens. *Journal of Applied Microbiology* 106: 1231-1243.
- Ottem, K.F., Nylund, A., Isaksen, T.E., Karlsbakk, E. and Bergh, Ø. 2008. Occurrence of *Francisella piscida* in farmed and wild Atlantic cod, *Gadus morhua* L., in Norway. *Journal of Fish Diseases* 31: 525-534.

- Øye, A.K. and Rimstad, E. 2001. Inactivation of infectious salmon anaemia virus, viral haemorrhagic septicaemia virus and infectious pancreatic necrosis virus in water using UVC irradiation. *Diseases of Aquatic Organisms* 48: 1-5.
- Palacios, G., Lovoll, M., Tengs, T., Hornig, M., Hutchison, S., Hui, J., Kongtorp, R.T., Savji, N., Bussetti, A.V., Solovyov, A., Kristoffersen, A.B., Celone, C., Street, C., Trifonov, V., Hirschberg, D.L., Rabadan, R., Egholm, M., Rimstad, E. and Lipkin, W.I. 2010. Heart and skeletal muscle inflammation of farmed salmon is associated with infection with a novel reovirus. *PLoS ONE* 5:7.
- Palmer, R., Carson, J., Rutledge, M., Drinan, E., Wagner, T. 1997. Gill disease associated with *Paramoeba*, in sea reared Atlantic salmon in Ireland. *Bulletin of the EAAP* 17:112-114.
- Plarre, H., Devold, M., Snow, M. And Nylund, A. 2005. Prevalance of infectious salmon anaemia virus (ISAV) in wild salmonids in western Norway. *Diseases of Aquatic Organisms* 66: 71-79.
- Raja-Halli, M., Vehmas, T.K., Rimaila-Pärnänen, E., Sainmaa, S., Skall, H.F., Olesen, N.J. and Tapiovaara, H. 2006. Viral haemorrhagic septicaemia (VHS) outbreaks in Finnish rainbow trout farms. *Diseases of Aquatic Organisms* 72: 201-211.
- Raynard, R.S., Murray, A.G. and Gregory, A. 2001. Infectious salmon anaemia virus in wild fish from Scotland. *Diseases of Aquatic Organisms* 46: 93-100.
- Roberts, R.J. 2012. Fish Pathology. 4th ed. Blackwell publishing, Chichester, UK.
- Roberts, R.J. and Pearson, M.D. 2005. Infectious pancreatic necrosis in Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases* 28: 383-390.
- Rodger, H.D. 2014. Amoebic gill disease (AGD) in farmed salmon (*Salmo salar*) in Europe. *Fish Veterinary Journal* 14: 16-27.
- Rodger, H. and Mitchell, S. 2007. Epidemiological observations of pancreas disease of farmed Atlantic salmon, *Salmo salar* L., in Ireland. *Journal of Fish Diseases* 30: 157-167.
- Rodger, H.D., McCleary, S.J. and Ruane, N.M. 2014. Clinical cardiomyopathy syndrome in Atlantic salmon, *Salmo salar* L. *Journal of Fish Diseases* 37: 935-939.
- Ruane, N.M., McCarthy, L.J., Swords, D. and Henshilwood, K. 2009. Molecular differentiation of infectious pancreatic necrosis virus isolates from farmed and wild salmonids in Ireland. *Journal of Fish Diseases* 32: 979-987.
- Ruane, N.M., McCleary, S.J., McCarthy, L.J. and Henshilwood, K. 2015. Phylogenetic analysis of infectious pancreatic necrosis virus in Ireland reveals the spread of a virulent genogroup 5 subtype previously associated with imports. *Archives of Virology* 160: 817-824.
- Ruane, N.M., Bolton-Warberg, M., Rodger, H.D., Colquhoun, D.J., Geary, M., McCleary, S.J., O'Halloran, K., Maher, K., O'Keeffe, D., Mirimin, L., Henshilwood, K., Geoghegan, F. and Fitzgerald, R.D. 2015. An outbreak of francisellosis in wild-caught Celtic Sea Atlantic cod, *Gadus morhua* L., juveniles reared in captivity. *Journal of Fish Diseases* 38: 97-102.
- Siah, A., Morrison, D.B., Fringuelli, E., Savage, P., Richmond, Z., Johns, R., Purcell, M.K., Johnson, S.C. and Saksida, S.M. 2015. Piscine reovirus: genomic and molecular phylogenetic analysis from farmed and wild salmonids collected on the Canada/US Pacific coast. *PLoS ONE* 10(11): e0141475.
- Skall, H.F., Olesen, N.J. and Møllergaard, S. 2005. Viral haemorrhagic septicaemia virus in marine fish and its implications for fish farming – a review. *Journal of Fish Diseases* 28: 509-529.
- Smail, D.A., Huntly, P.J. and Munro, A.L.S. 1993. Fate of four fish pathogens after exposure to fish silage containing fish farm mortalities and conditions for the inactivation of infectious pancreatic necrosis virus. *Aquaculture* 113: 173-181.
- Smail, D.A., Grant, R., Simpson, D., Bain, N. and Hastings, T.S. 2004. Disinfectants against cultured infectious salmon anaemia (ISA) virus: the virucidal effect of three iodophors, chloramines T, chlorine dioxide and peracetic acid/hydrogen peroxide/acetic acid mixture. *Aquaculture* 240: 29-38.

- Snow, M., Bain, N., Black, J., Taupin, V., Cunningham, C.O., King, J.A., Skall, H.F. and Raynard, R.S. 2004. Genetic population structure of marine viral haemorrhagic septicaemia virus (VHSV). *Diseases of Aquatic Organisms* 61: 11-21.
- Snow, M., Black, J., Matejusova, I., McIntosh, R., Baretto, E., Wallace, I.S. and Bruno, D.W. 2010. Detection of salmonid alphavirus RNA in wild marine fish: implications for the origins of salmon pancreas disease in aquaculture. *Diseases of Aquatic Organisms* 91: 177-188.
- Steinum, T., Kvellestad, A., Rønneberg, L.B., Nilsen, H., Asheim, A., Fjell, K., Nygård, S.M.R., Olsen, A.B., Dale, O.B. 2008. First cases of amoebic gill disease (AGD) in Norwegian sea-water farmed Atlantic salmon, *Salmo salar* L., and phylogeny of the causative amoeba using 18S cDNA sequences. *Journal of Fish Diseases* 31:205-214.
- Stewart, D.J., Waldemariam, K., Dear, G. and Mochaba, F.M. 1983. An outbreak of "Sekiten-byo" among cultured European eels, *Anguilla Anguilla* L., in Scotland. *Journal of Fish Diseases* 6: 75-76.
- Stone, D.M., Ferguson, H.W., Tyson, P.A., Savage, J., Wood, G., Dodge, M.J., Woolford, G., Dixon, P.F., Feist, S.W. and Way, K. 2008. The first report of viral haemorrhagic septicaemia in farmed rainbow trout, *Oncorhynchus mykiss* (Walbaum), in the United Kingdom. *Journal of Fish Diseases* 31: 775-784.
- Studer, J. and Janies, D.A. 2011. Global spread and evolution of viral haemorrhagic septicaemia virus. *Journal of Fish Diseases* 34: 741-747.
- Taksdal, T., Olsen, A.B., Bjerkås, I., Hjortaas, M.J., Dannevig, B.H., Graham, D.A. and McLoughlin, M.F. 2007. Pancreas disease in farmed Atlantic salmon, *Salmo salar* L., and rainbow trout, *Oncorhynchus mykiss* (Walbaum), in Norway. *Journal of Fish Diseases* 30: 545-558.
- Taylor, R.S., Muller, W.J., Cook, M.T., Kube, P.D. and Elliott, N.G. 2009. Gill observations in Atlantic salmon (*Salmo salar*) during repeated amoebic gill disease (AGD) field exposure and survival challenge. *Aquaculture* 290: 1-8.
- Tobback, E., Decostere, A., Hermans, K., Haesebrouck, F. and Chiers, K. 2007. *Yersinia ruckeri* infections in salmonid fish. *Journal of Fish Diseases* 30: 257-268.
- Vardić Smrzlić, I., Kapetanović, D., Valić, D., Teskeredžić, E., McLoughlin, M.F. and Fringuelli, E. 2013. First laboratory confirmation of sleeping disease virus (SDV) in Croatia. *Bulletin of the EAAP* 33: 78-83.
- Vincent, B.N., Adams, M.B., Crosbie, P.B.B., Nowak, B.F., Morrison, R.N. 2007. Atlantic salmon (*Salmo salar* L.) exposed to cultured gill-derived *Neoparamoeba branchiphila* fail to develop amoebic gill disease (AGD). *Bulletin of the EAAP* 27:112-115.
- Wakabayashi, H. and Egusa, S. 1972. Characteristics of a *Pseudomonas* sp. from an epizootic of pond-cultured eel (*Anguilla japonica*). *Bulletin of the Japanese Society of Scientific Fisheries* 38: 577-587.
- Wheeler, R.W., Davies, R.L., Dalsgaard, I., Garcia, J., Welch, T.J., Wagley, S., Bateman, K.S. and Verner-Jeffreys, D.W. 2009. *Yersinia ruckeri* biotype 2 isolates from mainland Europe and the UK likely represents different clonal groups. *Diseases of Aquatic Organisms* 84: 25-33.
- Wiik-Nielsen, C.R., Løvoll, M., Sandlund, N., Faller, R., Wiik-Nielsen, J. and Bang Jensen, B. 2012a. First detection of piscine reovirus (PRV) in marine fish species. *Diseases of Aquatic Organisms* 97: 255-258.
- Wiik-Nielsen, C.R., Ski, P.-M.R., Aunsmo, A. and Løvoll, M. 2012b. Prevalence of viral RNA from piscine reovirus and piscine myocarditis virus in Atlantic salmon, *Salmo salar* L., broodfish and progeny. *Journal of Fish Diseases* 35: 169-171.
- Wiklund, T. and Bylund, J. 1990. *Pseudomonas anguilliseptica* as a pathogen of salmonid fish in Finland. *Diseases of aquatic Organisms* 8: 13-19.
- Wiklund, T. and Lönnström, L. 1994. Occurrence of *Pseudomonas anguilliseptica* in Finnish fish farms during 1986-1991. *Aquaculture* 126: 211-217.

- Wong, F.Y.K., Carson, J., Elliott, N.G. 2004. 18S ribosomal DNA-based PCR identification of *Neoparamoeba pemaquidensis*, the agent of amoebic gill disease in sea-farmed salmonids. *Diseases of Aquatic Organisms* 60:65-76.
- Young, N.D., Dyková, I., Snekvik, K., Nowak, B.F., Morrison, R.N. 2008. *Neoparamoeba perurans* is a cosmopolitan aetiological agent of amoebic gill disease. *Diseases of Aquatic Organisms* 78:217-223.
- Zerihun, M.A., Feist, S.F., Bucke, D., Olsen, A.B., Tandstad, N.M. and Colquhoun, D.J. 2011. *Francisella noatunensis* subsp. *noatunensis* is the aetiological agent of visceral granulomatosis in wild Atlantic cod *Gadus morhua*. *Diseases of Aquatic Organisms* 95: 65-71.

5 Viral Diseases in Farmed Molluscs

5.1 Ostreid herpesvirus 1 in bivalves

5.1.1 Description of Agent

Ostreid herpesvirus 1 (OsHV-1) is the aetiological agent of a contagious viral disease affecting bivalve species including the Pacific oyster, *Crassostrea gigas*. OsHV-1 particles have been purified from French *C. gigas* larvae (Le Deuff and Renault, 1999) and were observed by transmission electron microscopy to be enveloped icosahedral with electron dense cores and a diameter around 120 nm. The entire virus DNA was sequenced and OsHV-1 capsids appear structurally similar to those of other herpesviruses that have been studied (Davison *et al.*, 2005). The virus was classified under the name Ostreid herpesvirus 1 (OsHV-1) as the first known species in the family Malacoherpesviridae, order Herpesvirales.

Although the aetiological agent is represented by all specimens of OsHV-1 (Arzul *et al.*, 2001; Davison *et al.*, 2005; Moss *et al.*, 2007; Segarra *et al.*, 2010; Martenot *et al.*, 2011; Renault *et al.*, 2012), increased mortality outbreaks recently reported in Europe, Australia and New Zealand among *C. gigas* spat in association with all OsHV-1 μ Var viral variants suggest differences in terms of virulence among OsHV-1 lineages. However, the detection of variants related to OsHV-1 μ Var have also been reported in the absence of mortality events (Dundon *et al.*, 2011; Shimahara *et al.*, 2012) suggesting the involvement of several factors in disease expression.

5.1.2 Geographical Distribution and Temporal Trends

OsHV-1 representatives have been reported in Europe (France, Ireland, Italy, the Netherlands, Spain, Sweden, United Kingdom), Australia, Brazil, China (People's Rep. of), Korea, Japan, Morocco, Tunisia, Mexico, New Zealand and the United States of America.

Since 2008, widespread mortality was reported among *C. gigas* stocks in different Member States of the European Union in association with emergence of the hypervirulent OSHV-1 μ Var variant (Segarra *et al.*, 2010; Peeler *et al.*, 2012; Roque *et al.*, 2012). The outbreaks are seasonal and highly temperature dependent with mortality rates of up to 100% (Clegg *et al.*, 2014). Mortality events attributed to OsHV-1 μ Var variants were also reported beginning in 2010 in Australia and New Zealand (Jenkins *et al.*, 2013).

5.1.3 Short Description of Clinical Signs

The virus can be found in adult bivalves in the absence of mortality. Infection-associated lesions in juveniles are mainly observed in connective tissues of all organs in which fibroblastic-like cells exhibit enlarged nuclei with perinuclear chromatin (Renault *et al.*, 1994; Schikorski *et al.*, 2011).

5.1.4 Control/Preventative Measures

The herpes infection affecting bivalves is not listed by the World Organisation for Animal Health (WOAH) (OIE) and the EU (Directive 2006/88/EC). Pacific cupped oyster families less susceptible to OsHV-1 including the variant OsHV-1 μ Var can be produced for aquaculture use (Sauvage *et al.*, 2009). Biosecurity may be successfully applied in confined and controlled facilities such as hatcheries and nurseries in order to protect the facility and the surrounding environment from the introduction of the virus.

5.1.5 Host Species

OsHV-1 infection causes mortality in larvae and juveniles of several bivalve species (Renault *et al.*, 1994; Garcia *et al.*, 2011) including the Pacific oyster, *C. gigas*, Portuguese oyster, *Crassostrea angulata*, suminoe oyster, *Crassostrea ariakensis*, European flat oyster, *Ostrea edulis*, Manila clam, *Ruditapes philippinarum*, carpet shell clam, *Ruditapes decussatus*, and great scallop, *Pecten maximus*. However, the variant μ Var (Segarra *et al.*, 2010) primarily infects *C. gigas*.

6 Bacterial Diseases of Farmed Molluscs

6.1 *Vibrio* sp. infecting marine molluscs

6.1.1 *Vibrio splendidus* infecting Pacific oysters

6.1.1.1 Description of Agent

Gram-negative bacteria related to *Vibrio splendidus* are frequently found in coastal areas and can infect and induce mortalities in the Pacific oyster, *C. gigas* (Lacoste *et al.*, 2001; Samain *et al.*, 2004; Saulnier *et al.*, 2010). Through epidemiological studies a high genetic diversity was observed in this group suggesting that the *splendidus* clade is diverse and possibly polyspecific (Le Roux *et al.*, 2002).

6.1.1.2 Geographical Distribution and Temporal Trends

Since 2008, widespread mortality events have been reported among *C. gigas* in France. The outbreaks are seasonal and most frequently affect spat and juveniles (< 18 months). Mortality rates of 40-100% were experienced. While the mortality in young oysters has generally been attributed to emergence of the OsHV-1 μ Var (EFSA, 2010; Segarra *et al.*, 2010), *V. splendidus* was frequently detected in affected, as well as unaffected, oysters (EFSA, 2010).

6.1.1.3 Short Description of Clinical Signs

The main sign of *V. splendidus* infections in the Pacific oyster remains reports of mortality events (Saulnier *et al.*, 2010). There are no reliable clinical indicators of *V. splendidus* infection in oysters.

6.1.1.4 Control/Preventative Measures

V. splendidus infection is not listed by the World Organisation for Animal Health (WOAH) and the EU (Directive 2006/88/EC).

6.1.1.5 Host species

V. splendidus-related species were reported in association with mortality outbreaks affecting different mollusc species (Sugumar *et al.*, 1998; Macian *et al.*, 2000; Lacoste *et al.*, 2001; Gay *et al.*, 2003; Gay *et al.*, 2004).

6.1.2 *Vibrio aestuarianus* infecting oysters

6.1.2.1 Description of Agent

The gram-negative bacterium *Vibrio aestuarianus* is frequently found in coastal areas and can infect and induce mortality outbreaks among the Pacific oyster, *C. gigas* (Samain *et al.*, 2004; Garnier *et al.*, 2008; Saulnier *et al.*, 2010).

6.1.2.2 Geographical Distribution and Temporal Trends

V. aestuarianus has been reported to be associated with mortality of oyster reared in open marine waters in France. Some bacterial isolates related to this species were demonstrated to be pathogenic to *C. gigas* under experimental conditions (Garnier *et al.*, 2007; Azandegbe *et al.*, 2010).

6.1.2.3 Short Description of Clinical Signs

V. aestuarianus has frequently been among the pathogens associated with massive mortality events occurring during summer in *C. gigas* oysters. These events often occur

when seawater temperatures reach 19°C on the French Atlantic coast (Garnier *et al.*, 2007; Labreuche *et al.*, 2010). Classic bacteriology studies revealed that moribund animals were predominantly infected with *V. aestuarianus*, found in the hemolymph but also in other oyster tissues (Azandegbe *et al.*, 2010).

6.1.2.4 Control/Preventative Measures

V. aestuarianus infection is not listed by the World Organisation for Animal Health (WOAH) or the EU (Directive 2006/88/EC).

6.1.2.5 Host species

The gram-negative bacterium *V. aestuarianus* is mainly reported infecting the *C. gigas*. However, *V. aestuarianus* has also been recently detected in the cockle *Cerastoderma edule* in France in association with mortality outbreaks (C. Garcia *et al.*, IFREMER, pers. comm.).

6.1.3 *Vibrio harveyi* in abalone

6.1.3.1 Description of Agent

The gram-negative bacterium *Vibrio harveyi* is known to be highly pathogenic for the European abalone *Haliotis tuberculata*. Since 1998, specific strains of *V. harveyi* have been implicated in mortality outbreaks in French farms and field stocks of abalone (Nicolas *et al.*, 2002). *V. harveyi* has been widely recognized as a common pathogen of many commercially cultured fish and shellfish species worldwide (Gomez *et al.*, 2004) including abalone in Australia and Japan (Nishimori *et al.*, 1998; Handler *et al.*, 2005; Sawabe *et al.*, 2007).

6.1.3.2 Geographical Distribution and Temporal Trends

The bacterium has been involved in recurrent mortality outbreaks occurring seasonally, at the end of warm season, since 1998 in farms and field stocks of *H. tuberculata* in France (Nicolas *et al.*, 2002).

6.1.3.3 Short Description of Clinical Signs

Although non-specific, clinical signs of *V. harveyi* infection include a loss of muscular strength occurring concomitantly with the appearance of white pustules on the foot. Subsequently, diseased animals develop a fatal septicemia leading to up to 80% mortality within a few days to 3 weeks. Vibriosis outbreaks in *H. tuberculata* cultivated in France were shown to be driven by seawater temperature exceeding a 17°C threshold (Huchette and Clavier, 2004) and host physiology such as gametogenesis and reduced immune defense capacities (Travers *et al.*, 2008).

6.1.3.4 Control/Preventative Measures

V. harveyi infection is not listed by the World Organisation for Animal Health (WOAH) (OIE, 2012) and the EU (Directive 2006/88/EC).

6.1.3.5 Host species

Vibrio harveyi infects members of the genus *Haliotis*, but has been also reported in bivalve molluscs (Saulnier *et al.*, 2010).

6.2 *Nocardia crassostreae* in oysters

6.2.1 Description of Agent

Nocardia crassostreae, a gram-positive actinomycete, is the causative agent of Pacific Oyster Nocardiosis (PON). The cycle of *N. crassostreae* in the environment between oysters is unknown. As most *Nocardia* species are soil bacteria this may suggest that *N. crassostreae* is acquired from the environment as an opportunistic invader of live oysters (Bower *et al.*, 2005). The soil substrate could be a natural source of *N. crassostreae*.

6.2.2 Geographical Distribution and Temporal Trends

The disease has been reported since the late 1940s in Japan and the west coast of North America in association with *C. gigas* field mortality outbreaks (Friedman *et al.*, 1991; Bower *et al.*, 2005). The extent of associated mortalities has not been accurately measured but estimated at about 35% in some localities. In British Columbia (Canada), European flat oysters *O. edulis* cultured alongside infected *C. gigas* have been found infected by *N. crassostreae* but mortality rate is unknown.

Over the last decade, the geographical distribution of *N. crassostreae* has extended outside the North Pacific with the report of *N. crassostreae* in *C. gigas* from the Netherlands (Engelsma *et al.*, 2008) and in *Mytilus galloprovincialis* and *O. edulis* from Italy (Carella *et al.*, 2013).

6.2.3 Short Description of Clinical Signs

The bacterium can be found all year as bacterial foci primarily in gonad follicles, vesicular connective tissue, gills, heart and adductor muscle, but they can invade every tissue. Bacteria are usually associated with mortalities during the late summer and fall. *C. gigas* oysters experimentally infected showed clinical signs and mortality (Friedman *et al.*, 1991; Friedman *et al.*, 1998).

6.2.4 Control/Preventative Measures

N. crassostreae infection is not listed by the World Organisation for Animal Health (OIE) or the EU (Directive 2006/88/EC). Bottom culture of oysters may possibly expose oysters more to *N. crassostreae* than other types of culture and so may be avoided to potentially mitigate infection pressure where it is intense.

6.2.5 Host species

N. crassostreae causes infection in the oysters *C. gigas* and *O. edulis* and was recently described in the mussel *M. galloprovincialis* (Carella *et al.*, 2013).

6.3 *Candidatus xenohaliotis californiensis* in abalone

6.3.1 Description of Agent

Candidatus xenohaliotis californiensis is an intracellular bacterium (332 × 1550 nm in the bacillus form and an average of 1405 nm in the spherical morphotype) in the family Anaplasmataceae and is closely related to members of the genera *Ehrlichia*, *Anaplasma* and *Cowdria* (Friedman *et al.*, 2000). Infection has been reported in wild and farmed abalone, *Haliotis* spp. (Archeogastropoda: Mollusca) (Gardner *et al.*, 1995; Friedman *et al.*, 2000). The disease caused by this bacterium is known as withering syndrome and may be more appropriately termed abalone rickettsiosis.

6.3.2 Geographical Distribution and Temporal Trends

Candidatus xenohaliotis californiensis occurs along the south-west coast of North America in California, USA and Baja California, Mexico. However, as infected abalone have been transported to Chile, Japan, Israel, Iceland and possibly other countries, a large geographical distribution of the bacterium is suspected. Infections have resulted in severe economic impacts to abalone culturists along the west coast of North America. Reduced profits have been associated with farm closures as well as contributing to the closure of commercial fisheries in California. Moreover, recurring disease outbreaks are also implicated in failure of wild abalone to repopulate historic habitats (Friedman and Finley, 2003). *Candidatus xenohaliotis californiensis* has now been detected in Europe including Ireland, Spain and France in the European abalone, *H. tuberculata* (Balseiro *et al.*, 2006).

6.3.3 Short Description of Clinical Signs

Clinical disease is typically observed in abalone over 12 months of age. Clinical disease has only been observed in infected individuals exposed to elevated seawater temperatures (e.g. 18°C). Gross signs of the infection include pedal atrophy, mottled digestive gland, anorexia, weakness, and lethargy before death. Associated losses may reach 99% of the population and depending on seawater temperatures and host species.

6.3.4 Control/Preventative Measures

The most effective prevention is avoidance of the pathogen. Infection caused by *Candidatus xenohaliotis californiensis* in molluscs is not listed as a notifiable disease under EU legislation (Directive 2006/88/EC). Infection is however included in the Manual of Diagnostic Tests for Aquatic Animals (www.oie.int) and an OIE Reference Laboratory for this pathogen has been designated. Should infection occur, holding abalone at temperatures below 15°C may reduce pathogen transmission and subsequent disease development. Application of oxytetracycline reduces losses.

6.3.5 Host species

Candidatus xenohaliotis californiensis infects members of the genus *Haliotis* and natural infections have been observed in black abalone (*H. cracherodii*), white abalone (*H. sorenseni*), red abalone (*H. rufescens*), pink abalone (*H. corrugata*), green abalone (*H. fulgens*), the small abalone (*H. diversicolor supertexta*), and the European abalone (*H. tuberculata*) in the wild or in culture facilities, as well as flat (*H. wallalensis*) and Japanese abalone (*H. discus-hannai*) in laboratory challenges. Other abalone species have not been tested.

7 Diseases of Mollusc: Parasitic Diseases

7.1 *Bonamia exitiosa*

7.1.1 Description of Agent

Bonamia exitiosa is a protozoan parasite in the Haplosporida, phylum Cercozoa (Cavalier-Smith and Chao, 2003), infecting haemocytes of several oyster species and inducing physiological disorders and eventually death of the animal (Dinamani *et al.*, 1987; Cranfield *et al.*, 2005). It is an intrahaemocytic protozoan, but it can be observed extracellularly (Dinamani *et al.*, 1987). This intrahaemocytic protozoan quickly becomes systemic and can be found in different organs, especially in connective tissues (Hine, 1991).

7.1.2 Geographical Distribution and Temporal Trends

Infection with *B. exitiosa* is found in oyster *Ostrea chilensis* in the Foveaux Strait and other locations around South Island, New Zealand (Dinamani *et al.*, 1987); and in *Ostrea angasi* and *Saccostrea glomerata* in New South Wales, Victoria, Tasmania, and Western Australia (Hine and Jones, 1994; Hine, 1996; Corbeil *et al.*, 2006; Carnegie *et al.*, 2014).

More recently, infection with *B. exitiosa* was reported in *O. edulis* in Galicia (Spain) (Abollo *et al.*, 2008), in the Adriatic Sea in Italy (Narcisi *et al.*, 2010), in the Mediterranean Sea in France and in Cornwall in the United Kingdom as well as in *Ostrea stentina* in Tunisia (Hill *et al.*, 2010). Molecular analyses have confirmed the presence of the parasite in the Americas as well, infecting *O. stentina* (= *Ostreola equestris*) along the southeastern Atlantic coast of the USA, *Ostrea lurida* in California, USA, and *Ostrea puelchana* in Argentina (Hill *et al.*, 2014).

7.1.3 Short Description of Clinical Signs

Infection can be lethal. In *O. chilensis*, death usually occurs as infections peak in intensity, particularly in association with high intensity apicomplexan infections (Hine and Wesney, 1994; Hine *et al.*, 2002). In one episode the disease killed more than 80% of *O. chilensis* as a wave of infection passed through an oyster bed over 2–3 years (Cranfield *et al.*, 2005). The impact of *B. exitiosa* in *O. edulis* or *O. stentina* has not yet been evaluated.

In *O. angasi*, the parasite is epitheliotropic, and apparently very light infections may cause a massive focal haemocyte infiltration with necrotic foci. In *O. edulis*, the parasite is associated with heavy haemocytic infiltration and appears in the connective tissue of different organs mostly within haemocytes, but sometimes outside host cells (Abollo *et al.*, 2008). In *O. stentina*, marked haemocytosis was not observed in animals found to be infected with the parasite (Hill *et al.*, 2010).

7.1.4 Control/Preventative Measures

Infection caused by *B. exitiosa* in molluscs is listed as a notifiable disease by the EU legislation (Directive 2006/88/EC). The infection is also listed by the OIE and included in the Manual of Diagnostic Tests for Aquatic Animals (www.oie.int).

It has been considered that in New Zealand development of lighter dredges and less damaging fishing strategies may reduce the chance of disease outbreaks by lowering disturbance (Cranfield *et al.*, 2005). Avoiding stressors such as exposure to extreme temperatures (below 7 or above 26°C) and salinity (40%), starvation, handling, or

heavy infection with other parasites, as well as decreasing density, should help to reduce the impact of the disease in that system (Cranfield *et al.*, 2005; Hine *et al.*, 2002). Control elsewhere should focus on preventing the introduction of *B. exitiosa* to areas in which it has not yet become established.

7.1.5 Host species

Oyster species *O. chilensis* (= *Tiostrea chilensis* = *T. lutaria*) (Dinamani *et al.*, 1987), *O. angasi* (Hine and Jones, 1994; Hine, 1996; Corbeil *et al.*, 2006), *O. edulis* (Abollo *et al.*, 2008; Narcisi *et al.*, 2010), *O. stentina* (Hill *et al.*, 2010) and *O. puelchana*, *O. lurida*, and *S. glomerata* (Hill *et al.*, 2014).

7.2 Bonamia ostreae

7.2.1 Description of Agent

Bonamia ostreae is a protozoan parasite in the Haplosporida (Pichot *et al.*, 1979; Comps *et al.*, 1980; Carnegie *et al.*, 2000), phylum Cercozoa (Cavalier-Smith and Chao, 2003), infecting haemocytes of flat oysters, *O. edulis*, and inducing physiological disorders and eventually death of the animal (Grizel, 1985). It is an intrahaemocytic protozoan, but it can be observed extracellularly.

7.2.2 Geographical Distribution and Temporal Trends

Infection with *B. ostreae* has been found in Europe (France, Ireland, Italy, The Netherlands, Portugal, Spain and the United Kingdom), Canada (British Columbia), the United States of America (California, Maine and Washington states), and New Zealand (Lane *et al.*, 2016). The parasite was also reported for the first time in flat oysters in Denmark in 2014 (ICES, 2015).

7.2.3 Short Description of Clinical Signs

B. ostreae is an intrahaemocytic protozoan but it can be observed extracellularly between epithelial or interstitial cells in the gills and stomach or in necrotic connective tissue areas. Intraepithelial localisation has also been reported in gills (Montes *et al.*, 1994) and the parasite has been reported in ovarian tissue (Van Banning, 1990). Advanced infections can become systemic. *O. edulis* of a year or less in age can develop a high prevalence and intensity of infection with associated mortality within six months of exposure to *B. ostreae* (Lynch *et al.*, 2005). However, individuals older than two years appear to be more susceptible to the disease (Culloty and Mulcahy, 1996; Grizel, 1985; Engelsma *et al.*, 2010). Seed from natural settlements appear to be significantly more parasitised than oyster seed from hatcheries (Conchas *et al.*, 2003). Infection of wild and cultured flat oysters is often lethal, and death usually occurs concurrently with the highest intensity infection level.

7.2.4 Control/Preventative Measures

Infection caused by *B. ostreae* in molluscs is listed as a notifiable disease under EU legislation (Directive 2006/88/EC) and also by the OIE (www.oie.int). Mortalities caused by bonamiosis can be reduced using suspension culture, lower stocking densities or by culturing *O. edulis* with *C. gigas*, which is far less susceptible to infection. Oyster seed from hatcheries are preferred for aquaculture use over seed from natural settlements as the latter appear to be significantly more parasitised (Conchas *et al.*, 2003). Resistant strains of *O. edulis* developed through selective breeding may offer an alternative in

infected areas. Selective breeding has been shown to be effective in reducing susceptibility and mortality caused by *B. ostreae* (Naciri-Graven *et al.*, 1998).

7.2.5 Host species

O. edulis is the only known species that is significantly naturally susceptible to *B. ostreae*, *C. gigas* being only scarcely susceptible at most (Lynch *et al.*, 2010). Infection intensity increases concurrently to mortality with age and/or size of the oysters (Culloty and Mulcahy, 1996; Grizel, 1985).

7.3 Marteiliosis

7.3.1 Description of Agent

Marteiliosis is a disease of marine bivalve molluscs caused by protozoan parasites in the genus *Marteilia* (order Paramyxida, phylum Cercozoa; Cavalier-Smith and Chao, 2003), including *M. refringens*, *M. sydneyi*, *M. chungmuensis*, *M. cochillia* among other described species. *Marteilia* species display a cell-within-a-cell structure in which triceellular spores are produced; the pattern of secondary cells produced within primary cells and triceellular spores within secondary cells distinguishes individual *Marteilia* species (Feist *et al.*, 2009). Sporulation occurs extracellularly within digestive tubule epithelia in *M. refringens* (Herrbach, 1971), *M. sydneyi* (Wolf, 1972), and *M. cochillia* (Carrasco *et al.*, 2011), but within oocytes in Asian *M. chungmuensis* (Comps *et al.*, 1986).

7.3.2 Geographical Distribution and Temporal Trends

M. refringens occurs from France southward along the Atlantic coast of Europe as well as in the Mediterranean Sea at least as far east as Greece, and in Tunisia (Virvilis and Angelidis, 2003; Carrasco *et al.*, 2007; Elgharsalli *et al.*, 2013; Arzul *et al.*, 2014). It contributed to the decline of *O. edulis* oyster populations and remains an economically significant pathogen of flat oysters and mussels. Reports over the last decade have noted the occurrence of *M. refringens* in Sweden (ICES 2010, 2011) and southern England (ICES 2012). *M. cochillia* was discovered in the context of significant cockle (*C. edule*) mortality in the Ebro Delta of Mediterranean Spain in 2008 (Carrasco *et al.*, 2011), and was associated with cockle mortality that approached 100% in the Ría de Arousa of the Atlantic coast of Spain in 2012 (Villalba *et al.*, 2014). It represents a notable new threat to an important European fishery species. *M. sydneyi* and *M. chungmuensis* are economically significant pathogens but remain absent from the ICES area, *M. sydneyi* occurring in eastern Australia (Wolf, 1979) and *M. chungmuensis* in Korea and Japan (Comps *et al.*, 1986; Itoh *et al.*, 2002).

7.3.3 Short Description of Clinical Signs

Marteilia parasites sporulating in host digestive tubule epithelia migrate to that tissue from portals of entry elsewhere, primarily the epithelia of palps or stomach in the case of *M. refringens* (Grizel, 1974; Berthe *et al.*, 2004). Infections can be difficult to detect before colonization of the digestive tubules, but parasite proliferation afterward produces sharp increases in infection intensity and marked disruption to digestive tubule structure and function, with emaciation of the host and significant mortality a common result (Figueras and Montes, 1988; Villalba *et al.*, 1993). Gonadal infection by *M. chungmuensis* produces gross nodular lesions in affected Pacific oysters (Bower *et al.*, 2011).

7.3.4 Control/Preventative Measures

Infection with *M. refringens* is a disease listed by both the World Organisation for Animal Health (OIE) and the EU (Directive 2006/88/EC). It is on the US National List of Reportable Animal Diseases and on Canada's list of Federally Reportable Aquatic Animal Diseases - Molluscs. Infection with *M. chungmuensis* is also on the Canadian list, and infection with *M. chungmuensis* and infection with *M. sydneyi* are both under review for US listing. Infection with *M. cochillia* is not listed. As *M. refringens* primarily is a pathogen of warmer estuarine systems, parasite impacts may be minimized by culturing susceptible hosts in higher salinity outer coastal waters and in cooler areas, the parasite requiring temperatures of 17°C or greater to infect oyster hosts (Berthe *et al.*, 2004). *M. refringens* is known to infect copepods *Paracartia grani* (Audemard *et al.*, 2001, 2002) and *P. latisetosa* (Arzul *et al.*, 2014) in addition to its molluscan hosts, but limited understanding of its presumably indirect life cycle otherwise limits options for its control.

7.3.5 Host Species

M. refringens primarily infects oyster *O. edulis* and mussels *M. edulis* and *M. galloprovincialis*, though numerous other molluscs have been proven to be hosts, and there is molecular evidence suggestive of infection in others (www.oie.int). Vigorous debate continues about whether *M. refringens* infecting mussels represents a distinct species, *M. maurini* (Comps *et al.*, 1982). Two distinct genetic lineages of *M. refringens* exist, with parasites on one more likely to infect oysters and parasites on the other to infect mussels (LeRoux *et al.*, 2001; López-Flores *et al.*, 2004, Novoa *et al.*, 2005). While the divergence of these parasite lineages may reasonably be interpreted as a speciation event, imperfect fidelity to host type among the parasites on the two lineages would argue that they should conservatively be regarded as a single species, *M. refringens*, for most effective health management.

M. cochillia infects cockle *C. edule* (Carrasco *et al.*, 2013), *M. sydneyi* infects the Sydney rock oyster *S. glomerata* (Perkins and Wolf, 1976), and *M. chungmuensis* infects the Pacific oyster *C. gigas* (Comps *et al.*, 1986), and possibly suminoe oyster *C. ariakensis* and Manila clam *R. philippinarum* (Yanin *et al.*, 2013).

7.4 Haplosporidiosis

7.4.1 Description of Agent

Haplosporidiosis is caused by protozoan parasites in the Haplosporida, phylum Cercozoa (Cavalier-Smith and Chao, 2003). Members of two genera, *Haplosporidium* and *Minchinia*, are responsible for haplosporidiosis in various molluscs. The parasites present uni- and binucleate cells and plasmodia of varying nuclear counts, in addition to sporogonic forms, with spore structure being used to assign parasites to genus (Burrenson and Ford, 2004; Burrenson and Reece, 2006). All are believed to be indirectly transmissible through intermediate hosts, none of which have ever been identified. Mollusc pathogens causing haplosporidiosis in the region include *H. nelsoni*, *H. costale*, *H. armoricanum*, *H. edule*, and *M. tapetis*.

7.4.2 Geographical Distribution and Temporal Trends

H. nelsoni is the most significant agent of haplosporidiosis. It is native to Pacific oyster *C. gigas* populations in Asia (Burrenson *et al.*, 2000), and is associated with established *C. gigas* populations in other parts of the world, including Europe (Renault *et al.* 2000, Lynch *et al.*, 2013) and western Canada (ICES, 2008), though without causing much

disease. It has been far more pathogenic in eastern oyster *C. virginica* along the Atlantic coast of North America, however, where it ranges from Florida to Maine, USA (Ford, 1996) and occurs also in Nova Scotia, Canada. The most significant *H. nelsoni* impacts in recent years have been in the northern part of its distribution, with major epizootics in Nova Scotia beginning in 2002 (ICES, 2003) and in Maine in 2010. In the Chesapeake and Delaware Bays of the Mid-Atlantic coast of the USA, however, *H. nelsoni* impacts have been declining with increasing disease resistance in the oyster host (Carnegie and Bureson, 2011; Ford and Bushek, 2012). The increased activity of *H. nelsoni* in the north probably represents the influence of climate change, with the parasite (and possibly its unidentified intermediate host) expanding its distribution northward and encountering relatively naive host populations. In the Mid-Atlantic, the parasite has been long established, providing the host more time to adapt to its presence (Ford and Bushek, 2012).

H. costale occurs from the Mid-Atlantic coast of the USA to the Atlantic Provinces of Canada (ICES, 2003) and reportedly has also been observed more frequently in northern waters, but this could reflect increased surveillance more than differences in pathogen dynamics in those areas. There is no evidence that *H. costale* contributes more than occasionally to substantial oyster mortality in eastern North America. *H. costale* was observed in 2008 in *C. gigas* in British Columbia at 4-10% prevalence (ICES, 2009).

The distribution of *H. armoricanum* includes the Netherlands and both Atlantic and Mediterranean coasts of France (Hine *et al.*, 2007) as well as Spain (Azevedo *et al.*, 1999), with a likely recent observation in Ireland (Lynch *et al.*, 2013). *H. edule* occurs in northwest Spain (Azevedo *et al.*, 2003) and *M. tapetis* occurs in northwest Spain and Portugal (Vilela, 1951; Azevedo, 2001), but both of these pathogens were detected in 2009 in Wales (ICES, 2010). An undescribed haplosporidian basal to the described genera was observed in 2002 in *R. decussatus* from Spain at high prevalence (to 71%) but without associated mortality (Novoa *et al.*, 2004); this pathogen has not been reported since.

7.4.3 Short Description of Clinical Signs

Haplosporidiosis generally involves invasion of hemal spaces in host connective tissues by uninucleate or plasmodial forms, followed by the production of masses of operculate spores in those tissues. One species of the derived haplosporidian genus *Bonamia*, *B. perspora*, presents a typical haplosporidiosis in its oyster host (Carnegie *et al.*, 2006), but none of the other *Bonamia* species do. *H. nelsoni*, sporulating in host digestive tubule epithelia, is a notable exception with regard to tissue tropism (Couch *et al.*, 1966).

7.4.4 Control/Preventative Measures

No haplosporidiosis is listed as notifiable by either the World Organisation for Animal Health (OIE) or the EU (Directive 2006/88/EC). Infection with *H. nelsoni* is on the list of Federally Reportable Aquatic Animal Diseases – Molluscs for Canada. Limited understanding of the presumably indirect life cycles of these pathogens limits options for their control. Breeding for resistance to *H. nelsoni* in the eastern USA has been highly successful (Haskin and Ford, 1979; Ragone Calvo *et al.*, 2003).

7.4.5 Host Species

H. nelsoni infects oysters *C. virginica* and *C. gigas* (Haskin *et al.* 1966; Bureson *et al.* 2000). *H. costale* infects *C. virginica* (Wood and Andrews, 1962) and has been reported recently from *C. gigas* in British Columbia, Canada. *H. armoricanum* infects oyster *O.*

edulis (Van Banning, 1977). *H. edule* infects cockle *C. edule* (Azevedo *et al.*, 2003). *M. tapetis* infects clam *R. decussatus* (Vilela, 1951).

7.5 Perkinsosis

7.5.1 Description of Agent

Perkinsosis is caused by protozoan parasites in the genus *Perkinsus*, order Perkinsida (Levine, 1978). Infections by *Perkinsus* parasites have been noted in a number of marine molluscs, primarily of the classes Bivalvia and Gastropoda. Seven species are presently accepted, *P. marinus*, *P. olseni*, *P. chesapeaki*, *P. mediterraneus*, *P. beihaiensis*, *P. honshuen-sis*, and *P. qugwadi*. Of these, *P. marinus* and *P. olseni* have been the most impactful on host populations and associated industries. *Perkinsus* parasites typically occur extracellularly in host connective tissues, with both uninucleate trophozoites and multinucleate schizonts observed, but *P. marinus* displays a distinct tropism for digestive epithelia (Carnegie and Burrenson, 2012), a habit that it may share with *P. beihaiensis* (Moss *et al.*, 2008). *P. qugwadi* is unique in expressing zoospore forms within host tissues (Blackbourn *et al.*, 1998).

7.5.2 Geographical Distribution and Temporal Trends

Of the two major *Perkinsus* pathogens, *P. marinus* occurs from the northeastern USA to Mexico in the western Atlantic Ocean and Gulf of Mexico, with colonization of the Pacific coast of Mexico presumed to be a recent event (Caceres-Martinez *et al.*, 2008). It was recently detected as well in Brazil (Da Silva *et al.*, 2013). *P. olseni* was described from Australia (Lester and Davis, 1981) but occurs widely in Asia and southern Europe as well as in New Zealand (Dungan *et al.*, 2007), and more recent work has found it in Uruguay and Brazil along the Atlantic coast of South America (Cremonte *et al.*, 2005; Da Silva *et al.*, 2014). Anthropogenic contributions to the distribution of *P. olseni* are not known.

The recent observations of *P. chesapeaki*, described from the eastern USA, in Europe (Arzul *et al.*, 2012; Carrasco *et al.*, 2014) and Australia (Dang *et al.*, 2015), and of *P. beihaiensis*, described from China (Moss *et al.*, 2008), in Brazil (Sabry *et al.*, 2009) and India (Sanil *et al.*, 2012) are noteworthy. As for *P. olseni*, however, it is not clear to what extent these observations represent emergence of established pathogens in new locations; the discoveries may be a product of increased surveillance. *P. qugwadi* reemerged at one location in British Columbia, Canada, in 2011 after not having been observed anywhere since 1997 (Itoh *et al.*, 2013).

7.5.3 Short Description of Clinical Signs

While very small (2–3 µm) *P. marinus* can be phagocytosed by host hemocytes and distributed generally through host hemal spaces, larger *Perkinsus* species are frequently observed as clusters or masses of parasite cells surrounded and sometimes encapsulated by host hemocytes (Blackbourn *et al.*, 1998; McLaughlin and Faisal, 1998; Burrenson *et al.*, 2005; Park *et al.*, 2006). Infections reaching high intensities can be lethal, and infection by *P. marinus* has caused mortality exceeding 70% (Burrenson and Ragone Calvo, 1996). Emaciation has long been considered a hallmark of perkinsosis caused by *P. marinus* and continues to be observed in association with heavy infections, although this clinical presentation can have numerous other causes (Carnegie and Burrenson, 2012).

7.5.4 Control/Preventative Measures

Both *P. marinus* and *P. olsenii* are notifiable to the World Organisation for Animal Health (OIE). Infection with *P. marinus* is listed by the EU (Directive 2006/88/EC) and on the list of Federally Reportable Aquatic Animal Diseases – Molluscs for Canada. Infection with *P. olsenii* is on the Canadian list and the US National List of Reportable Animal Diseases as well. As *Perkinsus* parasites are directly transmissible and easily spread with transfer of infected stocks, avoiding introduction to new areas with movement of infected shellfish is critical. In *P. marinus*-enzootic areas of the USA, control of *P. marinus* in aquaculture populations through selective breeding for disease-resistant oysters has been successful (Ragone Calvo *et al.*, 2003).

7.5.5 Host Species

P. marinus infects the eastern oyster *C. virginica* (Mackin *et al.*, 1950) but also oysters *C. corteziensis* (Caceres-Martinez *et al.*, 2008), *S. palmula* (Caceres-Martinez *et al.*, 2012), and *C. rhizophorae* (Da Silva *et al.*, 2013). *P. olsenii* (= *P. atlanticus*) was described from the abalone *H. ruber* (Lester and Davis, 1981) but infects a wide range of bivalve and gastropod species, notably including Manila clam *R. philippinarum* in Asia (Choi and Park, 1997) and Europe (Navas *et al.* 1992). *P. chesapeaki* (= *P. andrewsi*) infects clams *Macoma balthica* (Coss *et al.*, 2001), *Mya arenaria* and *Tagelus plebeius* (Dungan *et al.*, 2002), *Cyrtopleura costata* (Reece *et al.* 2008), and *R. decussatus* and *R. philippinarum* (Arzul *et al.*, 2012), cockle *C. edule* (Carrasco *et al.*, 2014), and the ark *Anadara trapezia* (Dang *et al.*, 2015). *P. beihaiensis* infects oysters *C. hongkongensis* and *C. ariakensis* (Moss *et al.* 2008), *C. rhizophorae* (Sabry *et al.*, 2009), and *C. madrasensis* (Sanil *et al.*, 2012), and clam *Anomalocardia brasiliensis* (Ferreira *et al.*, 2015). *P. mediterraneus* infects oyster *O. edulis* (Casas *et al.*, 2004) and ark *Arca noae* and scallop *Chlamys varia* (Ramilo *et al.*, 2015). *P. honsuensis* infects the clam *R. philippinarum* (Dungan and Reece, 2006). *P. qugwadi* infects the scallop *Patinopecten yessoensis* (Blackbourn *et al.*, 1998).

Molluscan References

- Abollo, E., Ramilo, A., Casas, S.M., Comesana, P., Cao, A., Carballal, M.J. and Villalba A. 2008. First detection of the protozoan parasite *Bonamia exitiosa* (Haplosporidia) infecting flat oyster *Ostrea edulis* grown in European waters. *Aquaculture* 274: 201–207.
- Arzul, I., Nicolas, J. L., Davison, A. J., Renault, T. 2001. French scallops: a new host for ostreid herpesvirus 1. *Virology* 290: 342–349.
- Arzul, I., Chollet, B., Michel, J., Robert, M., Garcia, C., Joly, J.P., Francois, C. and Miossec, L. 2012. One *Perkinsus* species may hide another: characterization of *Perkinsus* species present in clam production areas of France. *Parasitology* 139: 1757–1771.
- Arzul, A., Chollet, B., Boyer, S., Bonnet, D., Gaillard, J., Baldi, Y., Robert, M., Joly, J.P., Garcia, C. and Bouchoucha, M. 2014. Contribution to the understanding of the cycle of the protozoan parasite *Marteilia refringens*. *Parasitology* 141: 227–240.
- Audemard, C., Barnaud, A., Collins, C.M., Le Roux, F., Sauriau, P.G., Coustau, C., Blachier, P. and Berthe, F.C.J., 2001. Claire ponds as an experimental model for *Marteilia refringens* life-cycle studies: new perspectives. *Journal of Experimental Marine Biology and Ecology* 257: 87–108.
- Audemard, C., Le Roux, F., Barnaud, A., Collins, C., Sautour, B., Sauriau, P.G., De Montaudouin, X., Coustau, C., Combes, C. and Berthe, F. 2002. Needle in a haystack: involvement of the copepod *Paracartia grani* in the life-cycle of the oyster pathogen *Marteilia refringens*. *Parasitology* 124: 315–323.
- Azevedo, C. 2001. Ultrastructural description of the spore maturation stages of the clam parasite *Minchinia tapetis* (Vilela, 1951) (Haplosporida: Haplosporidiidae). *Systematic Parasitology* 49: 189–194.
- Azevedo, C., Conchas, R.F. and Montes, C. 2003. Description of *Haplosporidium edule* n. sp. (Phylum Haplosporidia), a parasite of *Cerastoderma edule* (Mollusca, Bivalvia) with complex spore ornamentation. *European Journal of Protistology* 39: 161–167.
- Azevedo, C., Montes, J. and Corral, L. 1999. A revised description of *Haplosporidium armoricanum*, parasite of *Ostrea edulis* L. from Galicia, northwestern Spain, with special reference to the spore-wall filaments. *Parasitology Research* 85: 977–983.
- Azandégbé, A., Garnier, M., Andrieux-Loyer, F., Kérouel, R., Philippon, X., Nicolas, J. L. 2010. Occurrence and seasonality of *Vibrio aestuarianus* in sediment and *Crassostrea gigas* haemolymph at two oyster farms in France. *Diseases of Aquatic Organisms* 91: 213–221.
- Balseiro, P., Aranguren, R., Gestal, C., Novoa, B. and Figueras, A. 2006. *Candidatus Xenohalictis californiensis* and *Haplosporidium montforti* associated with mortalities of abalone *Haliotis tuberculata* cultured in Europe. *Aquaculture* 258: 63–72.
- Berthe, F.C.J., Le Roux, F., Adlard, R.D. and Figueras, A. 2004. Martelliosis in molluscs: A review. *Aquatic Living Resources* 17: 433–448.
- Blackbourn, J., Bower, S.M. and Meyer, G.R. 1998. *Perkinsus qugwadi* sp. nov. (incertae sedis), a pathogenic protozoan parasite of Japanese scallops, *Patinopecten yessoensis*, culture in British Columbia, Canada. *Canadian Journal of Zoology* 76: 942–953.
- Bower, S. M., Goh, B. and Meyer, G. R. 2005. Epizootiology and detection of nocardiosis in oysters. In P. Walker, R. Lester and M.G. Bondad-Reantaso (eds). *Diseases in Asian Aquaculture V*, pp 249–262. Fish Health Section, Asian Fisheries Society, Manila.
- Bower, S. M., Itoh, N., Choi, D.-L., Park, M.S. (2011): Synopsis of Infectious Diseases and Parasites of Commercially Exploited Shellfish: *Marteilioides chungmuensis* of Oysters. <http://www.dfo-mpo.gc.ca/science/aah-saa/diseases-maladies/mcoy-eng.html>.
- Burreson, E.M. and Ford, S.E. 2004. A review of recent information on the Haplosporidia, with special reference to *Haplosporidium nelsoni* (MSX disease). *Aquatic Living Resources* 17: 499–517.

- Burreson, E.M. and Ragone Calvo, L.M. 1996. Epizootiology of *Perkinsus marinus* disease of oysters in Chesapeake Bay, with emphasis on data since 1985. *Journal of Shellfish Research* 15: 17–34.
- Burreson, E.M. and Reece, K.S. 2006. Spore ornamentation of *Haplosporidium nelsoni* and *Haplosporidium costale* (Haplosporidia), and incongruence of molecular phylogeny and spore ornamentation in the Haplosporidia. *Journal of Parasitology* 92: 1295–1301.
- Burreson, E.M., Dungan, C.F. and Reece, K.S. 2005. Molecular, morphological, and experimental evidence support the synonymy of *Perkinsus chesapeaki* and *Perkinsus andrewsi*. *Journal of Eukaryotic Microbiology* 52: 258–270.
- Burreson, E.M., Stokes, N.A. and Friedman, C.S. 2000. Increased virulence in an introduced pathogen: *Haplosporidium nelsoni* (MSX) in the eastern oyster *Crassostrea virginica*. *Journal of Aquatic Animal Health* 12: 1–8.
- Cáceres-Martínez, J., Vásquez-Yeomans, R., Padilla-Lardizábal, G. and del Río Portilla, M.A. 2008. *Perkinsus marinus* in pleasure oyster *Crassostrea corteziensis* from Nayarit, Pacific coast of México. *Journal of Invertebrate Pathology* 99: 66–73.
- Cáceres-Martínez, J., García Ortega, M., Vásquez-Yeomans, R., Pineda García, T.J., Stokes, N.A. and Carnegie, R.B. 2012. Natural and cultured populations of the mangrove oyster *Saccostrea palmula* from Sinaloa, Mexico, infected by *Perkinsus marinus*. *Journal of Invertebrate Pathology* 110: 321–325.
- Carella, F., Carrasco, N., Andree, K. B., Lacuesta, B., Furones, D., and De Vico, G. 2013. Nocardiosis in Mediterranean bivalves: first detection of *Nocardia crassostreae* in a new host *Mytilus galloprovincialis* and in *Ostrea edulis* from the Gulf of Naples (Italy). *Journal of Invertebrate Pathology* 114: 324–328.
- Carnegie, R.B. and Burreson, E.M. 2011. Declining impact of an introduced pathogen: *Haplosporidium nelsoni* in the oyster *Crassostrea virginica* in Chesapeake Bay. *Marine Ecology Progress Series* 432: 1–15.
- Carnegie, R.B. and Burreson, E.M. 2012. *Perkinsus marinus* and *Haplosporidium nelsoni*. Pages 92–108 in: *Fish Parasites: Pathobiology and Protection* (P.T.K. Woo and K. Buchmann, eds.). CABI Publishing, Oxfordshire, UK.
- Carnegie, R.B., Hill, K.M., Stokes, N.A. and Burreson, E.M. 2014. The haplosporidian *Bonamia exitiosa* is present in Australia, but the identity of the parasite described as *Bonamia roughleyi* is uncertain. *Journal of Invertebrate Pathology* 115: 33–40.
- Carnegie, R.B., B.J. Barber, S.C. Culloty, A.J. Figueras, and D.L. Distel. 2000. Development of a PCR assay for detection of the oyster pathogen *Bonamia ostreae* (Pichot *et al.* 1980), and support for its inclusion in the Haplosporidia. *Diseases of Aquatic Organisms* 42: 199–206.
- Carnegie, R.B., Burreson, E.M., Hine, P.M., Stokes, N.A., Audemard, C., Bishop, M.J. and Peterson, C.H., 2006. *Bonamia perspora* n. sp. (Haplosporidia), a parasite of the oyster *Ostreola equestris*, is the first *Bonamia* species known to produce spores. *Journal of Eukaryotic Microbiology* 53: 232–245.
- Carrasco, N., López-Flores, I., Alcaraz, M., Furones, M.D., Berthe, F. and Arzul, I., 2007. Dynamics of the parasite *Marteilia refringens* (Paramyxia) in *Mytilus galloprovincialis* and zooplankton populations in Alfacs Bay (Catalonia, Spain). *Parasitology* 134: 1541–1550.
- Carrasco, N., Rojas, M., Aceituno, P., Andree, K.B., Lacuesta, B. and Furones, M.D. 2014. *Perkinsus chesapeaki* observed in a new host, the European common edible cockle *Cerastoderma edule*, in the Spanish Mediterranean coast. *Journal of Invertebrate Pathology* 117: 56–60.
- Carrasco, N., Roque, A., Andree, K.B., Rodgers, C., Lacuesta, B. and Furones, M.D. 2011. A *Marteilia* parasite and digestive epithelial virosis observed during a common edible cockle *Cerastoderma edule* mortality event in the Spanish Mediterranean coast. *Aquaculture* 321: 197–202.

- Carrasco, N., Hine, P.M., Durfort, M., Andree, K.B., Malchus, N., Lacuesta, B., González, M., Roque, A., Rodgers, C. and Furones, M.D. 2013. *Marteilia cochillia* sp. nov., a new *Marteilia* species affecting the edible cockle *Cerastoderma edule* in European Waters. *Aquaculture* 412-413: 223–230.
- Casas, S.M., Grau, A., Reece, K.S., Apakupakul, K., Azevedo, C. and Villalba, A. 2004. *Perkinsus mediterraneus* n. sp., a protistan parasite of the European flat oyster *Ostrea edulis* from the Balearic Islands, Mediterranean Sea. *Diseases of Aquatic Organisms* 58: 231–244.
- Cavalier-Smith, T. and Chao E.E.-Y. 2003. Phylogeny of Choanozoa, Apusozoa, and other protozoa and early eukaryote megaevolution. *Journal of Molecular Evolution* 56: 540–563.
- Choi, K.S. and Park, K.I. 1997. Report on the occurrence of *Perkinsus* sp. in the Manila clams, *Ruditapes philippinarum* in Korea. *Journal of the Korean Aquaculture Society* 10: 227–237.
- Clegg, T.A., Morrissey, T., Geoghegan, F., Martin, S.W., Lyons, K., Ashe, S. and More, S.J. 2014. Risk factors associated with increased mortality of farmed Pacific oysters in Ireland during 2011. *Preventive Veterinary Medicine* 113: 257–267.
- Comps, M., Park, M.S. and Desportes, I. 1986. Ultrastructural study of *Marteilioides chungmuensis* gen. nov., sp. nov., a parasite of the ovocytes of the oyster *Crassostrea gigas* Th. *Protistologica* 22: 279–285.
- Comps M., Pichot, Y. and Papagianni, P. 1982. Recherche sur *Marteilia maurini* n. sp. parasite de la moule *Mytilus galloprovincialis* Lmk. *Rev. Trav. Inst. Pêches Marit.* 45: 211–214.
- Comps, M., Tigé, G., and Grizel, H. 1980. Etude ultrastructurale d'un protiste parasite de l'huître *Ostrea edulis* L. *LCR Academy of Science Paris, Série D* 290, 383–385.
- Conchas, R.F., Santamarina, J., Lama, A., Longa, M.A., and Montes, J. 2003. Evolution of bonamiosis in Galicia (NW Spain). *Bulletin of the European Association of Fish Pathologists* 23, 265–272.
- Corbeil, S., Arzul, I., Robert, M., Berthe, F.C.J., Besnard-Cochennec, N. and Crane, M.S.J. 2006. Molecular characterization of an Australian isolate of *Bonamia exitiosa*. *Diseases of Aquatic Organisms* 71 : 81–85.
- Coss, C.A., Robledo, J.A.F., Ruiz, G.M. and Vasta, G.R. 2001. Description of *Perkinsus andrewsi* n. sp. isolated from the Baltic clam (*Macoma balthica*) by characterization of the ribosomal RNA locus, and development of a species-specific PCR-based diagnostic assay. *Journal of Eukaryotic Microbiology* 48: 52–61.
- Couch, J.A., Farley, C.A and Rosenfield, A. 1966. Sporulation of *Minchinia nelsoni* (Haplosporida, Haplosporidiidae) in *Crassostrea virginica* (Gmelin). *Science* 153: 1529–1531.
- Cremonte, F., Balseiro, P. and Figueras, A. 2005 Occurrence of *Perkinsus olseni* (Protozoa: Apicomplexa) and other parasites in the venerid commercial clam *Pitar rostrata* from Uruguay, southwestern Atlantic coast. *Diseases of Aquatic Organisms* 64: 85–90.
- Cranfield, H.J., Dunn, A., Doonan, I.J. and Michael, K.P. 2005. *Bonamia exitiosa* epizootic in *Ostrea chilensis* from Foveaux Strait, southern New Zealand between 1986 and 1992. *ICES Journal of Marine Science* 62 : 3–13.
- Culloty, S.C. and Mulcahy, M.F. 1996. Season-, age-, and sex-related variation in the prevalence of bonamiosis in flat oysters (*Ostrea edulis* L.) on the south coast of Ireland. *Aquaculture* 144, 53–63.
- Dang, C., Dungan, C.F., Scott, G.P. and Reece, K.S. 2015. *Perkinsus* sp. infections and *in vitro* isolates from *Anadara trapezia* (mud arks) of Queensland, Australia. *Diseases of Aquatic Organisms* 113: 51–58.
- da Silva, P.M., Scardua, M.P., Vianna, R.T., Mendonça, R.C., Vieira, C.B., Dungan, C.F., Scott, G.P., and Reece, K.S. 2014. Two *Perkinsus* spp. infect *Crassostrea gasar* oysters from cultured and wild populations of the Rio São Francisco estuary, Sergipe, northeastern Brazil. *Journal of Invertebrate Pathology* 119: 62–71.

- da Silva, P.M., Vianna, R.T., Guertler, C., Ferreira, L.P., Santana, L.N., Fernandez-Boo, S., Ramilo, A., Cao, A. and Villalba, A. 2013. First report of the protozoan parasite *Perkinsus marinus* in South America, infecting mangrove oysters *Crassostrea rhizophorae* from the Paraíba River (NE, Brazil). *Journal of Invertebrate Pathology* 113: 96-103.
- Davison, A.J., Trus, B.L., Cheng, N., Steven, A.C., Watson, M.S., Cunningham, C., Le Deuff, R.M. and Renault, T. 2005. A novel class of herpesvirus with bivalve hosts. *Journal of General Virology* 86: 41-53.
- Dinamani, P., Hine, P.M. and Jones, J.B. 1987. Occurrence and characteristics of the haemocyte parasite *Bonamia* sp. in the New Zealand dredge oyster *Tiostrea lutaria*. *Diseases of Aquatic Organisms* 3: 37-44.
- Dundon, W. G., Arzul, I., Omnes, E., Robert, M., Magnabosco, C., Zambon, M., Gennari, L., Toffan, A., Terregino, C., Capua, I., and Arcangeli, G. 2011. Detection of Type 1 Ostreid Herpes variant (OsHV-1 μ var) with no associated mortality in French-origin Pacific cupped oyster *Crassostrea gigas* farmed in Italy. *Aquaculture* 314: 49-52.
- Dungan, C.F. and K.S. Reece. 2006. In vitro propagation of two *Perkinsus* spp. parasites from Japanese Manila clams *Venerupis philippinarum*, and description of *Perkinsus honshuensis* n. sp. *Journal of Eukaryotic Microbiology* 53: 316-326.
- Dungan, C.F., Hamilton, R.M., Hudson, K.L., McCollough, C.B. and Reece, K.S. 2002 Two epizootic diseases in Chesapeake Bay commercial clams, *Mya arenaria* and *Tagelus plebeius*. *Diseases of Aquatic Organisms* 50: 67-78.
- Dungan, C.F., Reece, K.S., Moss, J.A., Hamilton, R.M. and Diggles, B.K. 2007. *Perkinsus olseni* in vitro isolates from the New Zealand clam *Austrovenus stutchburyi*. *Journal of Eukaryotic Microbiology* 54: 263-270.
- EFSA. 2010. Scientific Opinion of the Panel on Animal Health and Welfare on a request from the European Commission on the increased mortality events in Pacific oysters *Crassostrea gigas*. *EFSA Journal* 8:1894-1853.
- Elgharsalli, R., Aloui-Bejaoui, N., Salah, H., Chollet, B., Joly, J.-P., Robert, M., Couraleau, Y. and Arzul, I. 2013. Characterization of the protozoan parasite *Marteilia refringens* infecting the dwarf oyster *Ostrea stentina* in Tunisia. *Journal of Invertebrate Pathology* 112: 175-183.
- Engelsma, MY, Roozenburg, I. and Joly, J.P. 2008. First isolation of *Nocardia crassostreae* from Pacific oyster *Crassostrea gigas* in Europe. *Diseases of Aquatic Organisms* 80:229-234.
- Engelsma, M.Y., Kerkhoff, S., Roozenburg, I., Haenen, O.L.M., Van Gool, A., Istermans, W., Wijnhoven, S., and Hummel, H. 2010. Epidemiology of *Bonamia ostreae* infecting European flat oyster *Ostrea edulis* from Lake Grevelingen, The Netherlands. *Marine Ecology Progress Series* 409, 131-142.
- Feist, S.W., Hine, M.P., Bateman, K.S., Steinford, G.D. and Longshaw, M., 2009. *Paramarteilia canceri* sp. n. (Cercozoa) in the European edible crab (*Cancer pagurus*) with a proposal for the revision of the order paramyxida Chatton, 1911. *Folia Parasitologica* 56: 73-85.
- Ferreira, L.P., Sabry, R.C., da Silva, P.M., Gesteira, T.C.V., Romao, L.D., Paz, M.P., Feijó, R.G., Neto, M.P.D. and Maggioni, R. 2015. First report of *Perkinsus beihaiensis* in wild clams *Anomalocardia brasiliensis* (Bivalvia: Veneridae) in Brazil. *Experimental Parasitology* 150: 67-70.
- Figueras, A.J. and Montes, J. 1988. Aber disease of edible oysters caused by *Marteilia refringens*. *American Fisheries Society Special Publication* 18: 38-46.
- Ford, S.E. 1996. Range extension by the oyster parasite *Perkinsus marinus* into the northeastern United States: response to climate change? *Journal of Shellfish Research* 15: 45-56.
- Ford, S.E. and Bushek, D. 2012. Development of resistance to an introduced marine pathogen by a native host. *Journal of Marine Research* 70: 205-223.

- Friedman, C.S. and Finley C.A. 2003. Evidence for an anthropogenic introduction of “*Candidatus Xenohaliotis californiensis*”, the etiological agent of withering syndrome, into northern California abalone population via conservation efforts. *Canadian Journal of Fisheries and Aquatic Sciences* 60: 1424-1431.
- Friedman, C.S., Beattie, J. H., Elston, R.A. and Hedrick, R.P. 1991. Investigation of the relationship between the presence of a Gram-positive bacterial infection and summer mortality of the Pacific oyster, *Crassostrea gigas* Thunberg. *Aquaculture* 94: 1.
- Friedman, C.S., Beaman, B. L., Chun, J., Goodfellow, M., Gee, A. and Hedrick, R.P. 1998. *Nocardia crassostreae* sp. nov., the causal agent of nocardiosis in Pacific oysters. *International Journal of Systematic Bacteriology* 48 Pt 1: 237-46.
- Friedman, C.S., Andree, K.B., Beauchamps, K.A., Moore, J.D., Robbins, T.T., Shields, J.D., and Hedrick, R.P. 2000. “*Candidatus Xenohaliotis californiensis*” a newly described pathogen of abalone, *Haliotis* spp., along the west coast of North America. *International Journal of Systematic Evolution and Microbiology* 50: 847-855.
- Garcia, C., Thébault, A., Dégremont, L., Arzul, I., Miossec, L., Robert, M., Chollet, B., François, C., Joly, J.-P., Ferrand, S., Kerdudou, N. and Renault, T. 2011. OsHV-1 detection and relationship with *C. gigas* spat mortality in France between 1998 and 2006. *Veterinary Research* 42 : 73–84.
- Gardner, G.R., Harshbarger, J.C., Lake, J., Sawyer, T.K., Price, K.L., Stephenson, M.D., Haaker, P.L., Togstad, H.A. 1995. Association of prokaryotes with symptomatic appearance of withering syndrome in black abalone *Haliotis cracherodii*. *Journal of Invertebrate Pathology* 66: 111-120.
- Garnier, M., Y. Labreuche, C. Garcia, M. Robert and J.L. Nicolas. 2007. Evidence for the Involvement of Pathogenic Bacteria in Summer Mortalities of the Pacific Oyster *Crassostrea gigas*. *Microbial Ecology* 53: 187-196.
- Garnier, M., Y. Labreuche, and Nicolas, J. L. 2008 Molecular and phenotypic characterization of *Vibrio aestuarianus* subsp. *francensis* subsp. nov., a pathogen of the oyster *Crassostrea gigas*. *Systematic and Applied Microbiology* 31: 358-365.
- Gay, M., Renault, T., Pons, A.M. and Le Roux, F. 2004. Two *Vibrio splendidus* related strains collaborate to kill *Crassostrea gigas*: taxonomy and host alterations. *Diseases of Aquatic Organisms* 62: 65-74.
- Gay, M., Lancelot, G., Chollet, B., Renault, T., Cochenne, C., Berthe, F.J., Lambert, C., Choquet, G., Paillard, C., Gouy, M., Le Roux, F. and Goulletquer, P. 2003. Characterisation of *Vibrio* isolated from Pacific oysters spat suffering from Summer Mortality outbreaks. *Journal of Shellfish Research* 22: 331.
- Gomez-Gil, B., Soto-Rodriguez, S., Garcia-Gasca, A., Roque, A., Vazquez-Juarez, R., Thompson, F.L. and Swings, J., 2004. Molecular identification of *Vibrio harveyi*-related isolates associated with diseased aquatic organisms. *Microbiology* 150: 1769-1777.
- Grizel H. 1985. Etudes des récentes épizooties de l’huître plate *Ostrea edulis* L. et de leur impact sur l’ostréiculture bretonne. Thèse de doctorat, Université des Sciences et Techniques de Languedoc, Montpellier, France.
- Grizel, H., Comps, M., Bonami, J.R., Cousserans, F., Duthoit, J.L. and Pennec, M. 1974. Epizooty of the common oyster *Ostrea edulis*. Part 1. Study of the agent of digestive gland disease in *Ostrea edulis* (Linne). *Sciences Pêche* 1–30.
- Handlinger, J., Carson, J., Donachie, L., Gabor, L. and Taylor, D. 2005. Bacterial Infection in Tasmanian Farmed Abalone: Causes, Pathology, Farm Factors and Control Options. Diseases in Asian Aquaculture. P. Walker, R. Lester and M.G. Bondad-Reantaso (Eds). Fish Health Section.

- Haskin, H.H. and Ford, S.E. 1979. Development of resistance to *Minchinia nelsoni* (MSX) mortality in laboratory-reared and native oyster stocks in Delaware Bay. *Marine Fisheries Review* 41: 54–63.
- Haskin, H.H., Stauber, L.A. and Mackin, J.A. 1966 *Minchinia nelsoni* n. sp. (Haplosporida, Haplosporidiidae): causative agent of the Delaware Bay oyster epizootic. *Science* 153: 1414–1416.
- Herrbach, B. 1971. Sur une affection parasitaire de la glande digestive de l'huitre plate, *Ostrea edulis* Linne. *Revue des Travaux de l'Institut des Peches Maritimes*. 35: 79–87.
- Hill, K.M., Carnegie, R.B., Aloui-Bejaoui, N., El Gharsalli, R., White, D., Stokes, N.A. and Burrenson, E.M. 2010. Observation of a *Bonamia* sp. infecting the oyster *Ostrea stentina* in Tunisia, and a consideration of its phylogenetic affinities. *Journal of Invertebrate Pathology* 103: 179–185.
- Hill, K.M., Stokes, N.A., Webb, S.C., Hine, P.M., Kroeck, M.A., Moore, J.D., Morley, M.S., Reece, K.S., Burrenson, E.M. and Carnegie, R.B. 2014. Phylogenetics of *Bonamia* parasites based on small subunit and internal transcribed spacer region ribosomal DNA sequence data. *Diseases of Aquatic Organisms* 110: 33–54.
- Hine, P.M. 1991. The annual pattern of infection by *Bonamia* sp. in New Zealand flat oysters, *Tiostrea chilensis*. *Aquaculture* 93: 241–251.
- Hine, P. M. 1996. The ecology of *Bonamia* and decline of bivalve molluscs. *New Zealand Journal of Ecology* 20: 109–116.
- Hine, P.M. and Jones, J.B. 1994. *Bonamia* and other aquatic parasites of importance to New Zealand. *New Zealand Journal of Zoology* 21: 49–56.
- Hine, P.M. and Wesney, B. 1994. The functional cytology of *Bonamia* sp. (Haplosporidia) infecting oysters *Tiostrea chilensis*: an ultracytochemical study. *Diseases of Aquatic Organisms* 20: 207–217.
- Hine, P.M., Diggles, B.K., Parsons, M.J.D., Pringle, A. And Bull, B. 2002. The effects of stressors on the dynamics of *Bonamia exitiosus* Hine, Cochenne-Laureau & Berthe, infections in flat oysters *Ostrea chilensis* (Philippi). *Journal of Fish Diseases* 25: 545–554.
- Hine, P.M., Engelsma, M.Y. and Wakefield, S.J. 2007. Ultrastructure of sporulation in *Haplosporidium armoricum*. *Diseases of Aquatic Organisms* 77: 225–233.
- Huchette, S.M.H. and Clavier, J. 2004. Status of the ormer (*Haliotis tuberculata* L.) industry in Europe. *Journal of Shellfish Research* 23: 951–955.
- ICES. 2003. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 11–15 March 2003, Aberdeen, United Kingdom. ICES CM 2003/F:03. 101 pp.
- ICES. 2008. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 4–8 March 2008, Galway, Ireland. ICES CM 2008/MCC:01. 128 pp.
- ICES. 2009. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 24–28 February 2009, Riga, Latvia. ICES CM 2009/MCC:01. 119 pp.
- ICES. 2010. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 23–27 February 2010, Uppsala, Sweden. ICES CM 2010/SSGHIE:02. 66 pp.
- ICES. 2011. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 1–5 March 2015, Aberdeen, United Kingdom. ICES CM 2015/SSGHIE:04. 58 pp.
- ICES. 2012. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 31 January - 4 February 2012, Lisbon, Portugal. ICES CM 2012/SSGHIE:03. 74 pp.
- ICES. 2015. Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 24–28 February 2015, Helsinki, Finland. ICES CM 2015/SSGEPI:01. 124 pp.

- Itoh, N., Oda, T., Ogawa, K. and Wakabayashi, H. 2002. Identification and development of a paramyxean ovarian parasite in the Pacific oyster *Crassostrea gigas* (Thunberg). *Fish Pathology* 37: 23–28.
- Itoh, N., Meyer, G.R., Tabata, A., Lowe, G., Abbott, C.L. and Johnson, S.C. 2013. Rediscovery of the Yesso scallop pathogen *Perkinsus qugwadi* in Canada, and development of PCR tests. *Diseases of Aquatic Organisms* 104: 83–91.
- Jenkins, C., Hick, P., Gabor, M., Spiers, Z., Fell, S. A., Gu, X., Read, A., Go, J., Dove, M., O'Connor, W., Kirkland, P. D. and Francesn J. 2013. Identification and characterisation of an ostreid herpesvirus-1 microvariant (OsHV-1 μ -var) in *Crassostrea gigas* (Pacific oysters) in Australia. *Diseases of Aquatic Organisms* 105: 109–126.
- Labreuche Y, Le Roux F, Henry J, Zatylny C, Huvet A, Lambert C, Soudant P, Mazel D, Nicolas JL. 2010. *Vibrio aestuarianus* zinc metalloprotease causes lethality in the Pacific oyster *Crassostrea gigas* and impairs the host cellular immune defenses. *Fish & Shellfish Immunology* 29: 753–8.
- Lacoste, A., Jalabert, F., Malham, S., Cueff, A., Gélébart, F., Cordevant, C., Lange, C. and Poulet, S. 2001. A *Vibrio splendidus* strain associated with Summer Mortality of juvenile oysters *Crassostrea gigas* in the bay of Morlaix (North Brittany, France). *Diseases of Aquatic Organisms* 46: 139–145.
- Lane, H.S., Webb, S.C. and Duncan, J. 2016. *Bonamia ostreae* in the New Zealand oyster *Ostrea chilensis*: a new host and geographic record for this haplosporidian parasite. *Diseases of Aquatic Organisms* 118: 55–63.
- Le Deuff, R.-M. and Renault, T. 1999. Purification and partial genome characterization of a herpes-like virus infecting the Japanese oyster, *Crassostrea gigas*. *Journal of General Virology* 80: 1317–1322.
- Le Roux, F., Gay, M., Lambert, C., Waechter, M., Poubalanne, S., Chollet, B., Nicolas, J.L. and Berthe, F. 2002. Comparative analysis of *Vibrio splendidus*-related strains isolated during *Crassostrea gigas* mortality events. *Aquatic Living Resources* 15: 251–258.
- Le Roux, F., Lorenzo, G., Peyret, P., Audemard, C., Figueras, A., Vivares, C., Gouy, M. and Berthe, F. 2001. Molecular evidence for the existence of two species of *Marteilia* in Europe. *Journal of Eukaryotic Microbiology* 48: 449–454.
- Lester, R.J.G. and Davis, G.H.G. 1981. A new *Perkinsus* species (Apicomplexa, Perkinsea) from the abalone *Haliotis ruber*. *Journal of Invertebrate Pathology* 37: 181–187.
- Levine, N.D. 1978. *Perkinsus* gen. n. and other new taxa in the protozoan phylum Apicomplexa. *Journal of Parasitology* 64: 549.
- López-Flores, I., de la Herran, R., Garrido-Ramos, M., Navas, J., Ruiz-Rejon, C. and Ruiz-Rejon, M. 2004. The molecular diagnosis of *Marteilia refringens* and differentiation between *Marteilia* strains infecting oysters and mussels based on the rDNA IGS sequence. *Parasitology* 129: 411–419.
- Lynch, S.A., Armitage, D.V., Wylde, S., Mulcahy, M.F., and Culloty, S.C. 2005. The susceptibility of young prespawning oysters, *Ostrea edulis*, to *Bonamia ostreae*. *Journal of Shellfish Research* 24, 1019–1025.
- Lynch, S.A., Abollo, E., Ramilo, A., Culloty, S.C. and Villalba, A. 2010. Observations raise the question if the Pacific oyster, *Crassostrea gigas*, can act as either a carrier or a reservoir for *Bonamia ostreae* or *Bonamia exitiosa*. *Parasitology* 137, 1515–1526.
- Lynch, S.A., Villalba, A., Abollo, E., Engelsma, M., Stokes, N.A. and Culloty, S.C. 2013. The occurrence of haplosporidian parasites, *Haplosporidium nelsoni* and *Haplosporidium* sp., in oysters in Ireland. *Journal of Invertebrate Pathology* 112: 208–212.
- Macián, M.C., Garay, E., Gonzalez-Candelas, F., Pujalte, M.J. and Aznar, R. 2000. Ribotyping of vibrio populations associated with cultured oysters (*Ostrea edulis*). *Systematic Applied Microbiology* 23: 409–417.

- Martenot, C., Oden, E., Travaillé, E., Malas, J.P. and Houssin, M. 2011. Detection of different variants of Ostreid Herpesvirus 1 in the Pacific oyster *Crassostrea gigas*. *Virus Research* 160: 25-31.
- McLaughlin, S.M. and M. Faisal. 1998. Histopathological alterations associated with *Perkinsus* spp. infection in the softshell clam *Mya arenaria*. *Parasite – Journal de la Societe Francaise de Parasitologie* 5: 263-271.
- Montes, J., Anadron, R., and Azevedo, C. 1994. A possible life cycle for *Bonamia ostreae* on the basis of electron microscopy studies. *Journal of Invertebrate Pathology* 63, 1–6.
- Moss, J.A., Xiao, J., Dungan, C.F. and Reece, K.S. 2008. Description of *Perkinsus beihaiensis* n. sp., a new *Perkinsus* sp. parasite in oysters of southern China. *Journal of Eukaryotic Microbiology* 55: 117–130.
- Moss, J.A., Bureson, E.M., Cordes, J.F., Dungan, C.F., Brown, G.D., Wang, A., Wu, X. and Reece, K.S. 2007. Pathogens in *Crassostrea ariakensis* and other Asian oyster species: implications for non-native oyster introduction to Chesapeake Bay. *Diseases of Aquatic Organisms* 77: 207–223.
- Naciri-Graven, Y., Martin, A.G., Baud, J.P., Renault, T., and Gérard, A. 1998. Selecting the flat oyster *Ostrea edulis* (L.) for survival when infected with the parasite *Bonamia ostreae*. *Journal of Experimental Marine Biology and Ecology* 224, 91–107.
- Narcisi, V., Arzul, I., Cargini, D., Mosca, F., Calzetta, A., Traversa, D., Robert, M., Joly, J.P., Chollet, B., Renault, T. and Tiscar, P.G. 2010. Detection of *Bonamia ostreae* and *Bonamia exitiosa* (Haplosporidia) in *Ostrea edulis* from the Adriatic Sea (Italy). *Diseases of Aquatic Organisms* 89: 79–85.
- Navas, J.I., Castillo, M.C., Vera, P. and Ruizrico, M. 1992. Principal parasites observed in clams, *Ruditapes decussatus* (L), *Ruditapes philippinarum* (Adams et Reeve), *Venerupis pullastra* (Montagu) and *Venerupis aureus* (Gmelin), from the Huelva Coast (SW Spain). *Aquaculture* 107: 193-199.
- Nicolas, J.L., Basuyaux, O., Mazurie, J. and Thebault, A. 2002. *Vibrio carchariae*, a pathogen of the abalone *Haliotis tuberculata*. *Diseases of Aquatic Organisms* 50: 35-43.
- Nishimori, E., Hasegawa, O., Numata, T. and bayashi, T. 1998. *Vibrio carchariae* causes mass mortalities in Japanese abalone, *Sulculus diversicolor supertexta*. *Fish Pathology* 33: 495-502.
- Novoa, B., Balseiro, P. and Figueras, A. 2004. Molecular detection of a haplosporidian parasite in carpet shell clam *Ruditapes decussatus* from Spain. *Diseases of Aquatic Organisms* 61: 89-93.
- Novoa, B., Posada, D. and Figueras, A. 2005. Polymorphisms in the sequences of *Marteilia* internal transcribed spacer region of the ribosomal RNA genes (ITS-1) in Spain: genetic types are not related with bivalve hosts. *Journal of Fish Diseases* 2: 331–338.
- Park, K.I., Ngo, T.T.T., Choi, S.D., Cho, M. and Choi, K.S. 2006. Occurrence of *Perkinsus olseni* in the Venus clam *Protothaca jedoensis* in Korean waters. *Journal of Invertebrate Pathology* 93: 81-87.
- Peeler, J. E., Reese, R. A., Cheslett, D. L., Geoghegan, F., Power, A. and Trush, M. A. 2012. Investigation of mortality in Pacific oysters associated with Ostreid herpesvirus-1 μ Var in the Republic of Ireland in 2009. *Preventive Veterinary Medicine* 105: 136-143.
- Perkins, F.O. and Wolf, P.H. 1976. Fine structure of *Marteilia sydneyi* sp. n. - Haplosporidian pathogen of Australian oysters. *Journal of Parasitology* 62: 528–538.
- Pichot, Y., Comps, M., Tige, G., Grizel, H., and Rabouin, M.A. 1979. Recherches sur *Bonamia ostreae* gen. n., sp. n., parasite nouveau de l'huitre plate *Ostrea edulis* L. *Revue des Travaux de l'Institut des Peches Maritimes* 43: 131-140.
- Ragone Calvo, L.M., Calvo, G.W. and Bureson, E.M. 2003. Dual disease resistance in a selectively bred eastern oyster, *Crassostrea virginica*, strain tested in Chesapeake Bay. *Aquaculture* 220: 69–87.

- Ramilo, A., Carrasco, N., Reece, K.S., Valencia, J.M., Grau, A., Aceituno, P., Rojas, M., Gairin, I., Furones, M.D., Abollo, E. and Villalba, A. 2015. Update of information on perkinsosis in NW Mediterranean coast: identification of *Perkinsus* spp. (Protista) in new locations and hosts. *Journal of Invertebrate Pathology* 125: 37-41.
- Reece, K.S., Dungan, C.F. and Bureson, E.M. 2008. Molecular epizootiology of *Perkinsus marinus* and *P. chesapeakei* infections among wild oysters and clams in Chesapeake Bay, USA. *Diseases of Aquatic Organisms* 82: 237-248.
- Renault, T., Le Deuff, R.-M., Cochenec, N. and Maffart, P. 1994. Herpesviruses associated with mortalities among Pacific oyster, *Crassostrea gigas*, in France – comparative study. *Revue de Médecine Vétérinaire* 145: 735-742.
- Renault, T., Moreau, P., Faury, N., Pepin, J.-F., Segarra, A. and Webb, S. 2012. Analysis of clinical Ostreid Herpesvirus 1 (*Malacoherpesviridae*) specimens by sequencing amplified fragments from three virus genome areas. *Journal of Virology* 86: 5942-5947.
- Renault, T., Stokes, N.A., Chollet, B., Cochenec, N., Berthe, F., Gérard, A. and Bureson, E.M. 2000. Haplosporidiosis in the Pacific oyster *Crassostrea gigas* from the French Atlantic coast. *Diseases of Aquatic Organisms* 42: 207-214.
- Roque, A., Carrasco, N., Andree, K. B. Lacuesta, B., Elandalousi, L. Gairin, I., Rodgers, C. J. and Furones, M. D. 2012. First report of OsHV-1 microvar in Pacific oyster (*Crassostrea gigas*) cultured in Spain. *Aquaculture* 324-325, 303-306.
- Sabry, R.C., Rosa, R.D., Magalhaes, A.R.M., Barracco, M.A., Gesteira, T.C.V. and da Silva, P.M. 2009. First report of *Perkinsus* sp. infecting mangrove oysters *Crassostrea rhizophorae* from the Brazilian coast. *Diseases of Aquatic Organisms* 88: 14-23.
- Samain, J.F., Boudry, P., Degremont, L., Soletchnik, P., Ropert, M., Nicolas, J.L., Le Roux, L. Renault, T., Burgeot, T. and Bacher C. 2004. Summer mortality in the Pacific Oyster *Crassostrea gigas*, overview of 3-year results of the cooperative “MOREST” Project. *Journal of Shellfish Research* 23: 309-310.
- Sanil, N.K., Suja, G., Lijo, J. and Vijayan, K.K. 2012. First report of *Perkinsus beihaiensis* in *Crassostrea madrasensis* from the Indian subcontinent. *Diseases of Aquatic Organisms* 98: 209-220.
- Saulnier, D., De Decker, S., Haffner, P., Cobret, L., Robert, M. and Garcia, C. 2010. A large-scale epidemiologicval study to identify bacteria pathogenic to Pacific oyster *Crassostrea gigas* and correlation between virulence and metalloprotease-like activity. *Microbial Ecology* 59: 787-798.
- Sauvage, C., Pépin, J.F., Lapègue, S., Boudry, P. and Renault, T. 2009. Ostreid herpes virus 1 infection in families of the Pacific oyster, *Crassostrea gigas*, during a summer mortality outbreak: difference in viral DNA detection and quantification using real-time PCR. *Virus Research* 142: 181-187.
- Sawabe, T., Inoue, S., Fukui, Y., Yoshie, K., Nishihara, Y. and Miura, H., 2007. Mass mortalities of Japanese abalone *Haliotis diversicolor supertexta* caused by *Vibrio harveyi* infection. *Microbes and Environments* 22: 300-308.
- Schikorski, D., Renault, T., Saulnier, D., Faury, N., Moreau, P. and Pepin, J.-F. 2011. Experimental infection of Pacific oyster *Crassostrea gigas* spat by ostreid herpesvirus 1: demonstration of oyster spat susceptibility. *Veterinary Research* 42: 27-40.
- Segarra, A., Pepin, J.F., Arzul, I., Morga, B., Faury, N. and Renault, T. 2010. Detection and description of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Research* 153: 92-95.
- Shimahara, Y., Kurita, J., Kiryu, I., Nishioka, T., Yuasa, K., Kawana, M., Kamaishi, T. and Oseko, N. 2012. Surveillance of Type 1 Ostreid Herpesvirus (OsHV-1) variants in Japan. *Fish Pathology* 47: 129-136.

- Sugumar, G., Nakai, T., Hirata, Y., Matsubara, D. and Muroga, K. 1998. *Vibrio splendidus* biovar II as causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Diseases of Aquatic Organisms* 33: 111-118.
- Travers, M.A., Le Goic, N., Huchette, S., Koken, M. and Paillard, C., 2008. Summer immune depression associated with increased susceptibility of the European abalone, *Haliotis tuberculata* to *Vibrio harveyi* infection. *Fish and Shellfish Immunology* 25: 800-808.
- Van Banning, P. 1977. *Minchinia armoricana* sp. nov. (Haplosporida), a parasite of the European flat oyster, *Ostrea edulis*. *Journal of Invertebrate Pathology* 30: 199-206.
- Van Banning, P. 1990. The life cycle of the oyster pathogen *Bonamia ostreae* with a presumptive phase in the ovarian tissue of the European flat oyster, *Ostrea edulis*. *Aquaculture* 84 : 189-192.
- Vilela, H. 1951. Sporozoaires parasites de la Palourde, *Tapes decussatus* (L.). *Rev. Fac. Ciencias, Lisboa, Sér C*, 1, 379-386.
- Villalba, A., Mourelle, S.G., López, M.C., Carballal, M.J. and Azevedo, C. 1993. Marteiliasis affecting cultured mussels *Mytilus galloprovincialis* of Galicia (NW Spain). I. Etiology, phases of the infection, and temporal and spatial variability in prevalence. *Diseases of Aquatic Organisms* 16: 61-72
- Villalba, A., Iglesias, D., Ramilo, A., Darriba, S., Parada, J.M., No, E., Abollo, E., Molares, J. and Carballal, M.J. 2014. Cockle *Cerastoderma edule* fishery collapse in the Ría de Arousa (Galicia, NW Spain) associated with the protistan parasite *Marteilia cochillia*. *Diseases of Aquatic Organisms* 109: 55-80.
- Virvilis, C. and Angelidis, P. 2003. Presence of the parasite *Marteilia* sp. in the flat oyster (*Ostrea edulis* L) in Greece. *Aquaculture* 259: 1-5.
- Wood, J.L. and Andrews, J.D. 1962. *Haplosporidium costale* (Sporozoa) associated with a disease of Virginia oysters. *Science* 136: 710-711.
- Wolf, P. H. 1972. Occurrence of a haplosporidian in Sydney rock oysters (*Crassostrea commercialis*) from Moreton Bay, Queensland, Australia. *Journal of Invertebrate Pathology* 19: 416-417.
- Wolf, P.H. 1979. Life cycle and ecology of *Marteilia sydneyi* in the Australian oyster, *Crassostrea commercialis*. *Marine Fisheries Review* 41: 70-72.
- Yanin, L., Kang, H.-S., Hong, H.-K., Jeung, H.-D., Kim, B.-K., Le, T.C., Kim, Y.-O. and Choi, K.-S. 2013. Molecular and histological identification of *Marteilioides* infection in Suminoe oyster *Crassostrea ariakensis*, Manila clam *Ruditapes philippinarum* and Pacific oyster *Crassostrea gigas* on the south coast of Korea. *Journal of Invertebrate Pathology* 114: 277-284.

Author contact information

Anders Alfjorden

National Veterinary Institute
75189 Uppsala, Sweden
anders.alfjorden@sva.se

Stephen Feist

Centre for Environment, Fisheries &
Aquaculture Science (CEFAS)
Barrack Road, The Nothe
Weymouth, Dorset DT4 8UB, UK
Stephen.feist@cefasc.co.uk

Marlene Areskog

National Veterinary Institute
75189 Uppsala, Sweden
marlene.areskog@sva.se

Susan Ford

Rutgers University
6959 Miller Avenue
Port Norris, NJ 08349, USA
sford@hsrl.rutgers.edu

David Bruno

University of Aberdeen
Aberdeen AB24 2TZ, Scotland, UK
david.bruno@abdn.ac.uk

Simon Jones

Fisheries and Oceans Canada
3190 Hammond Bay Road
Nanaimo, BC, V9T 6N7, Canada
simon.jones@dfo-mpo.gc.ca

Ryan Carnegie

Virginia Institute of Marine Science
P.O. Box 934
Gloucester Point
Virginia 23062, USA
carnegie@vims.edu

Atle Lillehaug

Norwegian Veterinary Institute
P.O. Box 750 Sentrum
0106 Oslo, Norway
atle.lillehaug@vetinst.no

Deborah Cheslett

Marine Institute
Oranmore
Co Galway H91 R673, Ireland
deborah.cheslett@marine.ie

Lone Madsen

Technical University of Denmark
National Veterinary Institute
Bülowsvej 27, 1870 Frederiksberg C,
Denmark
loma@vet.dtu.dk

Tristan Renault

The French Research Institute for Exploitation of the Sea (IFREMER)

La Tremblade 17390, France

tristan.renault@ifremer.fr

Pia Vennerström

Finnish Food Safety Authority (EVIRA)

Mustialankatu 3

00790 Helsinki, Finland

pia.vennerstrom@evira.fi

Neil Ruane

Marine Institute

Oranmore

County Galway H91 R673, Ireland

neil.ruane@marine.ie