

**Biological effects of contaminants:
Determination of CYP1A-dependent mono-oxygenase activity in dab
by fluorimetric measurement of EROD activity**

R. STAGG and A. MCINTOSH

Fisheries Research Services
The Marine Laboratory
PO Box 101
Victoria Road
Aberdeen, AB11 9DB
United Kingdom

International Council for the Exploration of the Sea
Conseil International pour l'Exploration de la Mer

Palægade 2-4 DK-1261 Copenhagen K Denmark

September 1998

ISSN  2707-6997
ISBN 978-87-7482-279-0

<https://doi.org/10.17895/ices.pub.5049>

For permission to reproduce material from *ICES Techniques in Marine Environmental Sciences*,
please apply to the General Secretary.

TABLE OF CONTENTS

Section	Page
ABSTRACT.....	I
1 INTRODUCTION.....	1
1.1 Principle of the Method	2
2 PROTOCOLS.....	4
2.1 General Applicability of the Method.....	4
2.2 Fish Sampling	4
2.3 Dissection.....	5
2.4 Homogenization	5
2.5 Assay Conditions	6
2.6 Purity of Resorufin Standards.....	7
2.7 Normalization of Enzyme Activity to Protein and Wet Weight of Tissue	9
2.8 Expression of Results.....	9
2.9 Sources of Error	10
2.10 Quality Control	13
3 REFERENCES	13

**Biological effects of contaminants:
Determination of CYP1A-dependent mono-oxygenase activity in dab
by fluorimetric measurement of EROD activity**

R. Stagg and A. McIntosh

Stagg, R., and McIntosh, A. 1998. Biological effects of contaminants: Determination of CYP1A-dependent mono-oxygenase activity in dab by fluorimetric measurement of EROD activity. ICES Techniques in Marine Environmental Sciences, No. 23.

Abstract

This paper describes a method for the determination of cytochrome P4501A (CYP1A) in fish liver by the measurement of 7-ethoxyresorufin *O*-deethylase (EROD) activity. The proposed method is by fluorescent assay of resorufin using internal standardization. The method is specifically for measurements made in dab (*Limanda limanda* L.), but is suitable for adaptation to other species by small changes in assay conditions. The principle of the method, the sampling requirements, the assay procedures and the reporting of the results are described. Sources of error and quality control procedures are also specified.

Key Words: CYP1A, EROD, fish, dab (*Limanda limanda*), biological effects of contaminants

1 INTRODUCTION

Cytochrome P450-dependent mono-oxygenases (CYP P450) are membrane-bound enzymes which metabolize a range of exogenous (organic contaminants) and endogenous (steroids, fatty acids) substrates. They are ubiquitous throughout the animal kingdom, but in fish are often present at higher concentrations in tissues such as the liver. CYP P450s are Phase I enzymes which catalyse the insertion of an oxygen atom into a xenobiotic (foreign) molecule to facilitate conjugation to an endogenous substrate. The terminal oxygenase of this enzyme system is an iron-containing haemoprotein, namely, cytochrome-P450. The system is called P450 because the wavelength absorption maximum, after reduction with carbon monoxide, is at 450 nm. The conjugation reactions are carried out by a second group of enzymes (Phase II) such as uridinediphosphate (UDP)-glucuronyl transferase (UDP-GT) and glutathione S-transferase (GST). These combined reactions have the effect of making neutral, lipid-soluble xenobiotic compounds more water soluble, thus facilitating their elimination from the body. Although this sequence of events is generally a detoxification mechanism, it can result in the activation of relatively inert chemicals to highly reactive and damaging intermediates.

In fish, one form of CYP P450 termed CYP1A is induced by environmental exposure to a diverse range of planar molecules, including polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs), dioxins and dibenzofurans (Spies *et al.*, 1982; Luxon *et al.*, 1987; Addison and Edwards, 1988; Jimenez *et al.*, 1988; Payne *et al.*, 1988; Stegeman *et al.*, 1988; Van Veld *et al.*, 1990; Stein *et al.*, 1992; Stagg *et al.*, 1995). The enzymes responsible for this activity belong to the 3-methylcholanthrene inducible family of cytochrome P450 and in fish there appears to be only a single gene (CYP1A1) in this family (Nebert and Gonzalez, 1987; Nelson *et al.*, 1993). To date, this gene has only been fully characterized in two fish species (scup (*Stenotomus chrysops*) and rainbow trout (*Oncorhynchus mykiss*)) and, therefore, it is only in these species that the enzyme can truly be termed CYP1A1. It has been recommended (Stegeman, 1992) that, until the gene is fully sequenced in other fish species, the general terminology CYP1A be used.

The basis for the use of CYP1A in monitoring is the phenomenon of induction, whereby levels of the enzyme are increased by the synthesis of new protein following exposure and the levels of enzyme activity are measurably increased. The induction of CYP1A can therefore be used to monitor exposure to bioavailable planar contaminants and, for example, is of value in determining exposure to PAHs, such as are found in hydrocarbon discharges associated with the oil industry (Davies *et al.*, 1984; Stagg *et al.*, 1995).

Stegeman (1993a, 1993b) and Stegeman and Hahn (1994) provide recent reviews of fish P450s, and Goksøyr and Forlin (1992), Stegeman *et al.* (1992), and Livingstone (1993) have reviewed the application of CYP1A as a monitoring tool.

Induction of CYP1A has also been proposed to be a measure of deleterious biological effects in the sense that compounds such as PAHs can be activated to compounds which cause DNA damage (Stegeman, 1987). There is considerable support from field observations correlating CYP1A induction to contaminant levels, DNA damage, hepatic carcinogenesis, and other pathological conditions (see reviews by Livingstone, 1993, and Livingstone *et al.*, 1994). Many steps in this cascade, for example, the effects of PAHs on CYP1A induction, DNA damage, and carcinogenesis, have also been substantiated by experimental studies. Induction of CYP1A activity can therefore be seen as an essential process involved in the aetiology of tumour initiation and carcinogenesis by the activation of compounds such as PAHs into pre-carcinogens. Whether or not the reactive intermediates produced in Phase I reactions will cause damage depends in part on the capacity of Phase II enzymes to conjugate and detoxify the compounds formed. However, there are considerable differences between fish species in this regard, which partly explains

species susceptibility to carcinogenesis (Collier *et al.*, 1992). Therefore, although the mechanistic link between elevated CYP1A activity and deleterious biological effects is well established, there is still a need for considerable care in interpreting CYP1A induction.

CYP1A mono-oxygenase activity is most conveniently measured as 7-ethoxyresorufin *O*-deethylase (EROD) activity and the method described here is that originally developed by Burke and Mayer (1974). 7-Ethoxyresorufin is an artificial substrate but is used because the assay is simple, sensitive, highly specific for CYP1A and also presents a low hazard to the operator. Aryl hydrocarbon hydroxylase (AHH) was used in many earlier studies, but it is catalysed by a broader spectrum of P450s. A previous TIMES leaflet (Galgani and Payne, 1991) gives details of a microplate method using a fluorescent plate reader. Although advantageous from the point of view of being able to process many samples very quickly, it suffers from high levels of variability and the lack of internal standardization, and is not widely used.

The present document describes a method for use with conventional fluorimeters, employing internal standardization. Two recent interlaboratory comparisons of EROD activity measurements (Munkittrick *et al.*, 1993; Stagg and Addison, 1994) showed the importance of developing a standard methodology to allow comparable data to be obtained from different laboratories, both in terms of the absolute levels of EROD activity reported and in terms of the ability to detect differences between induced and non-induced groups of fish.

1.1 Principle of the Method

Microsomal mono-oxygenase activity is the overall activity of a multi-enzyme system residing in the sarcoplasmic reticulum which comprises an NADPH reductase and multiple cytochrome P450s. Other components involved in the reaction are membrane lipids, cytochrome B₅ reductase and cytochrome b₅. NADPH reductase is a flavoprotein which accepts electrons from NADPH and passes them on to P450. The cytochrome P450 component provides the system with its substrate specificity, which in the case of CYP1A is selective for planar aromatic molecules. 7-Ethoxyresorufin *O*-deethylase (EROD) reacts exclusively with CYP1A and is converted from 7-ethoxyresorufin to resorufin by the deethylation of the hydroxyl group on the *para* position (Figure 1).

To measure enzyme activity, the liver is dissected from the fish, a sub-sample is homogenized in an appropriate buffer and then centrifuged usually to yield a 10,000 × g supernatant (the 'S9 fraction'). Many investigators prefer to work with a more refined sample in which the S9 fraction is further centrifuged to yield a 100,000 × g pellet which is resuspended in buffer ('microsomal preparation'). However, preparation of this refined sample would exclude measurements on research vessels and restrict the application of the technique in monitoring the marine environment. Therefore, only the use of S9 fractions will be described here.

EROD is measured in the S9 fraction in the presence of NADPH using 7-ethoxyresorufin (ethoxyphenoxazone) as substrate; the product, resorufin, is determined fluorimetrically (Burke and Mayer, 1974). The reaction can be followed in the reaction cuvette because the characteristic emission spectra of 7-ethoxyresorufin (λ_{Ex} 470 nm; λ_{Em} 560 nm) and resorufin (λ_{Ex} 535 nm; λ_{Em} 585 nm) show not only good separation of the maxima, but also the relative fluorescence intensity for resorufin compared to that of 7-ethoxyresorufin is at least one order of magnitude greater (Prough *et al.*, 1978). An important source of variability in EROD measurements (Munkittrick *et al.*, 1993; Stagg and Addison, 1994) is the purity of the resorufin used to calibrate the assays. Therefore, resorufin standards should be made up taking into account the purity of the resorufin stocks used. This is most conveniently done by reference to




Figure 1. The microsomal 7-ethoxyresorufin O-deethylase (EROD) reaction.

the extinction coefficient of resorufin (Klotz *et al.*, 1983) and adjusting the weight of the standard used to account for the purity measured.

2 PROTOCOLS

2.1 General Applicability of the Method

This method has been written specifically for the dab (*Limanda limanda* L.). This species has been selected because it is the preferred species for monitoring offshore under the Oslo and Paris Commissions' Joint Assessment and Monitoring Programme (JAMP). The method is equally appropriate for other flatfish species such as the flounder (*Platichthys flesus* L.) or plaice (*Pleuronectes platessa* L.). Reaction conditions for these species should be checked before the method is applied universally since it has been reported (Munkittrick *et al.*, 1993) that sub-optimal conditions (e.g., inadequate 7-ethoxyresorufin concentrations) have been used by some workers when methods have been transposed from one species to another without validating the conditions. Within any particular national or international programme, identical methodology should be used. However, this does not preclude studies on other species, in different geographical regions with slightly different methods. Variations in method are tolerable with regard to the buffers used, homogenization, normalization processes, method of freezing, and type of assay as long as they are standardized in any particular programme.

2.2 Fish Sampling

Many intrinsic and extrinsic factors affect EROD activity in the liver of fish. Differences between sexes and the effect of seasonal variation due to differences in the state of sexual maturation are well documented (Edwards *et al.*, 1988; Förlin and Haux, 1990; Goksøyr *et al.*, 1992; Larsen *et al.*, 1992). EROD is a temperature-dependent reaction which is carried out at a specified temperature (usually 20 °C) but because of the adaptation of fish to ambient temperatures, strong correlations between EROD activity and environmental temperature, independent of the assay temperature, have been observed (e.g., off the Dutch coast (Sleiderink *et al.*, 1995)). Therefore, the sampling design must either avoid or account for the temperature effect. Disease and associated liver damage may suppress EROD activity; for example, fish with overt signs of lesions can have a reduced level of EROD activity (see Figure 2). Inhibition or inactivation of EROD activity has been reported at high concentrations of some contaminants, particularly heavy metals and some PCB congeners (e.g., George, 1989; Skaare *et al.*, 1991). In some cases, there have also been reports of enhanced activity following exposure to certain combinations of contaminants (e.g., aromatic hydrocarbons and resin acids (Croce *et al.*, 1995)). Although in 90 % of field studies (to date) using fish as a monitoring species, the use of EROD alone has been successful (Livingstone, 1993), it is prudent also to measure levels of CYP1A protein by ELISA (enzyme-linked immunosorbent assay) or western blot, particularly if negative results are found. Sampling strategies should attempt to take account of these factors in any monitoring programme, and the following guidelines should be applied to the sampling:

- 1) The sex of the fish sampled should be standardized or separate analyses should be reported for males and females.
- 2) Gonad-somatic index (GSI, $100 \times \text{weight of gonad} / \text{weight of fish}$) should be calculated and reported as a measure of the state of maturity.
- 3) Animals should not be collected during the spawning period.

- 4) The bottom temperature (for flatfish) should be measured at the time of capture of the fish and reported for each group of fish analysed.
- 5) Healthy fish with no external signs of disease should be sampled.
- 6) Fish should be sampled outside their normal migration periods.
- 7) The sample size should be selected to give a coefficient of variation that is less than 10 % of the mean from the log-transformed data. Normally this would require a minimum of ten individuals of each sex at every station.
- 8) The length of the fish should be within the size range 19–24 cm, if possible.
- 9) A reference population from a pristine, uncontaminated environment should be used.
- 10) If negative results are found, the use of western blots or ELISA techniques is recommended to check that the lack of response is not due to de-activation of the enzyme.

2.3 Dissection

Samples of live fish should be obtained and each fish should be processed, within 10 minutes of killing, up to the point where the sample is cryopreserved or homogenized. The liver is most conveniently removed by dissection from the blind side of the fish. There is some evidence that biliary metabolites may inhibit EROD activity and, therefore, the liver should be dissected free without contaminating the tissue with bile (Hodson *et al.*, 1991). Since the liver is a heterogeneous tissue, it is also prudent to standardize on the lobe or portion of the lobe removed for assay. Care should be taken to ensure that the tissue is kept cold at all times during dissection. Cryopreserved samples should be freeze-clamped in liquid nitrogen and either kept in liquid nitrogen or, more conveniently, transferred to a -70°C freezer.

2.4 Homogenization


All procedures should be carried out at $0-4^{\circ}\text{C}$. A small piece (200–500 mg) of fresh or cryopreserved liver is homogenized on ice in 1–2.5 ml homogenizing buffer (Table 1) using a Potter-Elvehjem homogenizer at 700 rpm.

The homogenates should be transferred to polyethylene eppendorf tubes and centrifuged at $10,000 \times g$ for 20 minutes at 4°C . An aliquot of the supernatant is then aspirated and transferred to a fresh eppendorf tube, kept on ice and either assayed immediately or after storage at -70°C . Care should be taken not to aspirate the fat layer overlying the supernatant, which is seen in some samples. These S9 supernatants should contain approximately $5-10 \text{ mg protein ml}^{-1}$.

Table 1. Homogenizing buffer for the determination of EROD activity.

Reagent	Concentration
$\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$ (pH 7.5)	100 mM
EDTA	1 mM
Dithiothreitol	1 mM
KCl	150 mM

Figure 2. Specific activity ($\text{nmol min}^{-1} \text{mg protein}^{-1}$) of 7-ethoxyresorufin *O*-deethylase (EROD) and 7-ethoxycoumarin *O*-deethylase (ECOD) in livers of female dab (*Limanda limanda*) with and without overt hepatic tumours.



2.5 Assay Conditions

The reaction rate will be determined by the conditions under which the assays are carried out and will be affected by factors such as: pH, ionic strength and type of buffer; the concentrations of enzyme, substrate and co-factors; and the temperature of the assay. For most temperate fish species, including dab, the temperature optima are usually between 20 °C and 25 °C and the pH optimum is 7.8. The effects of changes in the concentrations of NADPH and 7-ethoxyresorufin on EROD activity in liver from the dab are shown in Figure 3. The data show that in both induced and non-induced fish, concentrations of NADPH and 7-ethoxyresorufin of 0.25 mmol l^{-1} and 2.0 $\mu\text{mol l}^{-1}$, respectively, give maximal activity.

A temperature-controlled (20 °C) cuvette should be used with continuous stirring using a small magnetic flea. The assays should be performed in a 2 ml reaction volume comprising: 1.96 ml of buffer to give a final concentration of 100 mM $\text{K}_2\text{HPO}_4/\text{KH}_2\text{PO}_4$ (pH 7.5) and 100 mM KCl, 10 μl of 7-ethoxyresorufin in DMSO (dimethylsulphoxide) to give a final concentration of 2 $\mu\text{M l}^{-1}$, and 20 μl of S9 supernatant (see Table 2). NADPH (10 μl) in distilled water, to give a final concentration of 0.25 mM, is added to start the reaction and the rate of resorufin production is measured fluorimetrically over the first minute. NADPH is prepared fresh daily.

To measure true reaction rates, the instrument sensitivity should be set to give a full-scale deflection of approximately 1.0–1.5 nmol resorufin and reaction rates measured corresponding to resorufin production up to a maximum of 0.25–0.5 nmol min^{-1} . Higher reaction rates regularly exhibit non-linear kinetics and/or result in underestimation of the true reaction rate, especially in induced animals (see below). In such cases, it is necessary to dilute the S9 supernatant in a homogenizing buffer until an appropriate reaction rate is achieved. Due to the quenching effect of

variable amounts of protein, the assay is calibrated internally by the addition of an appropriate amount of resorufin in 10 μl of phosphate buffer. Typically, this would be 0.125 nmoles but may vary depending on the instrument settings required for any particular sample. Fluorescence of resorufin should be measured by excitation at 535 nm and recording the emission at 580 nm.

Stock solutions of both 7-ethoxyresorufin and resorufin should be made by dissolving weighed amounts of crystalline solid in DMSO and stored at 4 °C in the dark. Working solutions of these stocks should be prepared fresh daily by serial dilution into either DMSO or 0.1 M phosphate buffer (pH 8.0) for 7-ethoxyresorufin and resorufin, respectively. Both resorufin and 7-ethoxyresorufin degrade rapidly on exposure to light. These compounds should therefore be stored in the dark and protected from strong light during use. The preparation of standards and solutions should be carried out in subdued light.

Table 2. Reagent volumes and concentrations used in the determination of 7-ethoxyresorufin *O*-deethylase (EROD) activity in dab (*Limanda limanda*) liver.

Reagent	Reagent concentration	Volume	Final concentration
Buffer	102.0 mM	1.96 ml	100 mM
7-ethoxyresorufin*	0.4 mM	10 μl	2.0 μM
NADPH	100.0 mM	10 μl	0.25 mM
S9 Supernatant	5–10 mg protein ml^{-1}	20 μl	50–100 $\mu\text{g ml}^{-1}$

*Dissolved in DMSO

2.6 Purity of Resorufin Standards

The purity of commercially obtained resorufin varies widely depending on source, batch and storage conditions (Munkittrick *et al.*, 1993; Stagg and Addison, 1994). Since this will have a marked influence on the level of EROD activity reported, it is vital that the purity of the resorufin used to calibrate the assay is taken into account in calculating enzyme activity. Pure resorufin has an absorption maximum of 572 nm at pH 8.0 in 0.1 M phosphate buffer with a molar absorbance of 73.2 $\text{mM}^{-1} \text{cm}^{-1}$ (Klotz *et al.*, 1983). Calculation of the weight of resorufin required to make up standards should take into account the purity of the resorufin used and the standards should be made up on the basis of the absorbance at 572 nm.



The following equation should be used:

$$A = \epsilon CL$$

where A = the absorbance at 572 nm
 ϵ = the extinction coefficient (0.0732 $\mu\text{M}^{-1} \text{cm}^{-1}$ in phosphate buffer, pH = 8.0)
 C = the concentration (μM)
 L = the path length (cm).

For example, for a 25 μM working solution of resorufin, the absorbance A_{572} in a 1-cm cuvette should be 1.83 ± 0.04 . The absorbance of the actual stock solutions (in 0.1 M phosphate buffer at pH 8.0) used to calibrate the assays should therefore be measured at 572 nm, and reported with the EROD measurements as the molar absorbance in $\text{mM}^{-1} \text{cm}^{-1}$.

Figure 3. The dependence of hepatic 7-ethoxyresorufin *O*-deethylase (EROD) activity in induced (100 mg kg⁻¹ β -naphthoflavone, intraperitoneal injection) and control dab (*Limanda limanda*) on the concentration of (a) NADPH and (b) 7-ethoxyresorufin in the reaction medium.



2.7 Normalization of Enzyme Activity to Protein and Wet Weight of Tissue

Enzyme activity can be related either to the protein content of the S9 fraction or to the wet weight of the tissue used in the reaction (i.e., per mg protein or per gram wet weight tissue). The latter has advantages in that it eliminates the possibility of a change in EROD activity due to changes in unrelated background protein. However, the argument for using protein is that in some monitoring situations, particularly when carrying out EROD measurements at sea, it is not possible to weigh accurately small pieces of tissue. Both Munkittrick *et al.* (1993) and Stagg and Addison (1994) demonstrated that protein determination was a major source of differences in the EROD values reported from different laboratories. Variation can be due to differences in the method or standards used and it is, therefore, imperative that a standard protein assay is used. The most widely used traditional protein determination is the Lowry assay (Lowry *et al.*, 1951), but many workers now use the Coomassie blue dye-binding method (Bradford, 1976), which is less expensive, faster and can be performed on a 96-well microplate. However, the latter method has poor linearity, is prone to interference, and is also markedly affected by the standards used. Recently, a modified protein assay, based on the method by Lowry *et al.* (1951), has been introduced (BioRad Laboratories or Pierce) which is fast, has much improved linearity compared to the Bradford method, and is suitable for use with 96-well microplates. This method should be used with bovine serum albumin (BSA) as a standard.

S9 supernatants are diluted to give an expected protein concentration in the range 0.5 to 1.0 mg ml⁻¹ protein. Five µl of samples or standards (0.2–1.6 mg ml⁻¹ BSA) is pipetted into a microtitre plate, 25 µl of alkaline copper tartrate reagent is added to each, followed by 200 µl Folin reagent (BioRad Laboratories). The plate is mixed and the absorbance read at 750 nm after 15 minutes incubation.

2.8 Expression of Results

The output of a typical EROD assay is shown in Figure 4A and the method of determining the rate of resorufin production is presented in Figure 4B. In an active sample, the amount of fluorescence increases after the addition of NADPH, but often the rate decreases with time. The rate measured should be as close as possible to the initial linear rate (a) shown in Figure 4B. The amount of resorufin produced over time is computed from the rate of increase in fluorescence divided by the increase in fluorescence produced by the addition of a known standard amount of resorufin (b) in Figure 4B. Occasionally, traces are produced in which the fluorescence dips immediately after the addition of NADPH (Figure 4C). In such cases, the trace should be read on the linear part of the curve. In measurements with S9 supernatants, a rate may also be observed prior to the addition of NADPH (Figure 4D) which is most likely due to the presence of endogenous NADPH in the sample.

Results should be presented normalized to the protein content of the S9 fraction and reported as nanomoles or picomoles of resorufin liberated for one minute per mg of protein (nmol min⁻¹ mg protein⁻¹). Table 3 shows the specific activity obtained from S9 supernatants of male and female dab liver injected with model inducers and from sites sufficiently contaminated with hydrocarbons to cause induction.

Table 3. Range of specific activities of 7-ethoxyresorufin *O*-deethylase in S9 supernatants from the liver of the dab (*Limanda limanda* L.) under different treatments.

Fish pretreatment	Sex	Specific Activity (nmol min ⁻¹ mg protein ⁻¹)
Laboratory controls	M	0.1–0.6
	F	0.04–0.4
β-Naphthoflavone	M	2.0–3.0
	F	0.7–2.0
Control field site	M	1.0–2.0
	F	0.1–0.5
Oil-contaminated field site	M	3.0–5.0
	F	0.7–1.0


2.9 Sources of Error

The following sources of error have been identified and should be taken into account:

- 1) Sample integrity
 - contamination of the sample by bile salts during dissection;
 - deterioration during storage or processing owing to an increase in temperature;
 - deterioration due to long-term storage.
- 2) Sample preparation
 - changes in the speed and duration of centrifugation;
 - variation in type of homogenizer and the speed and duration of homogenization;
 - preparation of fractions other than an S9 supernatant.
- 3) Assay conditions
 - deviation from assay conditions (temperature, pH and substrate concentrations);
 - deterioration of substrate and co-factors;
 - deterioration of resorufin standards;
 - differences in the purity of the resorufin;
 - protein determination (method, standards used, etc.);
 - graphical errors in the interpretation of fluorescence data.

Figure 4. Fluorescent output during hepatic 7-ethoxyresorufin *O*-deethylase EROD assays from the dab (*Limanda limanda*).

A) Typical EROD trace showing the increase in fluorescence with time and the addition (▼) of an aliquot of resorufin to calibrate the assay.



B) The same trace as in (A) to show the rate measurement (a) and the calibrant spike (b) used to compute the rate of resorufin production.






Figure 4. Continued.

C) A typical trace demonstrating an initial decline in fluorescence values after the addition of NADPH to start the reaction. Resorufin is added at the arrow (▼).



D) Trace to show the effects of endogenous NADPH after the addition of 7-ethoxyresorufin at arrow (a) and resorufin at arrow (b).



2.10 Quality Control

Laboratories should build appropriate quality control into their measurements; the development of quality control charts for all aspects of the analysis is encouraged. The following sources of error are particularly relevant to the measurement of EROD activity and appropriate steps to control these errors should be taken.

1) Pipetting

Adjustable volume pipettes must not be used to dispense less than 10% of their maximum volume.

All pipetting must be done using pipettes which are within their external calibration/service period.

All pipettes should be checked gravimetrically (or using special capillary tubes) at least once a month or after they have been dropped or jolted.

2) Resorufin standards

The molar absorbance of the working solutions used to make up the standards should be reported with each batch of assays performed.

Daily calibration blanks should be run by replacing the protein in the assay with buffer and the spike height noted for each of the standard concentrations used to calibrate the assays.

3) In-house standards

Once a week, or with each batch of samples run, samples of known EROD activity should be included as a check on the integrity of the assay and the instrumentation. In the authors' laboratory, a stock of β -naphthoflavone-induced salmon microsomes, aliquoted into small vials, is used for this purpose.

3 REFERENCES

- Addison, R.F., and Edwards, A.J. 1988. Hepatic microsomal mono-oxygenase activity in flounder *Platichthys flesus* from polluted sites in Langesundfjord and from experimental mesocosms experimentally dosed with diesel oil and copper. *Marine Ecology Progress Series*, 46: 51–54.
- Bradford, M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72: 248–251.
- Burke, M.D., and Mayer, R.T. 1974. Ethoxyresorufin: direct fluorimetric assay of microsomal O-dealkylation which is preferentially inducible by 3-methylcholanthrene. *Drug Metabolism and Disposition*, 2: 583–588.
- Collier, T.K., Singh, S.V., Awasthi, Y.C., and Varanasi, U. 1992. Hepatic xenobiotic metabolizing enzymes in two species of benthic fish showing different prevalences of contaminant-associated neoplasms. *Toxicology and Applied Pharmacology*, 113: 319–324.

- Croce, B., Stagg, R.M., and Smith, M.A. 1995. Effects of exposure to a combination of β -naphthoflavone and resin acids on biotransformation enzymes in Atlantic salmon parr (*Salmo salar*). *Aquatic Toxicology*, 32: 161–176.
- Davies, J., Bell, J., and Houghton, C. 1984. A comparison of levels of hepatic hydrocarbon hydroxylase in fish caught close to and distant from North Sea oil fields. *Marine Environmental Research*, 14: 23–45.
- Edwards, A.J., Addison, R.F., Willis, D.E., and Renton, K.W. 1988. Seasonal variation of hepatic mixed function oxidases in winter flounder (*Pseudopleuronectes americanus*). *Marine Environmental Research*, 26: 299–309.
- Förlin, L., and Haux, C. 1990. Sex differences in hepatic cytochrome P-450 monooxygenase activities in rainbow trout during an annual reproductive cycle. *Journal of Endocrinology*, 124: 207–213.
- Galgani, F., and Payne, J.E. 1991. Biological effects of contaminants: Microplate method for measurement of ethoxyresorufin-O-deethylase (EROD) in fish. *ICES Techniques in Marine Environmental Sciences*, No. 13. 11 pp.
- George, S.G. 1989. Cadmium effects on plaice liver xenobiotic and metal detoxication systems; dose response. *Aquatic Toxicology*, 15: 303–310.
- Goksøyr, A., and Förlin, L. 1992. The cytochrome P450 system in fish: aquatic toxicology and environmental monitoring. *Aquatic Toxicology*, 22: 287–312.
- Goksøyr, A., Larsen, H.E., Blom, S., and Forlin, L. 1992. Detection of cytochrome P450 1A1 in North Sea dab liver and kidney. *Marine Ecology Progress Series*, 91: 83–88.
- Hodson, P.V., Kloepper-Sams, P.J., Munkitterick, K.R., Lockhart, W.L., Metner, D.A., Luxon, P.L., Smith, I.R., Gagnon, M.M., Servo, M., and Payne, J.F. 1991. Protocols for measuring mixed function oxygenases of fish liver. *Canadian Technical Report on Fisheries and Aquatic Sciences*, 1829. 51 pp.
- Jimenez, B.D., Burtis, L.S., Ezell, G.H., Egan, B.Z., Lee, N.E., Beauchamp, J.J., and McCarthy, J.F. 1988. The mixed-function oxidase system of bluegill sunfish, *Lepomis macrochirus*: correlation of activities in experimental and wild fish. *Environmental Toxicology and Chemistry*, 7: 53–59.
- Klotz, A.V., Stegeman, J.J., and Walsh C. 1983. An aryl hydrocarbon hydroxylating hepatic cytochrome P-450 from the marine fish *Stenotomus chrysops*. *Archives of Biochemistry and Biophysics*, 226: 578–592.
- Larsen, H.E., Celander, M., and Goksøyr, A. 1992. The cytochrome P450 system of Atlantic salmon (*Salmo salar*): II. Variations in hepatic catalytic activities and isozyme patterns during an annual reproductive cycle. *Fish Physiology and Biochemistry*, 10: 291–301.
- Livingstone, D.R. 1993. Biotechnology and pollution monitoring: use of molecular markers in the aquatic environment. *Journal of Chemical Technology and Biotechnology*, 57: 195–211.

- Livingstone, D.R., Förlin, L., and George, S. 1994. Molecular biomarkers and toxic consequences of impact by organic pollution in aquatic organisms. *In* Water quality and stress indicators in marine and freshwater systems: linking levels of organisation, pp. 154–171. Ed. by D.W. Sutcliffe. Freshwater Biological Association, Ambleside, UK.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randall, R.J. 1951. Protein measurement with a Folin phenol reagent. *Journal of Biological Chemistry*, 193: 265–275.
- Luxon, P.L., Hodson, P.V., and Borgmann, U. 1987. Hepatic aryl hydrocarbon hydroxylase activity of lake trout (*Salvelinus namaycush*) as an indicator of organic pollution. *Environmental Toxicology and Chemistry*, 6: 649–657.
- Munkittrick, K.R., van den Heuvel, M.R., Metner, D.A., Lockhart, W.L., and Stegeman, J.J. 1993. Interlaboratory comparison and optimisation of hepatic microsomal ethoxyresorufin O-deethylase activity in white sucker (*Catostomus commersoni*) exposed to bleached kraft pulp mill effluent. *Environmental Chemistry and Toxicology*, 12: 1273–1283.
- Nebert, D.W., and Gonzalez, F.J. 1987. P-450 genes: structure, evolution and regulation. *Annual Review of Biochemistry*, 56: 945–993.
- Nelson, D.R., Kamataki, T., Waxman, D.G., Guengerich, F.P., Estabrook, R.W., Feyereisen, R., Gonzalez, F.J., Coon, M.J., Gunsalus, R.C., Gotoh, O., Okuda, K., and Nebert, D.W. 1993. The P450 superfamily: update on new sequences, gene mapping, accession numbers, early trivial names of enzymes, and nomenclature. *DNA and Cell Biology*, 12: 1–51.
- Payne, J.F., Kiceniuk, J., Fancey, L., Williams, U., Fletcher, G.L., Rahimtula, A., and Fowler, B. 1988. What is a safe level of polycyclic aromatic hydrocarbons for fish: subchronic study on winter flounder (*Pseudopleuronectes americanus*). *Canadian Journal of Aquatic Science*, 45: 1983–1993.
- Prough, R.A., Burke, M.D., and Mayer, R.T. 1978. Direct fluorimetric methods for measuring mixed function oxidase activity. *Methods in Enzymology*, 52: 372–377.
- Skaare, J.U., Jensen, E.G., Goksøyr, A., and Egaas, E. 1991. Response of xenobiotic metabolising enzymes of rainbow trout (*Oncorhynchus mykiss*) to the mono-ortho substituted polychlorinated PCB congener 2,3',4,4',5-pentachlorobiphenyl, PCB-118, detected by enzyme activities and immunochemical methods. *Archives of Environmental Contamination and Toxicology*, 20: 349–352.
- Sleiderink, H.W., Beyer, J., Everaats, J.M., and Boon, J.P. 1995. Influence of temperature on cytochrome P450 1A in dab (*Limanda limanda*) from the southern North Sea: results from field surveys and a laboratory study. *Marine Environmental Research*, 39: 67–71.
- Spies, R.B., Felton, J.S., and Dillard, L. 1982. Hepatic mixed function oxidases in California flatfishes are increased in contaminated environments by oil and PCB injection. *Marine Biology*, 70: 117–127.
- Stagg, R.M., and Addison, R.A. 1994. An interlaboratory comparison of measurements of ethoxyresorufin O-deethylase activity in dab (*Limanda limanda*) liver. *Marine Environmental Research*, 38: 93–108.

- Stagg, R.M., McIntosh, A., and Mackie, P. 1995. The induction of hepatic monooxygenase activity in dab (*Limanda limanda* L.) in relation to environmental contamination with petroleum hydrocarbons in the northern North Sea. *Aquatic Toxicology*, 33: 245–264.
- Stegeman, J.J. 1987. Monooxygenase systems in marine fish. *In* Pollutant studies in marine animals, pp. 65–92. Ed. by C.S. Giam and L.E. Ray. CRC Press, Boca Raton, Florida.
- Stegeman J.J. 1992. Nomenclature for hydrocarbon-inducible cytochrome P450 in fish. *Marine Environmental Research*, 34: 133–138
- Stegeman, J.J. 1993a. The cytochromes P450 in fish. *In* Biochemistry and Molecular Biology of Fishes, 2. Molecular Biology Frontiers, pp. 137–158. Ed. by P.W. Hochachka and T.P. Mommsen. Elsevier, Amsterdam.
- Stegeman, J.J. 1993b. Cytochrome P450 forms in fish. *In* Cytochrome P450, pp. 279–291. Ed. by J.B. Schenkman and H. Greim. Springer-Verlag, Berlin.
- Stegeman, J.J., and Hahn, M. 1994. Biochemistry and molecular biology of mono-oxygenases: current perspectives on forms, functions and regulation of cytochrome P450 in aquatic species. *In* Aquatic toxicology: molecular, biochemical and cellular perspectives, pp. 87–206. Ed. by D. Malins and G. Ostrander. Lewis Publishers, Boca Raton, Florida.
- Stegeman, J.J., Woodin, B.R., and Goksøyr, A. 1988. Apparent cytochrome P-450 induction as an indication of exposure to environmental chemicals in flounder *Platichthys flesus*. *Marine Ecology Progress Series*, 46: 55–60.
- Stegeman, J.J., Brouwer, M., DiGiulio, R.T., Förlin, L., Fowler, B.A., Sanders, B.M., and Van Veld, P.A. 1992. Molecular responses to environmental contamination: enzyme and protein systems as indicators of chemical exposure and effect. *In* Biomarkers: Biochemical, Physiological and Histological Markers of Anthropogenic Stress. Ed. by R.J. Huggett, R.A. Kimmerle, P.M. Merhle Jr., and H.L. Bergman. Lewis Publishers, Michigan, USA.
- Stein, J.E., Collier, T.K., Reichert, W.L., Casillas, E., Hom, T., and Varanasi, U. 1992. Bioindicators of contaminant exposure and sublethal effects: studies with benthic fish in Puget Sound, Washington. *Environmental Toxicology and Chemistry*, 11: 701–714.
- Van Veld, P.A., Westbrook, D.J., Woodin, B.R., Hale, R.C., Smith, C.L., Huggett, R.J., and Stegeman, J.J. 1990. Induced cytochrome P-450 in intestine and liver of spot (*Leiostomus xanthurus*) from a polycyclic aromatic hydrocarbon contaminated environment. *Aquatic Toxicology*, 17: 119–132.

ICES Techniques in Marine Environmental Sciences

- No. 1 Cadmium and lead: Determination in organic matrices with electrothermal furnace atomic absorption spectrophotometry
- No. 2 Trace metals in sea water: Sampling and storage methods
- No. 3 Cadmium in marine sediments: Determination by graphite furnace atomic absorption spectroscopy
- No. 4 Lipophilic organic material: An apparatus for extracting solids used for their concentration from sea water
- No. 5 Primary production: Guidelines for measurement by ^{14}C incorporation
- No. 6 Control procedures: Good laboratory practice and quality assurance
- No. 7 Suspended particulate matter: Collection methods for gravimetric and trace metal analysis
- No. 8 Soft bottom macrofauna: Collection and treatment of samples
- No. 9 Sediments and suspended particulate matter: Total and partial methods of digestion (*videotape available*)
- No. 10 Organic halogens: Determination in marine media of adsorbable, volatile, or extractable compound totals
- No. 11 Biological effects of contaminants: Oyster (*Crassostrea gigas*) embryo bioassay
- No. 12 Hydrocarbons: Review of methods for analysis in sea water, biota, and sediments
- No. 13 Biological effects of contaminants: Microplate method for measurement of ethoxyresorufin-O-deethylase (EROD) in fish
- No. 14 Temporal trend monitoring: Introduction to the study of contaminant levels in marine biota
- No. 15 Temporal trend monitoring: Contaminant levels in tissues of Atlantic cod
- No. 16 Benthic communities: Use in monitoring point-source discharges
- No. 17 Nutrients: Practical notes on their determination in sea water
- No. 18 Contaminants in marine organisms: Pooling strategies for monitoring mean concentrations
- No. 19 Common diseases and parasites of fish in the North Atlantic: Training guide for identification
- No. 20 Temporal trend monitoring: Robust method for analysing contaminant trend monitoring data
- No. 21 Chlorobiphenyls in marine sediments: Guidelines for determination
- No. 22 Biological effects of contaminants: Cholinesterase inhibition by organophosphate and carbamate compounds
- No. 23 Biological effects of contaminants: Determination of CYP1A-dependent mono-oxygenase activity in dab by fluorimetric measurement of EROD activity