9.4.14 Advice June 2011

ECOREGION Widely distributed and migratory stocks STOCK Orange roughy (Hoplostethus atlanticus) in the Northeast Atlantic

Advice for 2012

The advice given in 2010 for orange roughy assessment units is biennial and valid for 2011 and 2012 (see ICES, 2010). Therefore, ICES advices that no directed fisheries for this species and measures to minimize bycatch should be taken.

Sources

ICES. 2010. Orange Roughy (*Hoplostethus atlanticus*) in the Northeast Atlantic. Report of the ICES Advisory Committee, 2010. ICES Advice, 2010. Book 9. 211–216pp.

Table 9.4.14.1 Orange roughy in the Northeast Atlantic. ICES advice, management, and landings.

Year	ICES advice	Predicted	TAC	TAC	TAC	Total	ICES
		catch	EU	EU	EU	TAC	Landings
		corresp. to	Subarea I	Subarea	Subarea	EU	
		advice	- V, VIII	VI	VII	I-XII and	
			- XII and			XIV	
			XIV				
2003	Exploitation strictly limited and	-	=	0.088	1.349	1.437	0.9
	populations closely monitored.						
	No direct fishery in Subarea VI						
2004	Biennal	-	-	0.088	1.349	1.437	1.3
2005	No fishery unless accompanied	0	0.102	0.088	1.148	1.338	0.7
	by programmes to collect data						
2006	Biennial	0	0.102	0.088	1.148	1.338	0.7
2007	No direct fishery, bycatch in	0	0.044	0.051	0.193	0.288	0.2
	mixed fishery as low as possible						
2008	Biennial	0	0.03	0.034	0.13	0.194	0.2
2009	No direct fishery, bycatch in	0	0.015	0.017	0.065	0.097	0.08
	mixed fishery as low as possible						
2010	Biennial	0	0	0	0	0	
2011	No directed fisheries and	0	0	0	0	0	
	measures to minimize bycatch						
2012	No new advice, same as 2011	0	0	0	0	0	

Weights in '000 t.

ICES Advice 2011, Book 9