

Stock Annex: Blackspot seabream (*Pagellus bogaraveo*) in Subarea 9 (Atlantic Iberian waters)

Stock specific documentation of standard assessment procedures used by ICES.

Stock: Blackspot seabream

Working Group: Working Group on Biology and Assessment of Deep-sea Fisheries Resources (WGDEEP)

Created:

Authors:

Last updated: April 2016

Last updated by: Juan Gil Herrera – WGDEEP

A. General

A.1. Stock definition

Stock limits are generally determined not only by biological considerations but also by agreed boundaries and coordinates. ICES considered three different components for this species: a) Areas 6, 7, and 8; b) Area 9, and c) Area 10 (Azores region). This separation does not pre-suppose that there are three different stocks of red sea bream, but it offers a better way of recording the available information" (ICES, 2007). The inter-relationships of the red sea bream from Areas 6, 7, and 8, and the northern part of Area 9.a, and their migratory movements within these areas have been observed by tagging methods (Gueguen, 1974). However, there is no evidence of movement to the southern part of 9.a where the main fishery currently occurs. Tagging has been done also in the Strait of Gibraltar area, where the majority of the fishery currently occurs. No significant movements are reported, although local migrations are also observed: feeding grounds are distributed along the entire Strait of Gibraltar and the species seems to remain in this area as a resident population (Gil, 2006). In 2007, Piñera *et al.* suggests no significant genetic differences are present along Spanish coasts (Mediterranean and Atlantic areas).

Stock structure of the species in ICES Subarea 9 still unknown. Besides, in the case of the Strait of Gibraltar, Red sea bream also inhabit in Morocco waters. In fact recaptures of tagged fish were also notified by Moroccan fishermen.


A.2. Fishery

Although *Pagellus bogaraveo* is caught by Spanish and Portuguese fleets in Subarea 9, only a more complete description of one of the fisheries has been provided to the Working Group, the corresponding to the Spanish fishery in the southern part of Subarea 9, close to the Strait of Gibraltar.

The majority of landings on deep-water species at mainland Portugal are conducted by the artisanal fleet, mainly longline fisheries. These operated in the Portuguese continental slope and located in ports as Peniche, Sesimbra and Sagres. Red sea bream landings reflect a seasonal activity probably related with a larger availability of the species or market demands that lead fishermen to spend some time targeting this species (I. Figueiredo, *pers. com.*).

In relation to the Spanish fishery in the southern ICES Subarea 9a, an updated description of it has been presented to the Working Group by Gil *et al.* (WD to the 2016 WGDEEP), that complete the information offered in the previous WGs (Gil *et al.*, 2000, 2003, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014 and 2015; Gil and Sobrino, 2001, 2002 and 2004). This artisanal longline fishery targeted Red sea bream has been developed along the Strait of Gibraltar area. In recent years this fishery covers more than the 70% of the landings for the species in the Subarea 9. The base and landing ports are two: Algeciras and mainly Tarifa (Cádiz, SW Spain). The “*voracera*”, a particular mechanised hook and line baited with sardine, is the gear used by the fleet (Table A.2.1). The mean technical characteristics of this fleet by port are 8.95 and 6.52 meters length and 5.84 and 4.0 tons G.T.R. for Tarifa and Algeciras, respectively (Gil *et al.*, 2000). Currently around 60 boats are involved in the fishery. Fishing grounds are located at both sides of the Strait of Gibraltar and quite close to the main ports (Figure A.2.1). Fishing is carried out taking advantage of the turnover of the tides in depths from 200 to 400 fathoms. Landings are distributed in categories due to the wide range of sizes and to market reasons (these categories have varied in time but from 2000 onwards still the same).

Table A.2.1. Red sea bream Spanish fishery of the Strait of Gibraltar: Fleet and gear summary descriptive.

FLEET ID	GEAR TYPE	N° BOATS	NUMBER OF LINES	HOOK TYPE AND SIZE	MEAN SOAKTIME	EFFORT (DAYS AT SEA)
LHM_DEF	Vertical mechanized handline ("voracera")	±60	Maximum of 30 lines per day (each line attached a maximum of 100 hooks, usually ±70)	L=3.95±0.39 cm S=1.40±0.14 cm 	±30 min	Maximum 140 days

From 2002 onwards artisanal boats from other port, Conil, have begun to direct its fishing activity to *P. bogaraveo* in different fishing grounds and with different fishing gear

(longlines) than the “*voracera*” fleet boats. Nowadays, only around 9 boats are developing this fishery.

In addition, Moroccan longliners have been fishing in the Strait of Gibraltar area since 2001. Around 102 boats are mainly based in Tangier and their average technical characteristics are: 20 GRT, 160 CV and about ten years old. Moreover, 435 artisanal boats (± 15 CV, ≤ 2 GRT and 4–6 m length) also target this species in the Strait of Gibraltar area (S. Benchoucha, *pers.com.*). The WG considers the account of Moroccan data appropriate as the fishery operates in the same area as the Spanish fishery and obviously targets the same stock. Unfortunately, no updated information was available in 2015 and no new information from the Moroccan fishery has been received in the last three years.




Figure A.2.1. Red sea bream Spanish fishery of the Strait of Gibraltar: Yearly soaking positions footprints from observers on-board programme (from Gil *et al.*, WD 19 to the 2011 ICES WGDEEP).

It should be noted that not all the catches/landings from Strait of Gibraltar do not come exclusively from ICES Subarea 9: however it was considered from the same stock unless

the fishing area is placed between different Regional Organizations/Commissions (ICES, GCFM and CECAF) borders (Figure A.2.2).




Figure A.2.2. Red seabream Spanish “*voracera*” fishery of the Strait of Gibraltar: spatial distribution of landings in 2014 (from Gil *et al.*, 2016)

Detailed information from Portuguese fisheries has been provided to the Working Group by Araújo *et al.* (WD to the WGDEEP 2016). As well as in other Spanish places in Subarea 9, it is admitted that there is no target fishery towards Red (blackspot) seabream in Portugal mainland: the species is usually caught as by-catch of fisheries targeting other species. The majority of deep-water species landings as fresh fish in mainland Portugal correspond to the polyvalent fleet, which uses mainly longlines, while landings from trawlers are the second more relevant. The main landing ports ($\approx 89\%$ of the species mainland Portugal total landings) from North to South are: Matosinhos, Aveiro, Nazaré, Peniche, Sesimbra and Sagres. Among those, higher landings were registered in the southern ones (Peniche, Sagres and Sesimbra), possibly due to a higher abundance of the species in the nearby fishing grounds.

Araújo *et al.* (WD to the WGDEEP 2016) presents information of Red (blackspot) seabream spatial distribution from Portuguese research surveys. In continental Portugal this species distributes along the coast, but has been preferentially caught in the southern waters. It seems to have a patchy distribution; occurring predominantly in the same areas over the years were the groundfish survey take place (Figure A.2.3).




Figure A.2.3. Red (blackspot) seabream in ICES Subarea 9: Species spatial distribution along the Portuguese slope based on the Portuguese Autumn Groundfish Survey (PT-GFS) information between 1990 and 2015 (no survey was conducted in 2012) (from Araújo *et al.*, WD to the 2016 WGDEEP).

A.3. Ecosystem aspects

Red sea bream is a benthic-pelagic species that inhabits various types of bottom (rock, sand, and mud) down to a depth of 900 m. It is found in the Northeast Atlantic, from South of Norway to Cape Blanc, in the Mediterranean Sea, and in the Azores, Madeira and Canary Archipelagos (Desbrosses, 1938; Pinho and Menezes, 2005). Hareide (2002) reported also occasional occurrence of this species along the Mid-Atlantic Ridge (north and south of the Azores).

Feeding habit of this species has been little studied. Morato *et al.* (2001) describes the diet of *Pagellus bogaraveo* and *Pagellus acarne* in the Azores and Olaso and Pereda (1986) describe the diet of 22 demersal fish in the Cantabrian Sea including *Pagellus bogaraveo*. In the Strait of Gibraltar fishery, feeding studies presents the difficulty of the use of bait (sardine), which should be ignored to describe the feeding habit of the species. A total of 1106 red sea bream stomachs contents were analysed: 725 stomachs were empty and 381 were fullness. Vacuity index (VI) was 66%. The trophic spectrum is composed of 24 prey taxa, 9 orders, eleven families and 15 species and genera are represented. Despite the trophic spectrum diversity observed, the overall diet is not very diverse. Red sea bream in the Strait of Gibraltar has only a main prey, *Sergia robusta* (J. Gil, *pers.com.*).

Main red sea bream predators are unknown in the Strait of Gibraltar waters but maybe dolphins' (personal communication from Ceuta veterinary) and Bluefin tuna's (personal communication from fishermen) predation should be taken into account. Studies in Azores (Gomes *et al.*, 1998) cite that *Conger conger*, *Raja clavata* and *Galeorhinus galeus* must be considered as potential predators (all three species are present in Strait of Gibraltar area). Besides, other considerations as the competition for food of Red sea bream with other big predators in the Strait of Gibraltar (i.e. Bluefin tuna, marine mammals, etc.) and those related with environmental parameters (variation and long-term atmospheric oscillations) might affect the stock status as well as its fisheries.

Deep-sea coral ecosystems represent true biodiversity hot spots. OSPAR identified cold-water coral ecosystems as one of the most vulnerable ecosystems where action is required now to mitigate further loss of biodiversity. Figure 2 shows the deep-water coral occurrences in the Strait of Gibraltar.




Figure A.3.1. Coral distribution in the Strait of Gibraltar (adapted from Álvarez-Pérez *et al.* in Freiwald and Roberts, Eds. 2005). Yellow points correspond to “voracera” fleet fishing grounds from observers on board programme. Legend refers to percentage cover of coral.

B. Data

B.1. Commercial catch

In Subarea 9, catches, most of them taken by lines, correspond to Spain and Portugal. Spanish landings data from this area are available from 1983 and Portuguese from 1988 onwards. The maximum catch in this period was obtained in 1993–1994 and 1997 (about 1000 t). Catches in 2009 amount to 718 t, but decreases again (484 t) over the last years (180 t in 2013, the minimum of the available time-series). Moroccan information shows a continuous increasing trend in landings from 2001 to 2011.

Almost all Spanish catches in this area are taken in waters close to the Gibraltar Strait. Until 2002 they were restricted to two ports (Tarifa and Algeciras), but from 2002 significant catches were obtained also by artisanal Spanish boats of a third port (Conil) in different fishing grounds of the same area. An increasing trend in landings was observed but since 2008 it only rates an average of 15 t, lower than in the early years.

In the Portuguese landings no clear tendency is observed. The maximum values took place in 1988 (370 t) and in 1998 (357 t) and the minimum one in 2000 (83 t). In 2013 landings was 90 t.

Without the Moroccan landings, compared to the minimum from 2013 last year (2015) landings increased more than the 160% in the whole Subarea 9. In addition Gil *et al.* (WD to the WGDEEP 2016) reported more than the 252 % in the Strait of Gibraltar fishery. Again, it should be noted that not every Spanish landings from the Strait of Gibraltar come from ICES Subarea 9. So, last years landings are presented above. Note that Strait of Gibraltar ones (from Spanish fishery) are assigned to three difernet areas.

Year	Portugal	Spain	Morocco	Unallocated	TOTAL
2010	105	379	146		630
2011	74	259	154		487
2012	143	60	146	92	295
2013	90	91	118		180
2014	59	203	n/a		262
2015	66	87 (142*)	n/a		153 (295*)

***Figures in brackets includes Red (blackspot) seabream from other areas (FAO 34.1.11. and FAO 37.1.1): 153 t in ICES 9, 64 t in FAO 34.1.11 and 78 t in FAO 37.1.1 for the Total (without Morocco) 2015 landings.**

Length frequencies of landings are only available for the Spanish red sea bream fishery in the Strait of Gibraltar. There was a decrease of the mean size from 1995 to 1998. It is necessary to point out that the red sea bream may have a variable length distribution depending on its geographic and bathymetric distribution, as suggests the different mean length of landings measured in ports (Tarifa and Algeciras). The mean length of the landings increased steadily in both ports from 1999 onwards then decreased but has been increasing again between 2006 and 2009. The mean length from both landing ports declined in 2010. However the median value is lower than the mean since 1995, and very close to the minimum landing size in Algeciras. Only one mean value (in 1998) is lower than the 2013 year's mean landing size. Median values are well below the mean in recent years. However, changes are small and gradual. There seem to be a long-term decline, but the mean length has been mostly stable over the last decade (Figure 13.1.3).




Figure B.1.1. Spanish “voracera” Red (blackspot) sea bream fishery of the Strait of Gibraltar: 1983–2015 landings mean length distribution (from Gil *et al.*, WD to the 2016 WGDEEP).

A Kolmogorov-Smirnoff test reflects significant differences ($p < 0.05$) between the length distributions from Spain and Morocco (Belcaid *et al.*, WD 20 to the 2011 ICES WGDEEP) and also within Spain (Gil *et al.*, WD 19 to the 2011 ICES WGDEEP). Differences among the sampling protocols may be the explanation to the observed difference. In Moroccan and Spanish observers’ programme the sampling covers certain the boats (random sampling) while in the Spanish first sale fish market the sampling covers the four market categories (stratified sampling). So raising the random sampling weight to the total landings did not take into account the difference due to the variability of the length composition related to bathymetric distribution of the species and the stratified sampling seems to be more appropriate.

B.2. Biological

Red sea bream is a protandric hermaphrodite species changing from males to females. Red sea bream have a low productivity and they change sex as they age, starting as males and becoming females between ages 4 and 6. Measures to ensure balanced exploitation between younger fish (males) and older fish (females) are essential.

An annual reproductive cycle has been described for the species in this area (Gil, 2006). The spawning season seems to take place during the first quarter of the year. The smallest specimens are mainly males, maturing at a $L_{50} = 30.15$ cm. At about 32.5 cm in total length,

an important percentage of individuals change sex and became females, maturing at $L_{50}=35.73$ cm. Thus, from age 5 all individuals can be considered mature, whether they are males or females.

Red sea bream is considered a slow growing species. A combined ALK was obtained by three agreed readings from 1497 otoliths collected from 2003 to 2008 (Gil *et al.*, 2009). It comprises lengths from 24 to 54 cm and ages between 3 and 10, but it has not been validated yet. According to the available information the maximum age recorded in Subarea 9 is ten years. However, the ages of older fish may be underestimated and it is possible that this species may be slower growing and longer-lived than current studies indicate. In fact, there was one recapture from tagging surveys notified more than ten years after its release (J. Gil, *pers. com.*). Table 2.1 presents different estimates of von Bertalanffy Growth Function (VBGF) parameters available from otoliths readings or tag-recapture data.

Table 2.1. Red sea bream of the Strait of Gibraltar: VBGF parameter estimates.

AUTHORS	STUDY AREA	METHODOLOGY	T ₀	K	L _∞
Sobrino and Gil, 2001	Strait of Gibraltar	Otoliths reading	-0.67	0.169	58.00*
Gil <i>et al.</i> , 2008	Strait of Gibraltar	Otoliths reading	-1.23	0.169	62.00*
Gil <i>et al.</i> , 2009	Strait of Gibraltar	Otoliths reading	-0.34	0.162	62.00*
Gil <i>et al.</i> , 2008	Strait of Gibraltar	Recaptures ⁽¹⁾		0.079	62.00*
Gil <i>et al.</i> , 2008	Strait of Gibraltar	Recaptures ⁽²⁾		0.098	62.00*
Gil <i>et al.</i> , 2008	Strait of Gibraltar	Recaptures ⁽³⁾		0.161	62.00*
Gil <i>et al.</i> , 2008	Strait of Gibraltar	Recaptures ⁽⁴⁾		0.080	62.00*

⁽¹⁾Gulland y Holt, 1959 ⁽²⁾Munro, 1982 ⁽³⁾Fabens, 1965 ⁽⁴⁾Appeldoorn, 1987.

* F9ed (from the largest observed sample).

Padillo *et al.* (WD17 to the 2011 WGDEEP) present new information based on Discriminant Analysis of several of the samples used to make the ALK, combining morphometric and morphological variables to re-estimate red sea bream ages. The reclassification success percentage was 85.3%, well above from the 70% adopted by other authors (Palmer *et al.*, 2004 and Galley *et al.*, 2006). Changes in otolith shape could be related to the growth rate and be also strongly influenced by environmental components. Therefore, future work should include the analysis of such factors throughout years and cohorts.

The natural mortality of *Pagellus bogaraveo* is uncertain because there are no data available to estimate M directly. A mortality rate of 0.2 year⁻¹ has been adopted by several authors in several studies from other areas (Silva, 1987; Silva *et al.*, 1994; Krug, 1994; Pinho *et al.*, 1999; Pinho, 2003) and also by Gil (2006) for the Strait of Gibraltar.

Age and growth based on otolith readings were revised along the ICES WKAMDEEP meeting (October, 2013): No more than ten years of age was estimated from otolith readings in the Strait of Gibraltar area but two recaptures from the tag-recapture programme have remained at sea for more than ten years (J. Gil, *pers. com.*). Moreover, growth estimates from tag-recapture experiments suggest that otolith readings may

underestimate age and that some hyaline rings are uncounted and/or missing. The use of biased age estimates may have important consequences.

B.3. Surveys

Only tagging surveys were carried out in the Strait of Gibraltar area. Several tagging surveys (56 days at sea in 2001, 2002, 2004, 2006 and 2008) have been conducted in the Strait of Gibraltar area. 4500 fish were tagged and 404 recaptures have been reported. No significant movements have been observed, although local migrations were noted: feeding grounds are distributed along the entire Strait of Gibraltar and the species seems to remain within this area as a resident population (Gil, 2006). Recaptures of tagged fish have also been reported by the Moroccan fishery.

Figure B.3.1 presents abundance indexes from records in Portuguese research surveys, particularly on the Portuguese crustacean surveys/Nephrops TV Surveys (PT-CTS [UWTV (FU 28-29)]) and the Portuguese Autumn Groundfish Survey (PT-GFS). Catch rate (n/h) is presented for the sector where the species was more regularly sampled (Arrifana, SW Portugal) and for the depth strata with most adequate data in each survey. Results highlight not only its patchy distribution but aggregation pattern: when sampled, the catch rates are relatively high in most cases (Araújo *et al.*, WD to the WGDEEP 2016).




Figure B.3.1. Red (blackspot) seabream in ICES Subarea 9: Catch rate of *P. bogaraveo* in the “Arrifana” sector SW Portugal), in depth strata with high catch rates from Portuguese Autumn Groundfish Survey (PT-GFS), between 1990 and 2015 (no survey was conducted in 1997, 1998 and 2012; in the remaining years this sector and depth stratum was always sampled) and Portuguese crustacean surveys/Nephrops TV Surveys (PT-CTS (UWTV (FU 28-29))) undertaken between 1997 and 2015 (no survey was conducted in 2012; this sector and depth stratum was not sampled in 1997, 1999 and 2015) (from Araújo *et al.*, WD to the 2016 WGDEEP).

The availability of Portuguese research surveys data (Araújo *et al.*, WD to the WGDEEP 2016) shows a clear increase of specimens’ length with depth. Mean length by depth strata is similar between the two Portuguese surveys (100 to 300 m depth strata). However, overall data suggest that Groundfish survey catches a greater amount of smaller specimens (Figure B.3.2).




Figure B.3.2. Red (blackspot) seabream in ICES Subarea 9: Boxplot of length distribution of *P. bogaraveo* by depth strata based on the the Portuguese Autumn Groundfish Survey (PT-GFS), between 1990 and 2015 (no survey was conducted in 2012) and the Portuguese crustacean surveys/Nephrops TV Surveys (PT-CTS (UWTV (FU 28-29))) undertaken between 1997 and 2015 (no survey was conducted in 2012) (from Araújo *et al.*, WD to the 2016 WGDEEP).

B.4. Commercial cpue

It should be noted that the effort unit from the historical series, number of sales, may be inappropriate, as it fails to consider the missing effort from boats that have not caught enough fish to go to the market. Thus, in the years this missing effort has increased substantially (fishing vessels with no catches and no sale sheet to be recorded) and its cpue values may be overestimated.

Gil *et al.* (WD 19 to the 2011 WGDEEP) presents a short series of cpue (2005–2009) from the observers' on-board programme in the red sea bream fishery of the Strait of Gibraltar. Sampling level was five boats and three trips per month. Number and length measurements of caught species were recorded. Values vary around three Red sea bream per ± 70 hooks but the general trend seems to be slightly decreasing throughout the years.

Burgos *et al.* (2013) demonstrated that a VMS system in operation since 2009 provided a reliable estimate of actual fishing effort and derived a cpue time-series. This cpue series is now used as a biomass index of the stock (see Section C below).

VMS information was updated within the 2016 WGDEEP (Figure B.4.1).




Figure B.4.1. Spanish “voracera” Red (blackspot) sea bream fishery of the Strait of Gibraltar (ICES Subarea 9): Estimated Cpue using sales sheets (dashed line) and VMS data as unit of effort (continued line) (from Gil et *al.*, WD to the 2016 WGDEEP).

B.5. Other relevant data

C. Assessment: data and method

Model used: No model was adopted for the assessment yet. Till the moment the assessments attempts were no accepted and only several trends (landings and length distributions) were used for the scientific advice.

So according to the available information, in 2014 DLS method 3.2 (ICES DLS Guidance Report 2012) was selected to estimate C_{y+1} as a reduction of C_{y-1} , applying the 20% Uncertainty Cap and the 20% Precautionary Buffer because the steep declining the catch and in the biomass index occurred over the last years.

Advice basis in 2016 was given according to the DLS 3.2 method again. Thus the quantitative advice increases in 20% (because the biomass index increase, precautionary cap) from the 2015 and 2016 advice (115 t).

Software used: None

Model Options chosen: None

Input data types and characteristics:

D. Short-term projection

Model used: None

Software used: None

Initial stock size:

Maturity:

F and M before spawning:

Weight-at-age in the stock:

Weight-at-age in the catch:

Exploitation pattern:

Intermediate year assumptions:

Stock–recruitment model used:

Procedures used for splitting projected catches:

E. Medium–term projections

Model used: None

Software used: None

Initial stock size:

Natural mortality:

Maturity:

F and M before spawning:

Weight-at-age in the stock:

Weight-at-age in the catch:

Exploitation pattern:

Intermediate year assumptions:

Stock–recruitment model used:

Uncertainty models used:

- 1) Initial stock size:
- 2) Natural mortality:
- 3) Maturity:
- 4) F and M before spawning:

- 5) Weight-at-age in the stock:
- 6) Weight-at-age in the catch:
- 7) Exploitation pattern:
- 8) Intermediate year assumptions:
- 9) Stock-recruitment model used:

F. Long-term projections

Model used: None

Software used: None

Maturity:

F and M before spawning:

Weight-at-age in the stock:

Weight-at-age in the catch:

Exploitation pattern:

Procedures used for splitting projected catches:

G. Biological reference points

	TYPE	VALUE	TECHNICAL BASIS
MSY	MSY Btrigger	N/A	
Approach	FMSY	F0.1	YpR Analysis
	Blim	N/At	
Precautionary	Bpa	N/A	
Approach	Flim	N/A	
	Fpa	N/A	

No biological reference points have been defined.

H. Other issues

H.1. Historical overview of previous assessment methods

Historical series of landings data available to the Working Group have been exploratory assessed by the WGDEEP since 2006. No discard data were available to the Working Group, but for this species this could be considered minor. The landings data used in the

assessment exercise of red sea bream in 9 included Spanish and Portuguese landings from 1990 onwards.

New assessment exercises were presented to the Group in 2011. An Extended Survivors' Analysis (XSA) attempt with the Strait of Gibraltar Spanish red sea bream fishery data is described by González and Gil (WD 18 to the 2011 WGDEEP). Belcaid *et al.* (WD 20 to the 2011 WGDEEP) presents the results obtained by a Yield-per-recruit analysis from 2005–2007 Spanish and Moroccan landings length distribution available information from the Strait of Gibraltar area.

In 2012, new assessment attempts for the Strait of Gibraltar fishery were presented to the Group by González *et al.* (WD to the 2012 WGDEEP). Simple assessment methods that use historical catches and available trend or size-composition information could potentially be applied to many data-poor stocks. The exercise, which includes two different approaches, appears to be enough indicative because this quite small area comprises more than the 80% from the total of the species in the ICES 9 Subarea:

- Depletion-Corrected Average Catch (DCAC) input parameters were: Sum of Catch along 29 years=12 723, Natural Mortality=0.2 (standard deviation of 0.5 and Lognormal distribution), F_{MSY} to $M=0.8$ (standard deviation of 0.2 and Lognormal distribution), Depletion Delta=0.47 (standard deviation of 0.3 and Lognormal distribution) and $B_{MSY}/B_0=0.4$ (standard deviation of 0.1 and Beta distribution). The number of iterations chosen was 10 000.
- Besides, from 2009–2011 available information (Spanish and Moroccan landings and length distributions) Length Cohort (LCA) and Yield-per-recruit (YpR) analysis were carried out. A plus group of 50 cm was established because the anomalous F values in larger length classes. After exploratory analysis using different F terminal random seed values (0.3–0.5–0.8) 0.5 was adopted as F terminal in the final run.

Another join assessment between Spain and Morocco took place in Tangier in November 2014: info update till 2013 and three assessment exercises (LCA, Production model and MSY Catch) using SP+MOR data were attempted (CopeMed II, 2015).

Besides, along the 2012 WGDEEP meeting several exercises were attempted:

- The Catch-MSY method (Martell and Froese) propose a new method for estimating maximum sustainable yield (MSY) from a time-series of catch data, resilience of the species, and estimations about depletion, i.e. relative stock abundances at the beginning and the end of the time-series by means of the Catch-MSY method. With the guidance and help from R. Froese a Catch MSY with the total landings in Subarea 9 was carried out.
- Two new functions within FLR (FLAdvice package) allow us the simulation of a fish stock based on its life-history parameters. From a species complete set of parameters: ages 1 to 17, VBGF growth model ($L_{inf}=62$ and $k=0.169$) and the length–weight relationship ($a=0.014$ and $b=3.014$) the functions derive in a set of biological reference points, including $F_{0.1}$ and $F_{30\%SPR}$.

- Another approach was considered along the WG to estimate BRPs using the Beverton–Holt function developed in R by Azevedo and Cadima (BHAC). As same as the previous, a set of life-history parameters and derives in an F vector (which includes F_{MAX} , $F_{0.1}$ and $F_{30\%SPR}$ for example).

And also several Biological Reference Points estimates from different approaches were available. Table below summarizes the methods adopted and its estimates:

METHOD/ESTIMATE	F _{MAX}	F _{0.1}	F _{30%SPR}	F _{40%SPR}	F _{SQ}
Gislason spreadsheet (WKLIFE) with AFC=3	0.61	0.26	0.36	0.24	-
Gislason spreadsheet (WKLIFE) with AFC=4	0.77	0.29	0.62	0.37	-
BHAC (WKLIFE)	0.39	0.17	0.20	0.14	-
FLAdvice (WKLIFE)		0.16	0.16		-
YpR Analysis (from LCA outputs)(1)	0.30	0.11		0.12	0.19

⁽¹⁾Landings from the Strait of Gibraltar only.

WKLIFE Gislason spreadsheet was applied using an L_{MAX} of 62 cm and $AFC = 3$ and 4. The parameters estimated by the mode were unrealistic based on what is known about this stock and the F_{MAX} value was substantially higher than that estimated by YPR. This may be because the underpinning empirical relationships may not apply to a protandric hermaphrodite species.

In the 2014 meeting no analytical assessment was attempted. Results from gadget exploratory analysis that was presented should be considered preliminary as the model needs a better parameterization. However it was decided the use of the cpue as a biomass index rather than the use of recent landings (like two years before). Thus, the quantitative advice was given according to the DLS 3.2 method instead of the 6.2 that appears in the ICES DLS Guidance Report 2012.

In summary, recent trends are fairly clear; despite Moroccan landings from the Strait of Gibraltar are not available in the years 2012 and 2013. It is however assumed that these landings follow a decreasing trend. Landings have declined significantly over the last four years and may be considered as a substantial reduction in exploitable biomass. Mean length distribution and lpue decreasing trends may also be consistent with overexploited population signals. However, since 2014 all signals (landings, lpue and length distribution) reverted to increase but without any evidence of its sustainability.

In 2016, an exploratory assessment (gadget model) was presented to the Group. Model definition and the estimated parameters are conditioned by the available information. It should be remarked that this preliminary model was developed only with the Spanish target fishery information (“voracera” fleet). So, the effect of the inclusion of Morocco data from the Strait of Gibraltar area is unknown but it is desirable its future incorporation to the model. More detailed information is available at the WD presented by Gil *et al.* to the 2016 WGDEEP.

I. References

- Alvarez-Pérez, G., P. Busquets, B. de Mol, N. G. Sandoval, M. Canals and J.L. Casamor. 2005. Deep-water coral occurrences in the Strait of Gibraltar. In Cold-Water corals and ecosystems. Freiwald A. and J. M. Roberts (eds.). Springer-Verlag Berlin Heidelberg: 207–221.
- Araújo, G., T. Moura and I. Figueiredo. 2016. Notes on *Pagellus bogaraveo* in the Portuguese continental waters (ICES Division 9.a). Working Document to the 2016 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Belcaid, S., J. Gil, J.L. Pérez, O. Kada, J. Baro and A. García. 2011. Summary results about the Red seabream fishery of the Strait of Gibraltar joint assessment exercise between Spain and Morocco. Working Document to the 2011 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Burgos, C., J. Gil and L.A. del Olmo. 2013. The Spanish blackspot sea bream (*Pagellus bogaraveo*) fishery in the Strait of Gibraltar: Spatial distribution and fishing effort derived from a small-scale GPRS/GSM based fisheries vessel monitoring system. Aquatic Living Resources, 26: 399–407.
- CopeMed II. 2015. Report of the third meeting of the CopeMed II Working Group between Spain and Morocco on Blackspot seabream (*Pagellus bogaraveo*) of the Strait of Gibraltar area. CopeMed II Technical Documents N°37 (GCP/INT/028/SPA – GCP/INT/006/EC). Málaga, 2015. 35pp
- Desbrosses, P. 1938. La dorade commune (*Pagellus centrodontus*) et sa pêche. Rev. Trav. Off. Pêches Marit., 5 (2): 167–222.
- Galley, E., P. Wright and F. Gibb. 2006. Combined methods of otolith shape analysis improve identification of spawning areas of Atlantic cod. ICES Journal of Marine Science, 63: 1710–1717 pp.
- Gueguen, J. 1974. Précisions sur le migrations de la dorade rosse *Pagellus bogaraveo* (Brünnich, 1768). Science et Pêche. Bull. Inst. Pêches Marit., 237: 1–9.
- Gil, J. 2006. Biología y pesca del voraz [*Pagellus bogaraveo* (Brünnich, 1768)] en el Estrecho de Gibraltar. Tesis Doctoral, Universidad de Cádiz (UCA, Spain).
- Gil, J., J.J. Acosta, C. Farias and M.M. Soriano. 2012. Updating the information about the Red seabream (*Pagellus bogaraveo*) Spanish fishery in the Strait of Gibraltar (ICES Subarea 9). Working Document to the 2012 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., J.J. Acosta, M.M. Soriano, C. Farias and C. Burgos. 2011. The Red seabream (*Pagellus bogaraveo*) fishery in the Strait of Gibraltar: ICES Subarea 9 updated data. Working Document to the 2011 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., J. Canoura, C. Burgos and C. Farias. 2005. Update of the Red seabream (*Pagellus bogaraveo*) fishery data in the Strait of Gibraltar (ICES 9a south) including biological information. Working Document to the 2005 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., J. Canoura, C. Burgos and C. Farias. 2010. The Red seabream (*Pagellus bogaraveo*) Spanish fishery in the Strait of Gibraltar: Useful information that should be considered for the ICES Subarea 9 assessment update exercise. Working Document to the 2010 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., J. Canoura, C. Burgos, C. Farias and V. Polonio. 2008. Red seabream (*Pagellus bogaraveo*) assessment of the ICES 9 from the information available of the fishery in the Gibraltar Strait.

- Working Document to the 2008 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., J. Canoura, C. Burgos and I. Sobrino. 2007. Red seabream (*Pagellus bogaraveo*) fishery of the Strait of Gibraltar (ICES 9a south): Update of the available information. Working Document to the 2007 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., S. Cerviño and B.T. Elvarsson. 2016. A preliminary gadget model to assess the Spanish Red seabream fishery of the Strait of Gibraltar. Working Document to the 2016 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., C. Farias, J. Canoura and J.J. Acosta. 2013. The Red seabream fishery in the Strait of Gibraltar: update of the available information from the fishery statistics and some considerations about the current knowledge on the target species growth. Working Document to the 2013 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., C. Farias, J. Canoura, J.J. Acosta, M. Soriano and C. Burgos. 2015. Updating the available information from Spanish Red seabream fishery in the Strait of Gibraltar. Working Document to the 2015 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., C. Farias, C. Burgos, J. J. Acosta and M. Soriano. 2016. Updating the available information from Spanish Red seabream fishery in the Strait of Gibraltar. Working Document to the 2016 ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., C. Farias, C. Burgos, J.J. Acosta and J. Canoura. 2014. The red seabream fishery in the Strait of Gibraltar: an update of the available information. Working Document to the 2014 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., I. Sobrino and M. P. Jiménez. 2000. A brief description of the Strait of Gibraltar red seabream (*Pagellus bogaraveo*) fishery. Working Document to the 2000 Report of the ICES Study Group on the Biology and Assessment of Deep-sea Fisheries Resources (SGDEEP).
- Gil, J. and I. Sobrino. 2001. New biological information about the red seabream (*Pagellus bogaraveo*) of the Strait of Gibraltar (ICES 9a). Working Document to the 2001 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J. and I. Sobrino. 2002. Update of the information about the red seabream (*Pagellus bogaraveo*) from the Strait of Gibraltar (ICES 9a south). Working Document to the 2002 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J. and I. Sobrino. 2004. Red seabream (*Pagellus bogaraveo*) fishery of the Strait of Gibraltar (ICES 9a south): Update of the information available. Working Document to the 2004 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gil, J., I. Sobrino and J. Canoura. 2003. Update of the information about the red seabream (*Pagellus bogaraveo*) fishery in the Strait of Gibraltar (ICES 9a south). Working Document to the 2003 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Gomes, T.M., E. Sola, M.P. Grós, G. Menezes and M.R. Pinho. 1998. Trophic relationships and feeding habits of demersal fishes from the Azores: importance to multispecies assessment. ICES C.M. 1998/O: 7.

- González, F. And J. Gil. 2011. Summary results from the XSA exercise attempted with the Red seabream available data from the Spanish fishery in the Strait of Gibraltar. Working Document to the 2011 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Hareide NR and Garnes G. 2001. The distribution and catch rates for deep-water fish along the Mid-Atlantic Ridge from 43 to 61°N. *Fish Res* 51:297–310.
- ICES. 2007. Report of the Working Group on the Biology and Assessment of Deep-sea Fisheries Resources. ICES CM 2007/ACFM: 28.
- Morato, T., E. Solá, M. Pitta Grós and G. Menezes. 2001. Feeding habits of two congener species of seabreams, *Pagellus bogaraveo* and *Pagellus acarne*, off the Azores (Northeastern Atlantic) during spring of 1996 and 1997. *Bulletin of marine science* 69 (3): 1073–1084.
- Olaso, I. and P. Pereda. 1986. First results of the studies about the feeding of the species of fishes accompanying the southern stock of hake. *International Council for the Exploration of the Sea, C.M.* 1986/G:34.
- Padillo, M.M., J. Gil and A.M. Juárez. 2011. Morphometric and morphological studies of Red seabream (*Pagellus bogaraveo*) otoliths from the Strait of Gibraltar: Exploratory analysis of its application for ageing. Working Document to the 2011 Report of the ICES Working Group on the Biology and Assessment of Deep-Sea Fisheries Resources (WGDEEP).
- Palmer, M., G. Pons and M. Linde. 2004. Discriminating between geographical groups of a Mediterranean commercial clam (*Chamelea gallina* (L.): Veneridae) by shape analysis. *Fisheries research* 67: 93–98 pp.
- Pinho, M. R.. 2003. Abundance estimation and management of Azorean demersal species. Tesis Doctoral. Universidad de Azores. 163 pp.
- Pinho, M.R. and Menezes, G. 2005. Azorean Deepwater Fishery: Ecosystem, Species, Fisheries and Management Approach Aspects. Deep Sea 2003: Conference on the Governance and Management of Deep-sea Fisheries, Conference Poster and Dunedin Workshop Papers. *FAO Fish. Proc.* 3/2.
- Pinho, M.R., G. Menezes and H. Krug. 1999. Estado de exploração dos recursos demersais dos Açores – Proposta de gestão. Relatório apresentado ao Workshop: “Gestão de Pescas”, Secretaria Regional de Agricultura Pescas e Ambiente, Direcção Regional das Pescas. Horta 20 de Março de 1999.
- Piñera, J., G. Blanco, E. Vázquez and J. Sánchez. 2007. Genetic diversity of blackspot seabream (*Pagellus bogaraveo*) populations off Spanish Coast: a preliminary study. *Marine Biology* 151(6): 2153–2158.
- Piñera, J., G. Blanco, E. Vázquez and J. Sánchez. 2007. Genetic diversity of blackspot seabream (*Pagellus bogaraveo*) populations off Spanish Coast: a preliminary study. *Marine Biology* 151(6): 2153–2158.
- Silva, H. M. 1987. Estado dos stocks de goraz e abrótea. In Relatório da VII Semana das Pescas dos Açores, pp.197–199. Ed. by Secretaria Regional de Agricultura e Pescas, Direcção Regional das Pescas dos Açores, Horta, Açores. 269 pp.
- Silva, H. M., H. Krug and G. Menezes. 1994. Bases para a regulamentação da pesca de demersais nos Açores. *Arquivos do DOP-Série Estudos*, 94. 99 pp.
- Krug, H. M. 1990. The Azorean blackspot seabream, *Pagellus bogaraveo* (Brünnich 1768) (Teleostei: Sparidae): Reproductive cycle, hermaphroditism, maturity and fecundity. *Cybio*, 14, 151–159.