

This Report not to be cited without prior reference to the Council^{*)}

International Council for the
Exploration of the Sea

C.M.1977/F:8
Demersal Fish (Northern) Committee



REPORT OF THE NORTH SEA ROUND FISH WORKING GROUP

Charlottenlund, 21-25 March 1977

This Report has not yet been approved by the International Council for the Exploration of the Sea; it has therefore at present the status of an internal document and does not represent advice given on behalf of the Council. The proviso that it shall not be cited without the consent of the Council should be strictly observed.

^{*)} General Secretary,
ICES,
Charlottenlund Slot,
2920 Charlottenlund,
DENMARK

REPORT OF THE NORTH SEA ROUNDFISH WORKING GROUP

| <u>Contents</u> | <u>Page</u> |
|---|-------------|
| 1. Participation | 1 |
| 2. Terms of reference | 1 |
| 3. Total Allowable Catches | 1 |
| 3.1. Recommended TACs | 2 |
| 3.2. Relationship between stocks in Sub-Area IV and Division IIIa | 2 |
| 3.3. Difficulties of controlling fishing mortality by means of a TAC | 2 |
| 4. State of exploitation | 3 |
| 5. Biology and distribution with reference to fisheries zones | 3 |
| 6. Method of determining catch predictions | 3 |
| 6.1. General | 3 |
| 6.2. Estimates of discards | 4 |
| 6.3. Natural mortality rate | 4 |
| 6.4. Yearclass strength | 4 |
| 6.5. Numbers landed | 4 |
| 6.6. Estimates of fishing mortality rate in recent years | 4 |
| 7. Haddock (Division VIb) | 5 |
| 8. Fishing Effort | 5 |
| 9. The effect on fishing mortalities of a TAC being set too high | 5 |
| 10. Notes on stock assessment and TAC calculations | 6 |
| 10.1. Cod | 6 |
| 10.1.1. North Sea (Sub-Area IV) | 6 |
| 10.1.1.1. VPA | 6 |
| 10.1.1.2. Yearclass strengths | 6 |
| 10.1.1.3. Catch predictions | 7 |
| 10.1.2. West of Scotland - Division VIa | 7 |
| 10.1.2.1. VPA | 7 |
| 10.1.2.2. Yearclass strengths | 7 |
| 10.1.2.3. Catch predictions | 7 |
| 10.1.3. Irish Sea (Division VIIa) | 7 |
| 10.1.3.1. VPA | 7 |
| 10.1.3.2. Yearclass strength | 8 |
| 10.1.3.3. Catch predictions | 8 |

| | | |
|--------------------|---|---------|
| 10.2. | Haddock | 8 |
| 10.2.1. | North Sea (Sub-Area IV) | 8 |
| 10.2.1.1. | Total international catch per age group | 8 |
| 10.2.1.2. | Mean weight at age | 8 |
| 10.2.1.3. | VPA | 8 |
| 10.2.1.4. | Yield per recruit | 9 |
| 10.2.1.5. | Catch predictions | 9 |
| 10.2.2. | West of Scotland (Division VIa) | 10 |
| 10.2.2.1. | Total international catch per age group | 10 |
| 10.2.2.2. | Mean weight at age | 10 |
| 10.2.2.3. | VPA | 10 |
| 10.2.2.4. | Yield per recruit | 10 |
| 10.2.2.5. | Catch predictions | 10 |
| 10.2.3. | Sub-Area VII | 11 |
| 10.3. | Whiting | 11 |
| 10.3.1. | North Sea (Sub-Area IV) | 11 |
| 10.3.1.1. | Total international catch per age group | 11 |
| 10.3.1.2. | Mean weight at age | 11 |
| 10.3.1.3. | VPA | 11 |
| 10.3.1.4. | Catch predictions | 12 |
| 10.3.2. | Sub-Area VI | 12 |
| 10.3.3. | Sub-Area VII | 13 |
| Tables 1.1. - 6.6. | | 14 - 54 |
| Figures 1 - 7 | | 55 - 61 |

- o - o -

Note: See also Doc. C.M.1977/F:8 - APPENDIX.

REPORT OF THE NORTH SEA ROUND FISH WORKING GROUP

1. Participation

| | |
|---------------------|-----------------------|
| D.W. Armstrong | UK (Scotland) |
| R. de Clerck | Belgium |
| N. Daan | Netherlands |
| J.P. Hillis | Ireland |
| A. Hylan | Norway |
| J. Janusz | Poland |
| J. Lahn-Johannessen | Norway |
| R. Jones (Chairman) | UK (Scotland) |
| H. Knudsen | Denmark |
| F. Lamp | Germany, Fed. Rep. of |
| G. Lefranc | France |
| C.T. Macer | UK (England) |
| V.P. Ponomarenko | USSR |
| J.G. Pope | UK (England) |
| P. Sparre | Denmark |
| I.G. Tsenker | USSR |
| G. Wagner | Germany, Fed. Rep. of |

V.M. Nikolaev, ICES Statistician, also attended the meeting.

2. Terms of Reference

At the 1976 Statutory Meeting of ICES in Copenhagen, it was decided (C.Res.1976/2:28) that:

"the North Sea Roundfish Working Group should meet at Charlottenlund from 21-25 March 1977 to:

- (a) summarise the gadoid data collected from the North Sea Young Herring Surveys;
- (b) assess TACs for 1978 for cod, haddock and whiting in Sub-Areas IV, VI and VII."

Also, as a result of a request from NEAFC, the Group was asked to provide information on the distribution, biology and state of exploitation of certain fish stocks with reference to 200 mile fishery zones.

3. Total Allowable Catches (TACs)

Total allowable catches for 1978 for different assumptions about changes in fishing effort are summarised below

TACs for 1978 (in '000 metric tons)

Option A

| <u>Sub-Area</u> | <u>Cod</u> | <u>Haddock</u> | <u>Whiting</u> |
|-----------------|------------|----------------|----------------|
| IV | 240(174) | 112(190) | 173(130) |
| VI | 21.9(9.3) | 13.4(12) | 17.8(21) |
| VII | 20.2(14) | 8.2(6.5) | 25.7(20) |

Table cont'd. ...

| <u>Sub-Area</u> | <u>Cod</u> | <u>Option B</u> | | |
|-----------------|------------|-----------------|----------------|--|
| | | <u>Haddock</u> | <u>Whiting</u> | |
| IV | 220 | 106 | 161 | |
| VI | 20.4 | 12.4 | 16.6 | |
| VII | 19.5 | 8.2 | 25.7 | |

The values in brackets under Option A are the recommended TACs for 1977. Differences between the recommended TACs for 1977 and 1978 are largely due to differences in yearclass strengths.

3.1. Recommended TACs

The TACs given under Options A and B were chosen from the predictions as the most appropriate for consideration by management.

Option A gives the TACs that should be adopted if the object is to prevent fishing mortality in 1978 from increasing above its 1976 - 1977 level.

Option B gives the TACs that should be adopted, if the object is to reduce fishing mortality in 1978 by 10% compared to the level in 1976 and 1977.

Because of uncertainties about the state of exploitation of the stocks and because, in an event, it would be inadvisable to try to reduce fishing effort too rapidly, it is recommended that TACs be chosen so as to reduce the fishing mortality rate, but by no more than 10% in the first instance (Option B).

3.2. Relationship between stocks in Sub-Area IV and Division IIIa

The Sub-Area IV catch predictions and TACs are for Sub-Area IV only, and do not include Division IIIa.

It was noted however that there is a certain amount of interchange between the stocks of cod, haddock and whiting in the North Sea and those in Division IIIa (see Appendix).

3.3. Difficulties of controlling fishing mortality by means of a TAC

The Working Group wishes to draw attention to the difficulty of controlling fishing mortality by means of a TAC:

- 1) For years with relatively large variation in recruitment in which the recruiting yearclasses make up a large proportion of the exploitable stock estimates of yearclass strength are an essential part of a catch prediction. Reliable catch predictions are impossible if average yearclass strengths have to be assumed in the assessments.

The case of Division VIa cod for example, the appearance of a good yearclass in 1974, necessitated a revision of the 1977 catch prediction to 21 000 tons (Table 6.4). This compared with a value of only 9 400 tons calculated in last year's Working Group Report (ICES, C.M.1976/F:9, Table 33), a value that was determined before it was known that the 1974 yearclass was a good one.

- 2) For fisheries with relatively large variations in discarding, in which the discards make up a significant proportion of the catch, estimates of the rate of discarding are an essential part of a catch prediction. For example, in the case of North Sea haddock and whiting, discarding rates were estimated for the Netherlands and the UK (Scotland). These showed that the quantities discarded could be particularly high at times when good yearclasses are entering Recommendation 4 fisheries. This happened in the North Sea in 1976

and in that year it was estimated that the Netherlands and the UK (Scotland) alone discarded about 40 000 tons of haddock and 34 000 tons of whiting.

Attention is drawn to a previous report (C.M.1975/F:5) in which an increase in mesh size for vessels fishing for cod, haddock and whiting in the North Sea was recommended.

An increase in mesh size would appear to be particularly appropriate in fisheries in which large-scale discarding is common practice.

4. State of exploitation

It is difficult to quantify the state of exploitation of cod, haddock and whiting stocks in Sub-Areas IV, VI and VII.

This is because criteria based on different assumptions lead to views that cannot easily be reconciled:

- 1) For some species, F values are greatly in excess of F_{max} values on yield per recruit curves (Table 5.4). According to this criterion, a number of the stocks under consideration are all seriously overexploited.
- 2) Yield per recruit curves are not necessarily the same as total yield curves however. It is therefore not certain to what extent changes in total yield would necessarily be the same as changes in yield per recruit, for changes in fishing mortality.
- 3) During the 1960's, stocks of cod, haddock and whiting in some areas and particularly in the North Sea and Division VIa increased significantly above their pre-1960 levels.

This was a consequence of good recruitment, and it is not known to what extent this, and other changes that took place in North Sea fish stocks at the time, were the result of natural processes or to what extent they were an indirect outcome of fishing.

Updated figures of cod, haddock and whiting nominal catches are given in Tables 1.1. - 1.8.

5. Biology and Distribution with reference to Fisheries Zones

An account of the distribution and biology of various species has been prepared by members of the Working Group. This is given in the Appendix to the report.

6. Method of Determining Catch Predictions

6.1. General

Catch predictions for each species were calculated using the methods described in the previous report of the Working Group (Anon 1976), i.e., for those stocks for which age composition data were available, an estimate was made of the age composition for 1976 and this was projected forward taking account of available recruitment estimates to determine values for the catches in 1977 and 1978. This method was used for the stocks of cod, haddock and whiting in Sub-Area IV and Division VIa, and also for cod in Division VIIa.

For haddock and whiting in the North Sea, account was taken of Recommendation 2 as well as Recommendation 4 fisheries when preparing the input data for the catch predictions. For the remaining stocks under consideration, precautionary TACs were calculated on the basis of previous catch predictions.

6.2. Estimates of discards

Estimates were available of the numbers of discarded haddock and whiting in Sub-Area IV by the following countries:

Haddock - Netherlands, UK(Scotland)
Whiting - Netherlands, UK(Scotland)

6.3. Natural mortality rate

For cod, haddock and whiting a constant value of $M = 0.2$ was used in the assessments.

6.4. Yearclass strength

For North Sea cod, haddock and whiting, estimates of yearclass strength were based on data obtained from the International Young Herring Surveys. These surveys showed that the 1975 yearclass appears to be poor for cod and haddock, and average for whiting. The 1976 yearclass appears to be average for cod and whiting, but poor for haddock. For cod, the 1976 yearclass, although average, is good when considered with the yearclasses prior to 1969. For cod, haddock and whiting in other areas, no direct estimates were available and average yearclass strengths were assumed in the catch predictions.

6.5. Numbers landed

Estimates of the numbers landed at each age or length were provided by the following countries:

Cod Sub-Area IV - Belgium, Denmark, France, Netherlands, Norway, Poland, UK(England and Scotland)
Division VIa - Ireland, UK(England and Scotland)
Division VIIa - Ireland, UK(England and Wales)

Haddock

Sub-Area IV - Belgium, Denmark, Netherlands, Norway, Poland, UK(England and Scotland)
Division VIa - Ireland, UK(England and Scotland)

Whiting

Sub-Area IV - Belgium, Denmark, Netherlands, Norway, Poland, UK(England and Scotland)
Division VIa - UK(Scotland) and Ireland

6.6. Estimates of fishing mortality rate in recent years

A difficulty with the estimation of fishing mortality rates is that VPA does not necessarily provide reliable estimates of these parameters for the 3 or 4 most recent years. To obtain these, additional information or assumptions are required.

One approach was to investigate the long-term relationship between fishing mortality and total international effort. For cod and whiting, the relationship obtained was good (Figs. 1 & 2). For haddock, the relationship between effort and mortality rate was

not very good and it was not possible to say from this investigation that fishing mortality had necessarily changed. (Figure 3).

7. Haddock (Division VIb)

The recorded landings of haddock from Division VIb in 1974, 1975 and 1976 were 49 000, 50 000 and 41 000 tons respectively. Prior to 1974, haddock landings from this area were about 1 000 - 2 000 tons annually.

The area of Rockall Bank where haddock are normally caught is about 2 700 square miles. The catch rates for the years 1974, 1975 and 1976 have therefore been 18.1, 18.5 and 15.2 tons per square mile annually. These catch rates are extremely high when compared with average haddock catch rates in other parts of the North Atlantic, and the Group felt that it would be unwise to continue fishing at so high an intensity.

As a precautionary measure, it is strongly recommended that a TAC of 2 000 tons should be adopted.

8. Fishing Effort (Tables 1.9 - 1.11)

Fishing effort data were available for English and Scottish vessels fishing in the North Sea and to the west of Scotland. English data were also available for the Irish Sea. In addition, other countries had some effort data but only for periods that were too short to give useful indications of trends in catch rate.

Raising the English data to total international fishing effort for cod, haddock and whiting in the North Sea, gave the time series shown in Table 1.11. These are given for the years for which VPA estimates of fishing mortality were also available.

Table 1.11 also shows the time series of international fishing effort obtained by raising Scottish effort data to total international catch for cod, haddock and whiting in the North Sea. Obvious relationships between these fishing effort series and the VPA estimates of fishing mortality were observed for North Sea cod (Figure 1) (for the English series), and for North Sea whiting for both series (Figure 2 shows the result obtained using Scottish data). North Sea haddock (Figure 3) fishing mortalities did not relate well to either the Scottish or English fishing effort series nor with an alternative series based on Danish industrial fishing effort.

Fishing effort series for cod, haddock and whiting west of Scotland did not relate well to fishing mortalities calculated by VPA, neither did fishing effort for cod in the Irish Sea.

It is possible that the measurement of fishing effort in these various areas could be improved by basing international effort estimates on more homogeneous sectors of national fishing fleets.

The Group recommended that research be carried out to see if more reliable effort series can be constructed.

9. The Effect on Fishing Mortalities of a TAC being set too high

If a TAC is overestimated, the percentage increase in fishing mortality will be greater than the percentage by which the TAC exceeds its correct level.

As an example, the 1978 TAC for the North Sea cod has been assessed at 240 000 tons (Option A). Figure 4 shows the relationship between the TAC set in 1978 and the fishing mortality in that year. The fishing mortality in 1978 is expressed as a percentage of the 1976 value. Two possible relationships are shown. The first (lower line) results from the Working Group's estimates of fishing mortality in 1976. The second (upper line) results from assuming that the 1976 fishing mortality was equal to the average in the period 1963-72. This latter assumption leads to more conservative results than does the Working Group's estimate. The figure shows that if the second relationship is correct, the TAC (for stabilising fishing mortality) should have been set at 211 000 tons. Clearly, if the Option A TAC of 240 000 tons were caught, this would lead to a 20% increase in fishing mortality in 1978. The effect on fishing mortality in 1978 of other incorrect TAC's, under either assumption, can also be determined from this figure. More generally, such graphs must always pass through the origin and rise asymptotically to a TAC level equal to the exploitable biomass of the stock. Consequently, the effect of successive unit percentage increases in TAC will produce increasingly large percentage increases in the fishing mortality.

10. Notes on Stock Assessment and TAC Calculations

10.1. Cod

10.1.1. North Sea (Sub-Area IV)

10.1.1.1. VPA (Tables 2.1, 3.1 and 4.1)

Since the fishing mortalities given in the 1975 Working Group report suggest that the change in F values with time might be small, a different approach was tried for determining terminal F values. This was done by calculating the ratios of the catches of each yearclass in consecutive years. If it is assumed that fishing mortalities on consecutive age groups in consecutive years are the same, it follows that

$$C_{a,t}/C_{a+1,t+1} = \exp(F+M), \text{ where } C_{a,t}$$

is the total catch of age group "a" in year "t". This permits F to be estimated directly. The average value obtained in this way for 7 year and older cod was 0.55, and this value has been applied as terminal F value for the oldest age group throughout the years 1963 to 1965. The total international effort for cod, as calculated from English cpue data, also indicated only minor variations over this period, but in 1976 the level of effort had apparently dropped below the average level observed before by some 25% (Figure 1). Since there is a reasonably good relationship between the F values for age groups 2-8, and the total international effort, the terminal F values in 1976 were reduced by 25% for all age groups after calculating the average fishing mortalities on each age group during the period 1963-73.

The resulting fishing mortalities from the VPA are relatively higher in the years 1971, 1972 and 1973, when fishing for cod was extremely profitable as a consequence of the presence of the two exceptionally strong 1969 and 1970 yearclasses. Apparently these yearclasses attracted much fishing effort and experienced a high mortality as a consequence. In recent years the fishing mortality appeared to have declined, as explained above.

10.1.1.2. Yearclass strengths

Predictive regressions of the VPA estimates of yearclass size were significantly correlated with IYHS abundance indices (Table 5.3.). Yearclasses 1975 and 1976 were estimated using this regression as 123×10^6 and 256×10^6 1-year-old fish

respectively. The fishing mortality on the 1975 yearclass in 1976 was adjusted to simulate the predicted yearclass strength. Yearclass 1977 entering the fishery in 1978 was assumed to be of average size (230×10^6 1-year-old fish).

10.1.1.3. Catch predictions (Tables 5.1 and 6.1)

The values of the different parameters used in the catch predictions are given in Table 6.1. Numbers landed are the provisional figures available for 1976. The F values correspond to the VPA input terminal F values. Weight at age data were the same as last year, but an adjustment of -6.26% was made to simulate the actual catch in 1976.

There seems to be no urgent need to reduce exploitation rates drastically, since there is above average recruitment and fishing mortalities are not excessively high. TACs were calculated: a) assuming that the fishing mortalities in 1976 were 25% lower than the average for the period 1963-1975, b) based on the more pessimistic view that fishing mortalities in 1976 were similar to the average values over the period 1963-1975. A TAC of 220 000 tons was selected by the Working Group as the most appropriate one for 1978.

10.1.2. West of Scotland - Division VIa

10.1.2.1. VPA (Tables 2.4, 3.4 and 4.4)

The terminal F value calculated from the catch ratios of 5 years and older cod over the entire period was 0.7. For younger age groups in 1976 the average values for each of these age groups were used.

10.1.2.2. Yearclass strengths

Since no direct estimates were available of the strengths of the yearclasses of 1975 and 1976, average values of the number of 1-year-old cod in 1966-73 were used for the catch predictions. From the catch data, the 1974 yearclass appears to be a particularly strong one.

10.1.2.3. Catch predictions (Tables 5.1 and 6.4)

Weight at age data were adjusted by +2.85% to simulate the actual catch in 1976. Catch predictions were carried out on the assumption that the average level of fishing mortality applied to the exploitation rate in 1976 and 1977 was at an average level.

The predicted catch for 1977 is 20 600 tons, which compares with a prediction in last year's Working Group report of only 9 400 tons. This increase is caused by the apparent strength of the 1974 yearclass in the 1976 catches. The latter contributed 16 000 tons instead of a predicted value of 9 700 tons, which illustrates the difficulties that can arise when assessing TACs without proper estimates of yearclass strength.

A TAC of 19 100 tons was recommended for Division VIa cod for 1978 (Table 5.1). For Division VIb cod, a precautionary TAC of 1 300 tons was recommended (Table 5.1).

10.1.3. Irish Sea (Division VIIa)

10.1.3.1. VPA (Tables 2.7, 3.7 and 4.7)

Similar procedures as for Division VIa were followed to obtain terminal F values, which were smaller than those obtained in the former Working Group report.

The resulting F estimates indicate that the level of exploitation has been relatively constant over the time period for which data are available.

10.1.3.2. Yearclass strength

K. Brander (personal communication) estimated the size of the 1975 yearclass as 2 500 000 1-year-old fish. To account for the number of fish of this yearclass taken in 1976 by the Irish fishery using this value, it would be necessary to increase the terminal F value on this age group to 1.4, which appears to be unrealistic. Therefore, this yearclass, as well as those of 1976 and 1977, was assumed to be of average strength (6 866 000 1 year olds).

10.1.3.3. Catch predictions (Tables 5.1 and 6.7)

Weights at age were adjusted by -5.88% to simulate the actual catch in 1976. Subsequent to the 1976 meeting, Brander pointed out that the 1974 yearclass was rather better than its presence in the 1975 catch as 1 year olds had indicated. Consequently, the catch in 1976 was higher than predicted, and as a result, the predicted catches in 1977 and 1978 continue to be better than previously estimated. A TAC of 8 600 tons was recommended for Division VIIa for 1978 (Table 5.1.). For Divisions VIIb-k, a precautionary TAC of 10 900 tons was recommended (Table 5.1).

10.2. Haddock

10.2.1. North Sea (Sub-Area IV)

10.2.1.1. Total international catch per age group

For the years 1959-71, the data presented in last year's report (ICES, CMI1976/F:9, Table 10) were used. The age compositions for these years are based solely on samples from Recommendation 4 fisheries by England, the Netherlands and Scotland, and therefore they probably underestimate the catches of younger age groups. From 1972, the data included the catches from the Recommendation 2 fisheries by Denmark and Norway, and, in addition, Dutch discards. However, as a result of new information on discarding rates, it was decided to revise the data base back to 1972.

In addition to revised Dutch data, new information on discarding by Scottish vessels in 1975 and 1976 was available, and it is believed that similar discarding by these vessels also occurred prior to 1975. Accordingly, estimates were made of Scottish discards from 1972 to 1974, based on the ratio Scottish discards at age in 1975 and 1976 to total numbers at age in international landings. No attempt was made to revise the data prior to 1972.

In the revised data, catches of age groups 1-3 are increased considerably by the inclusion of the discards.

The data for 1976 are provisional.

10.2.1.2. Mean weight at age (Table 6.2)

Data were available for the Danish and Norwegian Recommendation 2 fisheries, for the Scottish discards, and for the Scottish and Polish Recommendation 4 fisheries.

10.2.1.3. VPA (Table 2.2)

The catch at age was derived as described in 10.2.1.1. For natural mortality, a value of $M = 0.2$ was assumed for all age groups. A terminal F value of 1.1 was used for the oldest age group (age 10) in all years. This was derived by inspection of values of survival rates (e^{-Z}) for age groups 6-10 in successive years, under

the assumption that fishing mortality is relatively constant in successive years in the older fish.

With reference to F values in 1976, an attempt was made to estimate these by relating past F values to various measures of fishing effort, but no satisfactory relationship could be found (Figure 2). Therefore, for age groups older than 2, mean F values for the period 1970-72 were used and adjusted to give a smoothed exploitation pattern. For age groups 0 and 1, input F values were adjusted to produce population numbers which corresponded to those indicated by the International Young Herring Survey (Table 5.3). However, for age group 2, this procedure was not possible, since the catch had already exceeded the value indicated by the IYHS. For this age group, therefore, a value was obtained from the exploitation pattern by interpolation.

10.2.1.4. Yield per recruit (Table 5.4)

The effect of various F values on yield per recruit was investigated, using the 1976 assumed exploitation pattern and mean weights at age weighted by the numbers caught (Table 6.2). A reduction of about 70% of the present value ($F = 1.1$) is necessary to achieve the maximum yield per recruit at $F = 0.3$. The gain in yield per recruit is 20%.

10.2.1.5. Catch predictions

Input values for the catch predictions are given in Tables 6.2 and 5.5. The starting point was the catch at age in 1976 for the following categories: Rec.2 landings; Rec.4 landings; and discards. Age compositions for the first two categories were adjusted by sums of products (numbers x mean weight) to agree with preliminary catch data for 1976. Sums of products were used to estimate the weight of discards.

The following predictions were made:

Run 1. Fishing mortality constant 1976-78
($F_{76} = F_{77} = F_{78}$)

Run 2. Here it was assumed that the TAC of 190 000 tons for 1977 previously recommended by the Group would be taken.

This would result in an increase in F_{77} of 25%. It has been assumed that F_{78} then returns to the 1976 level.

$$(F_{77} = 1.25 \times F_{76} ; F_{78} = F_{76})$$

Run 3. Here it was assumed that the 1977 landings were 190 000 tons and that the increased F value was maintained in 1978.

$$(F_{78} = F_{77} = 1.25 \times F_{76})$$

Run 4. As for Run 2, except that the TAC in 1978 achieves a reduction of 10% in F compared to 1976.

$$(F_{77} = 1.25 \times F_{76} ; F_{78} = 0.9 \times F_{76})$$

Run 5. F constant in 1977 and reduced by 10% in 1978.

$$(F_{77} = F_{76} ; F_{78} = 0.9 \times F_{76})$$

Run 6. F constant in 1977 and reduced by 20% in 1978.

$$(F_{77} = F_{76} ; F_{78} = 0.8 \times F_{76})$$

Recruitment estimates were obtained from the IYHS data or were assumed to be average for future yearclasses. The average was calculated from yearclasses 1958-72, but excluding the exceptional yearclasses of 1962 and 1967.

The results of the prediction runs are given in Table 6.2.

TACs given in Section 3 of the report were based on Run 1 (Option A) and Run 5 (Option B).

10.2.2. West of Scotland (Division VIa)

10.2.2.1. Total international catch per age group

The same data base was used as in last year's report (ICES, GM1976/F:9, Table 13). The 1975 age compositions were updated and a provisional 1976 age composition was produced.

10.2.2.2. Mean weight at age

These are shown in Table 6.5 and are the same as those used last year.

10.2.2.3. VPA (Tables 2.5, 3.5 and 4.5)

Terminal F values in 1976 were obtained from average values in the period 1970-73. The exploitation pattern obtained indicated that F decreased on the older age groups and a terminal value of 0.15 was therefore selected for age 8 for all years.

For age groups 1 and 2, F values were adjusted so that population numbers corresponded to estimates of recruitment based on North Sea data.

The relation between effort data and F values was examined, but no satisfactory relations were obtained.

10.2.2.4. Yield per recruit

The same calculations were made as for the North Sea using the weight data and exploitation pattern given in Table 6.5. The present value of F (0.64) is close to that corresponding to F_{\max} with reference to yield per recruit.

10.2.2.5. Catch predictions

The input data for these calculations are given in Table 6.5. The starting point was the catch at age in 1976. Recruitment of 1-group fish in 1977 was estimated from the correlation between recruitment in Division VIa and in Sub-Area IV. Recruitment in 1978 was assumed to be average, based on the yearclasses 1964-72 (excluding the 1967 yearclass).

The following prediction runs were made:

Run 1. No change in fishing mortality

$$(F_{76} = F_{77} = F_{78})$$

Run 2. Here it was assumed that the TAC of 10 000 tons recommended by the Group for 1977 would be taken. This implies a reduction in fishing mortality of 40%. It was then assumed that F returned to the 1976 level in 1978.

$$(F_{77} = 0.6 \times F_{76} ; F_{78} = F_{76})$$

Run 3. No change in fishing mortality in 1977, but the reduction by 10% in 1978.

$$(F_{77} = F_{76} ; F_{78} = 0.9 \times F_{76})$$

Run 4. F constant in 1977 and reduced by 20% in 1978.

$$F_{77} = F_{76} ; F_{78} = 0.8 \times F_{76})$$

The results are given in Table 6.5.

TACs given in Table 5.1 were based on Run 1 (Option A) and Run 3 (Option B), and a precautionary TAC of 2 000 tons was recommended for Division VIb (Table 5.1).

10.2.3. Sub-Area VII

For Sub-Area VII, a precautionary TAC of 8 200 tons was recommended (Table 5.1).

10.3. Whiting

10.3.1. North Sea (Sub-Area IV)

10.3.1.1. Total international catch per age group

Estimates of the number of whiting discarded by Dutch vessels in each age group have been included in the total age composition used for the VPA assessments in earlier reports of the North Sea Roundfish Working Group. However, Dutch discard age compositions have been reassessed by Daan (1976). These new data have been included in the age compositions used during the present meeting of the Working Group.

Scottish discard data were also available at the meeting for 1975 and 1976. Provisional estimates for these 2 years show an average discarding rate by weight of 55% of the landed quantities. This rate has been used each year from 1967 to 1974 to calculate the quantity discarded by Scottish vessels. These figures, together with the weight of the Dutch discarded fish, estimated from their length compositions by numbers and a length/weight relation ($W = 0.008 L^3$), have been used to raise the Dutch age composition to a total age composition for the Netherlands and Scotland. Separate age compositions for Scotland and the Netherlands were available for both 1975 and 1976.

10.3.1.2. Mean weight at age

The mean weights at age used for the Recommendation 4 fisheries are based on Scottish landings in the period 1964-73. For the industrial fisheries age groups 0, 1 and 2, values from Danish landings in 1975 were used. The same values were used for the discards. The values in Table 6.3 are based on the numbers in the 3 components.

10.3.1.3. VPA

Based on the relationship between total international effort and F in Figure 2, it was decided to use mean values of fishing mortality for age groups 0-4 in

1972-73 as F values for 1976 in the VPA. For older age groups, the mean for all ages in the two years (0.85) was used (Table 3.3).

It was noted that with these values, the VPA estimate of strength of the year-class 1974 was much higher than the IYHS estimate. However, even an F value of 1.2 for two year olds in 1976 was not sufficient to make the two estimates agree. No adjustments were made to the F value to try to take account of this factor therefore.

10.3.1.4. Catch predictions

The input data for the catch prediction (Table 6.3) was based on three components in the catches in 1976, i.e., the catch in Recommendation 4 fisheries, the catch in Recommendation 2 fisheries and discards from the Dutch and Scottish Recommendation 4 fisheries (Table 5.6).

The numbers of recruits as 0-group fish were all taken to be $2\,300 \times 10^6$.

Predicted catches of each yearclass were reduced by a fraction based on ratios of discarded/total number landed in each age group to arrive at predictions of landings.

Six prediction runs were made as follows:

Run 1. $F_{76} = F_{77} = F_{78}$ (no change in fishing mortality)

Run 2. $F_{76} = F_{77}$; $F_{78} = 0.9 \times F_{76}$

Run 3. $F_{76} = F_{77}$; $F_{78} = 0.8 \times F_{76}$

Runs 4 - 6. As Runs 1 - 3, but with a 25% reduction of F_{76} .

The results are given in Table 6.3.

TACs given in Section 3 of the report were based on Run 4 (Option A) and Run 5 (Option B).

10.3.2. Sub-Area VI

In most years the catches from Division VIb were small compared to the catches from Division VIa, and the two Divisions were treated together.

For the years 1965 to 1976, age composition data for the Scottish landings were available. For 1976, Irish data were also available.

For the VPA, the exploitation pattern in 1976 was adjusted so that it corresponds to the average at 1971-73 (Table 3.6). The same F values were used for the catch prediction (Table 6.6). The strengths of the yearclasses 1976 and 1977 were taken to be average for the years 1964-73. Three prediction runs were made:

Run 1. No change in fishing mortality.

$$(F_{76} = F_{77} = F_{78})$$

Run 2. A reduction of fishing mortality in 1978 by 10%.

$$(F_{77} = F_{76} ; F_{78} = 0.9 \times F_{76})$$

Run 3. A reduction of fishing mortality in 1978 by 20%.

$$(F_{77} = F_{76} ; F_{78} = 0.8 \times F_{76})$$

The results are given in Table 6.6.

TACs given in Section 3 of the report were based on Run 1 (Option A) and Run 2 (Option B).

10.3.3. Sub-Area VII

For Sub-Area VII, a precautionary TAC of 25 700 tons was recommended (Table 5.1).

Table 1.1 Nominal catch of Cod, Haddock and Whiting (metric tons)
by Div. IIIa and Sub-areas IV, VI and VII, 1966-1976 (Bulletin Statistique)

| Sub-Area ^{a)} | | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 [*]) |
|------------------------|------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------------------|
| Cod | IIIa | 15 706 | 17 010 | 16 649 | 13 243 | 14 238 | 19 052 | 21 667 | 22 942 | 27 452 | 32 284 | 33 481 |
| | IV | 220 033 | 249 803 | 285 314 | 199 258 | 224 745 | 320 564 | 347 055 | 234 466 | 211 247 | 187 692 | 208 884 |
| | VI | 18 655 | 25 214 | 25 022 | 24 272 | 13 557 | 10 760 | 17 266 | 12 746 | 14 827 | 13 406 | 13 352 |
| | VII | 22 580 | 23 162 | 20 270 | 21 509 | 15 102 | 22 134 | 18 767 | 19 239 | 17 350 | 20 206 | ... |
| Haddock | IIIa | 695 | 469 | 582 | 1 056 | 942 | 2 249 | 8 989 | 3 091 | 4 618 | 6 115 | 3 019 |
| | IV | 269 205 | 167 408 | 139 469 | 639 195 | 671 833 | 258 220 | 213 556 | 196 079 | 193 640 | 184 003 | 205 654 |
| | VI | 31 816 | 21 176 | 21 429 | 27 398 | 35 018 | 46 920 | 50 518 | 32 848 | 67 258 | 63 611 | 59 017 |
| | VII | 9 655 | 7 343 | 3 726 | 5 392 | 5 931 | 6 518 | 11 248 | 12 480 | 10 585 | 8 638 | ... |
| Whiting | IIIa | 20 306 | 30 157 | 29 497 | 16 544 | 13 130 | 13 989 | 14 652 | 22 547 | 28 842 | 14 690 | 17 127 |
| | IV | 157 573 | 91 245 | 144 920 | 215 829 | 181 506 | 113 044 | 109 532 | 141 191 | 188 663 | 153 409 | 190 686 |
| | VI | 18 787 | 19 709 | 14 474 | 12 550 | 12 499 | 16 032 | 15 394 | 16 709 | 17 058 | 20 053 | 20 520 |
| | VII | 25 873 | 33 123 | 29 691 | 26 821 | 15 710 | 17 836 | 20 845 | 26 655 | 28 203 | 32 433 | ... |

*) provisional figures.

a) see footnotes on following page

Footnotes to Table 1.1.

Cod in Division IIa

Landings of German, Dem.Rep. in 1966, 1969-72 included in Sub-Area IV.
Landings of Sweden in 1966-74 included in Sub-Area IV.
Landings of GermanyFed.Rep. for 1968-70 include miscellaneous products.

Haddock in Division IIIa

Landings of German, Dem.Rep. in 1966, 1969-72 included in Sub-Area IV.
Landings of Sweden in 1966, 1968-74 included in Sub-Area IV.

Whiting in Division IIIa

Landings of German, Dem.Rep. in 1966 included in Sub-Area IV.
Landings of Sweden in 1966-74 included in Sub-Area IV.

Cod in Sub-Area IV

German, Dem.Rep. landings in 1966, 1969-72 include Division IIIa.
Sweden: landings 1966-74 include Division IIIa.
GermanyFed.Rep. landings in 1968-70 include miscellaneous products.
French figures for 1971-75 revised (March 1977).
Norway landings revised for 1974-75.
For Netherlands - not included for 1967 - 3 369 tons and 1968 - 1 132 tons.

Haddock in Sub-Area IV

French landings for 1971-75 figures revised.
Landings for Germany, Dem.Rep. for 1966, 1969-72 include Division IIIa.
Landings for Sweden for 1966, 1968-74 include Division IIIa.
Netherlands: Not included for 1967 - 720 tons and for 1968 - 306 tons caught mostly in Division IVb, rest in Division IVc.
Norway landings revised for 1974-75.

Whiting in Sub-Area IV

Landings for Germany, Dem.Rep. in 1966 include Division IIIa.
Landings for Sweden for 1966-74 includes Sub-Area IV and Division IIIa.
France - figures for 1971-1975 revised (March 1977).
Netherlands: Not included for 1967 - 913 tons and for 1968 - 267 tons.
Norway landings revised for 1974-75.

Cod in Sub-Area VI

Landings for GermanyFed.Rep. include miscellaneous products.
Landings for France 1971-75 revised.

Haddock in Sub-Area VI

French landings for 1971-75 figures revised.

Whiting in Sub-Area VI

French data for 1971-75 revised.

Footnotes to Table 1.1 (Continued)

Cod in Sub-Area VII

Landings for France for 1971-75 revised.

Haddock in Sub-Area VII

French landings for 1971-1975 figures revised.

Whiting in Sub-Area VII

French figures for 1971-75 revised.

Table 1.2 COD Div. IIIa and the Div. of Sub-areas IV, VI and VII
Nominal catch by Divisions in metric tons 1966 - 1976

| Area ^{a)} \ Year | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 [*]) |
|---------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------------------|
| IIIa | 15 706 | 17 010 | 16 649 | 13 243 | 14 238 | 19 052 | 21 667 | 22 942 | 27 452 | 32 284 | 33 481 |
| IVa | 69 440 | 89 923 | 74 051 | 56 015 | 79 606 | 67 370 | 80 650 | 69 557 | 72 362 | 59 582 | 56 189 |
| IVb | 125 233 | 134 258 | 175 949 | 122 027 | 110 271 | 184 957 | 215 160 | 134 953 | 114 087 | 107 227 | 135 705 |
| IVc | 25 360 | 25 622 | 35 314 | 21 216 | 34 868 | 68 237 | 51 245 | 29 956 | 24 798 | 20 883 | 16 990 |
| VIa | 17 133 | 23 025 | 24 357 | 21 739 | 12 682 | 10 666 | 14 699 | 12 263 | 13 652 | 13 163 | 11 690 |
| VIIb | 1 522 | 2 189 | 665 | 2 533 | 875 | 94 | 2 567 | 483 | 1 175 | 243 | 1 662 |
| VIIa | 5 249 | 12 652 | 8 541 | 7 967 | 6 257 | 9 540 | 9 173 | 11 787 | 10 190 | 9 790 | 8 142 |
| VIIb, c | 206 | 1 479 | 2 259 | 4 418 | 2 049 | 1 302 | 735 | 1 009 | 405 | 692 | ... |
| VIIId, e | 1 064 | 3 300 | 4 113 | 3 856 | 2 553 | 5 432 | 3 544 | 2 077 | 3 436 | 5 082 | ... |
| VIIIf | 1 188 | 1 321 | 1 514 | 856 | 925 | 797 | 969 | 976 | 594 | 998 | ... |
| VIIg-k | 14 873 | 4 410 | 3 843 | 4 412 | 3 318 | 5 063 | 4 346 | 3 390 | 2 725 | 3 644 | ... |
| Total | 273 974 | 315 189 | 347 255 | 258 282 | 267 642 | 372 510 | 404 755 | 289 393 | 270 876 | 253 588 | |

*) provisional figures

a) see footnotes on following page

Footnotes to Table 1.2.

Division IIIa

German, Dem.Rep. figures for 1966 and 1969-72 }
Swedish figures for 1966-74 } included in Division IVa.

GermanyFed.Rep. figures for 1968-70 include miscellaneous products.

Danish figure for 1976 including industrial catch only and is lacking some landings in foreign ports.

Division IVa

Danish figures for 1966-73 included in Division IVb.

German, Dem.Rep. figures for 1966 and 1969-72 include Divisions IIIa and IVb,c.

Swedish figures for 1966-74 include Divisions IIIa and IVb.

GermanyFed.Rep. figures for 1968-70 include miscellaneous products.

French figures for 1971-75 revised (March 1977).

Danish figure for 1976 included in Division IVb.

French figures for 1966 and 1976 }
Norwegian figures for 1966-68 and 1976 } include Division IVb,c
USSR figures for 1966-73 }

Norwegian figures for 1969-72 include Division IVb.

Norwegian figures 1974-75 revised (March 1977).

Norwegian figures for 1971 and 1972 not including catches from Recommendation 2 fisheries (1971 = 1 314 tons; 1972 = 1 656 tons).

Division IVb

Danish figures for 1966-73 included in Division IVa.

French figures for 1971-75 revised (March 1977).

Faroe Islands figure for 1976 }
French figures for 1966 and 1976 }
German, Dem.Rep. figures for 1966, 1969-72 and 1976 } included in Division IVa
Norwegian figures for 1966-72 and 1976 }
Swedish figures for 1966-74 }
USSR figures for 1966-73 }

Swedish figure for 1976; }
UK (Eng. + Wales) figure for 1976 } included Division IVa,c - from Data Form 5.

Netherlands: Not included for 1967 - 3 369 tons and 1968 - 1 132 tons caught mostly in Division IVb, rest in Division IVc.

GermanyFed.Rep. figures for 1968-70 include miscellaneous products.

Swedish figures for 1975 include Division IVa,c.

Danish figure for 1976 includes Division IVa,c. From Recommendation 12 form industrial catch only in Division IVa 988 tons; Division IVb 2 529 tons.

Division IVc

French figure for 1966 }
German, Dem.Rep. figures for 1966, 1969-72 and 1976 } included in Division IVa.
Norwegian figures for 1966-69 }
USSR figures for 1966-73 }

Germany Fed.Rep. figures for 1968-70 include miscellaneous products.

/Cont'd.

Footnotes to Table 1.2 (Continued)

| | | |
|-----------------------------------|---|---------------------------|
| Danish figure for 1976 | } | included in Division IVb. |
| French figure for 1976 | | |
| Swedish figures for 1975 and 1976 | | |
| UK (Eng. + Wales) figure for 1976 | | |

French figures for 1971-75 revised.

Division VIa

Swedish figure for 1968 includes Division VIb.
Germany, Fed.Rep. figures for 1968-70 include miscellaneous products.
French figures for 1971-75 revised.

Division VIb

Swedish figure for 1968 included in VIa.
French figures for 1971-75 revised.

Division VIIa

French figures for 1971-75 revised.
French figure for 1966 included in Division VIIg-k.
French figure for 1971 includes Division VIIf.

Division VIIb,c

French figure for 1966 included in Division VIIg-k.
French figure for 1971-75 revised.

Division VIIId,e

French figures for 1971-75 revised.

Division VIIIf

French figures for 1971-75 revised.
French figure for 1966)
Polish figure for 1976) included in Division VIIg-k

French figure for 1971 included in Division VIIa.

Division VIIg-k

French figure for 1971-75 revised.
French figure for 1966 includes Divisions VIIa,b,c and f.
Polish figure for 1976 includes Division VIIf.

Table 1.3 HADDOCK Div. IIIa and the Div. of Sub-areas IV, VI and VII
Nominal catch by Divisions in metric tons 1966 -1976

| Year Area a) | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 [*]) |
|-----------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------------------|
| IIIa | 695 | 469 | 582 | 1 056 | 942 | 2 249 | 2 989 | 3 091 | 4 618 | 6 115 | 3 019 |
| IVa | 197 518 | 122 531 | 75 347 | 271 953 | 455 649 | 197 306 | 135 095 | 131 819 | 128 818 | 120 688 | 161 015 |
| IVb | 71 283 | 44 823 | 62 696 | 361 836 | 212 646 | 58 270 | 75 325 | 62 288 | 63 695 | 62 761 | 44 306 |
| IVc | 404 | 54 | 1 426 | 5 406 | 3 538 | 2 644 | 3 136 | 1 972 | 1 127 | 554 | 333 |
| VIa | 29 881 | 20 302 | 20 526 | 26 273 | 34 178 | 46 299 | 41 044 | 28 830 | 17 970 | 13 683 | 15 698 |
| VIb | 1 935 | 874 | 903 | 1 125 | 840 | 621 | 9 474 | 4 018 | 49 288 | 49 928 | 43 319 |
| VIIa | 270 | 2 614 | 611 | 807 | 624 | 1 343 | 1 318 | 2 364 | 697 | 276 | 220 |
| VIIb,c | 245 | 787 | 433 | 758 | 1 922 | 1 141 | 1 419 | 931 | 2 090 | 2 565 | ... |
| VIIId,e | 37 | 111 | 88 | 811 | 421 | 170 | 411 | 359 | 633 | 971 | ... |
| VIIIf | 137 | 66 | 47 | 50 | 77 | 152 | 766 | 1 804 | 594 | 928 | ... |
| VIIg-k | 8 966 | 3 765 | 2 547 | 2 966 | 2 887 | 3 712 | 7 334 | 7 022 | 6 571 | 3 898 | ... |
| Total | 311 371 | 196 396 | 165 206 | 673 041 | 713 724 | 313 907 | 278 311 | 244 498 | 276 101 | 262 367 | |

^{*}) provisional figures

a) see footnotes on following page

Footnotes to Table 1.3.

Division IIIa

German, Dem.Rep. figures for 1966 and 1969-72 } included in Division IVa.
Swedish figures for 1966 and 1968-74 }

Danish figure includes industrial catch only and is lacking some landings in foreign ports.

Division IVa

Swedish figure for 1975 } included in Division IVb.
Danish figures for 1966-73 }

French figures for 1971-75 revised.

French figure for 1966 }
German, Dem.Rep. figure for 1976 } include Division IVb,c
Norwegian figures for 1966-69 and 1976 }
USSR figures for 1966-73 }

German, Dem.Rep. figures for 1966 and 1969-72 include Division IIIa and IVb,c.

Norwegian figures for 1969-72 } include Division IVb.
Swedish figure for 1967 }

Swedish figures for 1966 and 1968-74 include Divisions IIIa and IVb.

Danish figure for 1976 includes Divisions IVb,c - data from Data Form 5. From
Recommendation 12 industrial catch only Division IVa - 26 074; IVb - 12 785;
IVc - 12.

French figure for 1976 includes Division IVb,c from Data Form 5, up to November 1976.

Spanish }
Swedish } figure for 1976 includes Division IVb,c from Data Form 5.
UK (Eng. + Wales) }

Faroe Islands figure for 1976 includes Division IVb - the split in areas calculated from logbook returns up to 12 November 1976.

Norwegian figures for 1971 and 1972 not including catches from the Recommendation 2 fisheries (1971 = 4 512 tons; 1972 = 5 685 tons).

Norwegian landings revised for 1974-75.

Division IVb

Danish figures for 1966-73 include Division IVa.

Faroe Islands figure for 1976 }
French figures for 1966 and 1976 }
German, Dem.Rep. figures for 1966, 1969-72 and 1976 } included in Division IVa.
Norwegian figures for 1966-72 and 1976 }
Spanish figure for 1976 }
Swedish figures for 1966-74 and 1976 }
UK (Eng. + Wales) figure for 1976 }
USSR figures for 1966-73 }

Netherlands: Not included for 1967 - 720 tons and for 1968 - 306 tons caught mostly in Division IVb, rest in Division IVc.

Swedish figure for 1975 includes Division IVa,c.

Danish figure for 1976 includes Division IVb,c - data from Data Form 5. From
Recommendation 12 industrial catch only Division IVa - 26 074 tons; IVb - 12 785;
IVc - 12.

French figures for 1971-75 revised.

/Cont'd.

Footnotes to Table 1.3 (Continued)

Division IVc

| | | |
|---|---|---------------------------|
| French figures for 1966 and 1976 | } | included in Division IVa. |
| German, Dem.Rep. figures for 1966, 1969-72 and 1976 | | |
| Norwegian figures for 1966-68 and 1976 | | |
| Spanish figure for 1976 | | |
| Swedish figure for 1976 | | |
| UK (Eng. + Wales) figure for 1976 | | |
| USSR figures for 1966-73 | } | |

Netherlands: Not included for 1967 - 720 tons and for 1968 - 306 tons caught mostly in Division IVb, rest in Division IVc.

Swedish figure for 1975 included in Division IVb.

Danish figure for 1976 includes Division IVb,c - data from Data Form 5. From Recommendation 12 industrial catch only Division IVa - 26 074; Division IVb - 12 785; Division IVc - 12.

French figures for 1971-75 revised.

Division VIa

French figures for 1971-75 revised.

Division VIb

French figures for 1971-75 revised.

Division VIIa

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

French figure for 1971 includes Division VIIf.

Division VIIb,c

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

Divisions VIId,e

French figures for 1971-75 revised.

Division VIIf

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

French figure for 1971 included in Division VIIa.

Division VIIg-k

French figures for 1971-75 revised.

French figure for 1966 includes Divisions VIIa,b,c and f.

Table 1.4 WHITING Div. IIIa and the Div. of Sub-areas IV, VI and VII
Nominal catch by Divisions in metric tons 1966 - 1976

| Area \ Year a) | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 [*]) |
|-------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------------------|
| IIIa | 20 306 | 30 157 | 29 497 | 16 544 | 13 130 | 13 989 | 14 562 | 22 547 | 28 842 | 19 690 | 17 127 |
| IVa | 78 438 | 43 218 | 51 701 | 49 839 | 32 185 | 23 451 | 32 932 | 31 104 | 81 771 | 88 687 | 98 475 |
| IVb | 72 704 | 41 449 | 76 928 | 157 568 | 126 024 | 70 728 | 66 789 | 96 678 | 87 842 | 41 930 | 67 010 |
| IVc | 6 431 | 6 578 | 16 291 | 8 422 | 23 297 | 18 865 | 9 811 | 13 409 | 19 050 | 22 792 | 25 210 |
| VIa | 15 542 | 17 586 | 13 989 | 12 181 | 11 222 | 15 225 | 15 313 | 16 646 | 17 057 | 20 041 | 20 498 |
| VIIb | 3 245 | 2 123 | 485 | 369 | 1 277 | 807 | 81 | 63 | 1 | 12 | 22 |
| VIIa | 5 803 | 18 902 | 12 875 | 9 724 | 4 804 | 8 383 | 7 680 | 10 337 | 9 819 | 9 832 | 9 421 |
| VIIb,c | 276 | 2 246 | 3 249 | 3 595 | 1 507 | 287 | 1 056 | 1 091 | 1 243 | 1 829 | ... |
| VIIId,e | 1 307 | 5 554 | 6 640 | 5 066 | 4 825 | 3 592 | 3 676 | 5 647 | 8 572 | 11 400 | ... |
| VIIIf | 724 | 1 573 | 1 740 | 2 856 | 2 036 | 315 | 728 | 1 366 | 1 468 | 1 752 | ... |
| VIIg-k | 17 763 | 4 848 | 5 187 | 5 580 | 2 538 | 5 259 | 7 705 | 8 214 | 7 101 | 7 620 | ... |
| Total | 222 539 | 174 234 | 218 582 | 271 744 | 223 845 | 160 901 | 160 333 | 207 102 | 262 766 | 225 585 | 241 786 |

^{*}) provisional figures

a) see footnotes on following page

Footnotes to Table 1.4.

Division IIIa

German, Dem.Rep. figure for 1966 }
Swedish figures for 1966-74 } included in Division IVa.

Danish figure for 1976 includes industrial catch only and is lacking some landings in foreign ports.

Division IVa

French figures for 1971-75 revised.

German, Dem.Rep. figure for 1966 includes Divisions IIIa and IVb,c.

Danish figures for 1966-73 }
Swedish figure for 1975 } included in Division IVb.

French figures for 1966 and 1969 }
German, Dem.Rep. figure for 1976 }
Norwegian figures for 1966-69 and 1976 } include Divisions IVb,c.
USSR figures for 1966-73 }

Swedish figures for 1966-73 include Divisions IIIa and IVb.

Norwegian figures for 1969-72 include Division IVb.

Danish figure for 1976 includes industrial catch only and is lacking some landings in foreign ports.

Faroe Islands figure for 1976 includes Division IVb. The split on areas calculated from logbook returns up to 12 November 1976.

French figure for 1976 included in Division IVc.

Spanish }
Swedish } figure for 1976 includes Divisions IVb,c - from Data Form 5.
UK (Eng. + Wales) }

Norwegian figures for 1971 and 1972 not including catches from the Recommendation 2 fisheries (1971 = 1 605 tons; 1972 = 2 023 tons).

Norwegian landings revised for 1974-75.

Division IVb

French figures for 1971-75 revised.

Faroe Islands figure for 1976 }
French figures for 1966 and 1969 }
German, Dem.Rep. figures for 1966 and 1976 }
Norwegian figures for 1966-72 and 1976 } included in Division IVa.
Spanish figure for 1976 }
Swedish figures for 1966-74 and 1976 }
UK (Eng. + Wales) figure for 1976 }
USSR figures for 1966-73 }

Danish figures for 1966-73 include Division IVa.

Netherlands: Not included for 1967 - 913 tons and for 1968 - 257 tons caught mostly in Division IVb, rest in Division IVc.

Swedish figure for 1975 includes Divisions IVa,c.

Danish figure for 1976 includes industrial catch only and is lacking some landings in foreign ports.

French figure for 1976 included in Division IVa.

/Cont'd.....

Footnotes to Table 1.4 (Continued)

Division IVc

French figures for 1971-75 revised.

| | | |
|---|---|---------------------------|
| French figures for 1966 and 1969 | } | included in Division IVa. |
| German Dem.Rep. figures for 1966 and 1976 | | |
| Norwegian figures for 1966-69 and 1976 | | |
| Spanish figure for 1976 | | |
| Swedish figure for 1976 | | |
| UK (Eng. + Wales) figure for 1976 | | |
| USSR figures for 1966-73 | } | |

Netherlands: Not included for 1967 - 913 tons and for 1968 - 257 tons caught mostly in Division IVb, rest in Division IVc.

Swedish figure for 1975 included in Division IVb.

Danish figure for 1976 includes industrial catch only and is lacking some landings in foreign ports.

French figure for 1976 includes Division IVa and Division IVb - from Data Form 5, up to November 1976.

Division VIa

French figures for 1971-75 revised.

Division VIb

French figures for 1971-75 revised.

Faroe Islands: The split on areas calculated from logbook returns up to 12 November 1976.

Division VIIa

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

French figure for 1971 includes Division VIIf.

Division VIIb,c

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

Division VIId,e

French figures for 1971-75 revised.

Division VIIf

French figures for 1971-75 revised.

French figure for 1966 included in Division VIIg-k.

French figure for 1971 included in Division VIIa.

Division VIIg-k

French figure for 1971-75 revised.

French figure for 1966 includes Divisions VIIa,b,c and f.

Table 1.5 Nominal catches of Cod (metric tons) from Recommendation 2 fisheries in Sub-area IV ^{*)}
(data taken from NEAFC reports)

| Country | 1970 | | 1971 | | 1972 | | 1973 | | 1974 | | 1975 ^{*)} | |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|---------------------|-------------|----------------------|-------------|
| | legal-sized | under-sized | legal-sized | under-sized | legal-sized | under-sized | legal-sized | under-sized | legal-sized | under-sized | legal-sized | under-sized |
| Denmark | 20 | | 8 332 | 3 601 | 8 213 | 1 076 | 5 189 | 1 313 | 4 215 | 2 498 | 248 ^{d)} | |
| Germany, Fed. Rep. of | 1 202 | 3 140 | 4 125 | 970 | 555 | 54 | ? | ? | - | 1 | | |
| Netherlands | 6 731 | - | 8 199 | - | ? | - | 5 931 | 67 | 7 679 | - | 1 347 ^{d)} | |
| Norway (IVa) | 579 | 463 | 730 | 584 | 920 | 736 | 480 | 659 | 733 | 368 | | |
| Poland ^{a)} | 211 | 8 | 181 | 6 | 189 | 23 | ? | ? | 210 | 11 | 150 | 7 |
| Sweden ^{a)} | - | - | - | - | - | - | - | - | 8 260 | ... | 6 247 | ... |
| U.K. (England) | - | - | - | - | - | - | - | - | 6 | - | - | - |
| U.K. (Scotland) ^{a)} | - | - | - | - | - | - | - | - | 736 | - | 1 983 | - |
| Faroe Islands ^{a)} | - | - | - | - | - | - | - | - | 415 | 1 | - | - |
| Total ^{b)} | 8 723 | 3 611 | 21 567 | 5 161 | 9 877 | 1 889 | 11 600 | 2 039 | 22 254 | 2 879 | 10 032 | |
| Nominal catches of <u>Haddock</u> (metric tons) from Recommendation 2 fisheries in Sub-area IV (data taken from NEAFC reports) | | | | | | | | | | | | |
| Denmark | 145 | 201 | 13 657 | 7 651 | 9 088 | 11 568 | 771 | 3 155 | 9 364 | 27 785 | 8 621 ^{d)} | |
| Germany, Fed. Rep. of | 5 913 | 67 | 927 | 32 | 263 | - | ? | ? | - | - | | |
| Netherland | 7 003 | - | 6 341 | - | ? | - | 2 088 | 1 | 2 237 | - | 77 | |
| Norway (IVa) | 1 726 | 1 852 | 2 176 | 2 336 | 2 742 | 2 943 | 1 055 | 4 102 | 3 379 | 2 356 | | |
| Poland ^{a)} | - | - | 12 | 1 | 38 | 7 | ? | ? | 115 | 7 | 2 978 | 3 |
| Sweden ^{a)} | - | - | - | - | - | - | - | - | 2 954 | ... | 652 | ... |
| U.K. (Scotland) ^{a)} | - | - | - | - | - | - | - | - | 553 | 1 842 | 1 582 | - |
| Faroe Islands ^{a)} | - | - | - | - | - | - | - | - | 20 | 186 | - | - |
| Total ^{b)} | 14 642 | 1 919 | 23 113 | 10 020 | 12 131 | 14 518 | 3 914 | 7 258 | 18 622 | 32 176 | 13 940 | |
| Nominal catches of <u>Whiting</u> (metric tons) from Recommendation 2 fisheries in Sub-area IV (data taken from NEAFC reports) | | | | | | | | | | | | |
| Denmark | 102 | 141 | 20 462 | 34 493 | 29 446 | 20 035 | 57 194 | 16 081 | 84 448 | 24 578 | 366 ^{d)} | |
| Germany, Fed. Rep. of | 3 666 | 262 | 2 923 | 119 | 926 | 184 | ? | ? | 1 081 ^{c)} | - | | |
| Netherlands | 2 020 | - | 1 193 | - | ? | - | 2 153 | 14 | 4 281 | - | 13 794 ^{d)} | |
| Norway (IVa) | 789 | 484 | 995 | 610 | 1 254 | 769 | 1 322 | 166 | 4 710 | 312 | | |
| Poland ^{a)} | - | - | 2 | - | - | - | ? | - | 74 | 4 | 45 | 2 |
| Sweden ^{a)} | - | - | - | - | - | - | - | - | 860 | ... | 845 | ... |
| U.K. (Scotland) ^{a)} | - | - | - | - | - | - | - | - | 1 442 | 559 | 3 441 | 948 |
| Faroe Islands ^{a)} | - | - | - | - | - | - | - | - | 31 | 494 | - | - |
| Total ^{b)} | 6 475 | 746 | 25 575 | 35 222 | 31 626 | 20 988 | 60 669 | 16 261 | 95 847 | 25 947 | 19 538 | |

^{a)} IIIa inclusive ^{b)} total of available data only ^{c)} excluded from totals ^{d)} preliminary estimates ^{*)} for Belgium in IVc catches of legal-sized fish were 13 tons for cod and 94 tons for whiting; for German Democratic Republic catches of legal sized fish were 37 tons for cod, 27 tons for haddock and 3 tons for whiting.

Table 1.6 Nominal catch of COD for Div. IVa-IVc
by country in metric tons, 1971 - 1976
(Bulletin Statistique)

| Country | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 [*]) |
|--------------------------------|---------------------|---------------------|---------|---------------------|---------------------|---------------------|
| Belgium | 19 334 | 21 133 | 11 741 | 10 253 | 7 566 | 5 957 |
| Denmark | 68 179 | 72 520 | 47 950 | 54 207 | 46 344 | 53 971 |
| Faroe Islands | 123 | 284 | 803 | 416 | 732 | 400 |
| France ^{a)} | 24 769 | 24 038 | 13 247 | 7 275 | 8 667 | 5 646 |
| German Dem. Rep. ^{b)} | 18 | 122 | 343 | 132 | 223 | 69 |
| Germany, Fed. Rep. | 46 647 | 49 431 | 21 410 | 17 089 | 16 457 | 21 094 |
| Iceland | 1 | - | + | + | - | |
| Netherlands | 46 614 | 47 634 | 25 758 | 24 029 | 23 263 | 22 924 |
| Norway | 7 732 ^{c)} | 4 377 ^{c)} | 4 831 | 2 437 ^{d)} | 2 767 ^{d)} | 2 913 |
| Poland | 178 | 189 | 1 551 | 4 750 | 2 991 | 2 961 |
| Spain | - | 91 | 90 | 80 | 63 | |
| Sweden ^{b)} | 9 062 | 8 769 | 8 074 | 8 168 | 900 | 721 |
| U.K. (Engl. & Wales) | 55 525 | 62 503 | 47 327 | 39 857 | 33 615 | 46 475 |
| U.K. (Scotland) | 37 229 | 55 190 | 48 844 | 39 887 | 37 308 | 39 566 |
| U.S.S.R. | 5 153 | 774 | 2 497 | 2 667 | 6 796 | 6 187 |
| Total | 320 564 | 347 055 | 234 466 | 211 247 | 187 692 | 208 884 |

^{*}) provisional figures

a) French figures for 1971-1975 revised

b) GDR figures for 1971-1972 and Swedish figures for 1971-1974 include IIIa

c) Norwegian figures for 1971-1972 do not include cod caught in Recommendation 2 fisheries (1971 = 1 314; 1972 = 1 656 tons)

d) Norwegian figures for 1974-1975 revised for Div. IVa (March 1977).

Table 1.7 Nominal catch of HADDOCK for Div. IVa-IVc
by country in metric tons, 1971 - 1976
(Bulletin Statistique)

| Country | 1971 | 1972 | 1973 | 1974 | 1975 | 1976* |
|-------------------------------|---------------------|---------------------|---------|---------------------|----------------------|---------|
| Belgium | 971 | 1 601 | 2 385 | 1 137 | 2 209 | 1 605 |
| Denmark | 31 043 | 34 858 | 13 118 | 44 342 | 32 930 | 46 821 |
| Faroe Islands | - | 5 | 1 198 | 435 | 267 | 14 |
| France ^{e)} | 8 738 | 7 814 | 4 695 | 4 020 | 4 646 | 3 680 |
| German Dem.Rep. ^{a)} | 3 | 90 | 22 | 8 | 44 | 20 |
| Germany, Fed.Rep. | 3 045 | 4 020 | 4 587 | 3 478 | 2 396 | 3 204 |
| Iceland | 1 | - | - | - | - | - |
| Netherlands | 6 914 | 5 188 | 3 185 | 3 035 | 1 901 | 1 754 |
| Norway | 1 063 ^{b)} | 1 146 ^{b)} | 5 611 | 6 165 ^{d)} | 10 171 ^{d)} | 3 098 |
| Poland | - | 38 | 2 553 | 3 001 | 1 485 | 1 155 |
| Spain ^{c)} | - | - | 101 | 210 | - | 222 |
| Sweden ^{a)} | 5 857 | 5 305 | 4 550 | 3 098 | 2 083 | 3 091 |
| U.K. (Engl. & Wales) | 16 648 | 20 827 | 16 586 | 10 798 | 11 499 | 17 238 |
| U.K. (Scotland) | 121 539 | 96 197 | 88 132 | 71 679 | 64 686 | 80 062 |
| U.S.S.R. | 62 398 | 36 467 | 49 356 | 42 234 | 49 686 | 43 690 |
| Total | 258 220 | 213 556 | 196 079 | 193 640 | 184 003 | 205 654 |

*) provisional figures

a) German Democratic figures for 1971-72 and Swedish figures for 1971-1974 include IIIa

b) Norwegian figures for 1971 and 1972 do not include haddock caught in Recommendation 2 fisheries. (1971 = 4 512 tons; 1972 = 5 685 tons)

c) Spain reported 90 tons caught in 1975

d) Norwegian figures for 1974 and 1975 revised for Div. IVa (March 1977)

e) French figures for 1971-1975 revised.

Table 1.8 Nominal catch of WHITING for Div. IVa-IVc
by country in metric tons, 1971 - 1976

| Country | 1971 | 1972 | 1973 | 1974 | 1975 | 1976* |
|----------------------|------------------|------------------|---------|---------------------|----------------------|---------|
| Belgium | 2 108 | 2 745 | 3 387 | 3 156 | 3 279 | 2 186 |
| Denmark | 55 618 | 50 109 | 73 928 | 109 654 | 61 941 | 116 862 |
| Faroe Islands | - | - | 1 453 | 1 126 | 764 | 6 |
| France ^{a)} | 16 668 | 19 822 | 20 353 | 19 825 | 20 079 | 12 958 |
| German Dem. Rep. | - | - | 5 | - | 3 | 18 |
| Germany, Fed. Rep. | 233 | 264 | 403 | 454 | 446 | 359 |
| Iceland | - | - | - | - | - | - |
| Netherlands | 6 322 | 7 613 | 8 811 | 12 057 | 14 078 | 12 370 |
| Norway | 25 ^{b)} | 28 ^{b)} | 1 527 | 5 068 ^{c)} | 13 298 ^{c)} | 6 072 |
| Poland | - | - | 7 | 1 002 | 888 | 509 |
| Spain | - | 107 | 119 | 110 | 65 | 73 |
| Sweden ^{d)} | 616 | 596 | 2 328 | 2 440 | 255 | 847 |
| U.K. (Engl. & Wales) | 4 158 | 3 789 | 4 592 | 5 519 | 5 246 | 5 680 |
| U.K. (Scotland) | 26 755 | 23 846 | 20 756 | 25 274 | 27 969 | 26 038 |
| U.S.S.R. | 541 | 613 | 3 522 | 2 978 | 5 098 | 6 708 |
| Total | 113 044 | 109 532 | 141 191 | 188 663 | 153 409 | 190 686 |

*) provisional figures

a) French figures for 1971-1975 revised.

b) Norwegian figures for 1971 and 1972 do not include whiting caught in Recommendation 2 fisheries. (1971 = 1 605 tons; 1972 = 2 023 tons).

c) Norwegian figures for 1974 and 1975 revised for Div. IVa (March 1977).

d) Swedish figures for 1971-1974 include IIIa.

Table 1.9 U.K. (England and Wales) Fishing Effort data for different areas

| Area | | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|-----------|-------|-------|-------|-------|-------|-------|-------|
| North Sea | Hours | 819.5 | 855.1 | 884.9 | 852.9 | 781.3 | 694.5 | 725.8 |
| | Av. Tons | 56 | 54 | 60 | 56 | 58 | 52 | 59 |
| | Ton-Hours | 4 589 | 4 618 | 5 309 | 4 776 | 4 532 | 3 611 | 4 282 |
| W. of Scotland | Hours | 49.2 | 33.3 | 33.6 | 32.4 | 31.1 | 35.8 | 40.6 |
| | Av. Tons | 254 | 242 | 445 | 392 | 351 | 307 | 310 |
| | Ton-Hours | 1 250 | 806 | 1 495 | 1 270 | 1 092 | 1 099 | 1 259 |
| Irish Sea | Hours | 128.0 | 151.4 | 147.9 | 159.3 | 119.7 | 142.8 | 133.5 |
| | Av. Tons | 43 | 41 | 39 | 43 | 41 | 40 | 42 |
| | Ton-Hours | 550 | 621 | 577 | 585 | 491 | 571 | 561 |
| Bristol Channel | Hours | 44.1 | 47.4 | 38.4 | 37.0 | 32.2 | 34.3 | 27.4 |
| | Av. Tons | 56 | 49 | 52 | 57 | 62 | 41 | 45 |
| | Ton-Hours | 247 | 232 | 200 | 211 | 200 | 141 | 123 |

Note: HOURS are in thousands, TON-HOURS in 10 thousands

Table 1.10 U.K. (Scotland) Fishing Effort (000's hours fishing) for different Areas

| Sub-area | Gear | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 |
|----------|-------------|------|------|------|------|------|------|------|------|------|
| IV | Trawl | 206 | 203 | 112 | 110 | 149 | 177 | 176 | 179 | 150 |
| | Light trawl | 24 | 41 | 54 | 67 | 98 | 109 | 146 | 117 | 160 |
| | Seine | 499 | 537 | 479 | 411 | 399 | 379 | 405 | 350 | 342 |
| | Total | 729 | 781 | 645 | 588 | 646 | 665 | 727 | 646 | 654 |
| VI | Trawl | 54 | 50 | 43 | 41 | 42 | 56 | 55 | 44 | 37 |
| | Light trawl | 83 | 66 | 105 | 115 | 129 | 142 | 91 | 86 | 129 |
| | Seine | 159 | 150 | 140 | 96 | 99 | 71 | 60 | 56 | 56 |
| | Total | 296 | 266 | 288 | 252 | 270 | 269 | 206 | 186 | 222 |

Provisional figures for 1976 show that 1976 effort in Sub-area IV was about 90% of that in 1975. In Sub-area VI the 1976 effort was about the same as in 1975

Table 1.11 International Effort Data from the North Sea

| Year | Cod | | Haddock | | Whiting | |
|------|-----------------------------|------------------------------|-----------------------------|------------------------------|-----------------------------|------------------------------|
| | International Effort | | International Effort | | International Effort | |
| | English Units 100 hrs | Scottish Units 100 hrs | English Units 100 hrs | Scottish Units 100 hrs | English Units 100 hrs | Scottish Units 100 hrs |
| 1959 | | | 97 198 | | | |
| 1960 | | | 109 798 | | | |
| 1961 | | | 104 522 | | | |
| 1962 | | | 84 568 | | | |
| 1963 | 36 999 | | 75 241 | | | |
| 1964 | 43 587 | | 94 615 | | | |
| 1965 | 43 034 | | 133 697 | | 186 530 | |
| 1966 | 41 213 | | 196 357 | | 331 869 | |
| 1967 | 46 120 | 42 914 | 180 873 | 15 793 | 230 406 | 20 169 |
| 1968 | 42 318 | 48 291 | 144 842 | 16 680 | 424 087 | 37 371 |
| 1969 | 38 282 | 38 822 | 385 785 | 58 867 | 809 264 | 67 876 |
| 1970 | 47 883 | 44 934 | 282 342 | 34 974 | 437 740 | 50 629 |
| 1971 | 49 367 | 55 624 | 132 631 | 13 725 | 232 477 | 27 294 |
| 1972 | 49 135 | 41 818 | 90 736 | 14 763 | 255 806 | 30 545 |
| 1973 | 42 254 | 34 898 | 100 829 | 16 175 | 262 243 | 49 454 |
| 1974 | 41 418 | 34 220 | 139 957 | 17 433 | 266 971 | 48 202 |
| 1975 | 38 522 | 32 585 | 105 188 | 17 555 | 185 561 | 32 675 |
| 1976 | 32 620 | | 87 202 | | 266 068 | |

Table 2.1. Cod.
Sub-Area IV. Catch in numbers ('000 fish) by year and by age.

| Age | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|---------|---------|--------|--------|--------|---------|
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 274 | 317 |
| 1 | 18 622 | 47 311 | 40 500 | 75 633 | 65 388 | 9 941 | 5 109 | 47 304 | 61 347 | 6 317 | 33 809 | 15 715 | 35 086 | 12 182 |
| 2 | 37 798 | 23 681 | 68 149 | 65 705 | 81 282 | 79 589 | 23 009 | 27 373 | 149 128 | 195 922 | 30 551 | 53 537 | 54 771 | 105 109 |
| 3 | 6 192 | 15 976 | 14 441 | 26 341 | 26 741 | 36 676 | 31 590 | 16 392 | 14 385 | 43 709 | 52 648 | 11 799 | 17 597 | 22 510 |
| 4 | 3 069 | 3 439 | 6 715 | 5 896 | 9 265 | 11 078 | 14 959 | 12 179 | 5 952 | 5 095 | 13 163 | 5 180 | 4 078 | 9 805 |
| 5 | 2 360 | 1 513 | 1 783 | 2 513 | 2 698 | 5 623 | 5 190 | 6 867 | 6 028 | 2 406 | 1 905 | 4 397 | 6 401 | 1 550 |
| 6 | 1 404 | 1 652 | 873 | 1 065 | 1 750 | 1 275 | 2 842 | 1 963 | 2 394 | 2 802 | 1 038 | 974 | 1 662 | 2 374 |
| 7 | 67 | 433 | 510 | 409 | 655 | 623 | 688 | 1 051 | 760 | 1 449 | 988 | 472 | 378 | 737 |
| 8 | 485 | 99 | 275 | 362 | 304 | 314 | 379 | 207 | 394 | 545 | 486 | 373 | 144 | 114 |
| 9 | 4 | 390 | 14 | 77 | 148 | 154 | 170 | 221 | 182 | 339 | 38 | 310 | 175 | 63 |
| 10 | 5 | 1 | 81 | 64 | 36 | 103 | 54 | 136 | 82 | 102 | 41 | 65 | 73 | 44 |
| 11 | 1 | 1 | 1 | 25 | 2 | 21 | 110 | 46 | 53 | 5 | 64 | 35 | 29 | 22 |
| 12+ | 2 | 2 | 4 | 8 | 6 | 9 | 17 | 24 | 26 | 11 | 73 | 27 | 20 | 16 |

Table 2.2. Haddock.
Sub-Area IV. Catch in numbers ('000 fish) by year and by age.

| Age | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 161 900 | 41 800 | 387 000 | 62 303 | 144 891 |
| 1 | 199 600 | 2 000 | 73 000 | 11 700 | 102 000 | 376 000 | 96 500 | 6 300 | 48 300 | 292 350 | 33 000 | 361 800 | 653 682 | 91 166 |
| 2 | 118 100 | 430 500 | 3 700 | 6 700 | 25 400 | 190 100 | 17 285 | 119 100 | 22 700 | 288 860 | 344 760 | 101 530 | 262 828 | 552 299 |
| 3 | 13 500 | 146 400 | 460 800 | 17 700 | 3 300 | 26 700 | 181 800 | 15 011 | 37 500 | 27 400 | 240 900 | 252 100 | 38 852 | 220 267 |
| 4 | 12 200 | 17 100 | 33 200 | 410 500 | 6 700 | 2 300 | 26 800 | 34 600 | 372 300 | 20 100 | 9 000 | 49 000 | 96 419 | 12 792 |
| 5 | 6 500 | 9 500 | 6 800 | 24 600 | 194 800 | 2 200 | 5 200 | 600 | 11 400 | 147 500 | 6 100 | 2 600 | 10 489 | 35 363 |
| 6 | 500 | 4 300 | 3 800 | 4 300 | 4 800 | 66 100 | 2 300 | 500 | 700 | 3 300 | 1 600 | 1 100 | 652 | 5 578 |
| 7 | 400 | 300 | 700 | 400 | 500 | 600 | 42 500 | 200 | 200 | 100 | 40 | 9 600 | 434 | 242 |
| 8 | 900 | 500 | 300 | 80 | 300 | 70 | 5 100 | 2 600 | 1 800 | 400 | 1 | 200 | 2 888 | 83 |
| 9 | 10 | 60 | 20 | 5 | 40 | 10 | 10 | 20 | 900 | 8 | 4 | 20 | 108 | 800 |
| 10 | 10 | 10 | 10 | 1 | 3 | 6 | 1 | 3 | 200 | 100 | 20 | 40 | 42 | 87 |

Table 2.3. Whiting.

Sub-Area IV. Catch in numbers ('000 fish) by year and by age.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 0 | 35 800 | 26 864 | 225 344 | 149 071 | 114 392 | 105 852 | 969 531 | 478 565 | 201 785 | 492 277 | 181 773 | 298 317 |
| 1 | 80 050 | 267 347 | 187 736 | 425 514 | 513 060 | 486 258 | 208 832 | 642 039 | 638 510 | 873 497 | 602 340 | 274 648 |
| 2 | 53 023 | 187 031 | 163 927 | 317 412 | 790 117 | 172 353 | 90 844 | 235 436 | 446 112 | 745 235 | 273 809 | 663 190 |
| 3 | 222 525 | 72 901 | 123 885 | 101 396 | 133 868 | 401 920 | 22 821 | 41 610 | 108 925 | 190 795 | 255 145 | 124 264 |
| 4 | 61 271 | 188 881 | 28 061 | 48 832 | 30 646 | 34 378 | 115 699 | 6 816 | 18 653 | 32 495 | 60 267 | 69 475 |
| 5 | 8 466 | 33 896 | 59 486 | 10 730 | 11 183 | 10 568 | 13 065 | 51 901 | 5 905 | 5 000 | 11 565 | 14 106 |
| 6 | 3 873 | 3 226 | 7 714 | 23 612 | 3 807 | 4 051 | 2 241 | 5 971 | 18 094 | 1 779 | 2 487 | 3 402 |
| 7 | 928 | 1 540 | 923 | 2 190 | 7 248 | 504 | 801 | 843 | 2 638 | 5 409 | 781 | 954 |
| 8 | 141 | 451 | 150 | 138 | 3 499 | 1 673 | 662 | 575 | 635 | 578 | 1 651 | 154 |

Table 2.4 COD Division VIa
Catch in number ('000 fish) by year and by age

| Year Age | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-------------|-------|-------|-------|-------|------|------|-------|-------|-------|-------|-------|
| 1 | . 81 | 101 | 222 | 84 | 92 | 335 | 220 | 153 | 727 | 1 260 | 548 |
| 2 | 1 119 | 1 004 | 859 | 986 | 272 | 884 | 2 264 | 504 | 1 841 | 2 043 | 4 252 |
| 3 | 452 | 1 427 | 1 862 | 970 | 944 | 523 | 1 068 | 1 271 | 752 | 1 217 | 1 542 |
| 4 | 459 | 141 | 1 296 | 1 519 | 457 | 709 | 483 | 518 | 874 | 506 | 688 |
| 5 | 280 | 140 | 112 | 624 | 356 | 220 | 405 | 145 | 235 | 269 | 256 |
| 6 | 23 | 104 | 121 | 104 | 133 | 185 | 91 | 161 | 53 | 60 | 169 |
| 7 | 16 | 21 | 72 | 84 | 24 | 68 | 72 | 42 | 52 | 11 | 65 |
| 8+ | 2 | 12 | 18 | 53 | 39 | 36 | 47 | 47 | 22 | 19 | 15 |

Table 2.5 HADDOCK Division VIa
Catch in number ('000 fish) by year and by age

| Year Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-------|--------|
| 1 | 5 | 278 | 516 | 9 311 | 0 | 230 | 2 448 | 590 | 1 208 | 1 970 | 4 861 | 915 |
| 2 | 1 654 | 359 | 11 419 | 7 387 | 48 921 | 164 | 2 844 | 22 221 | 6 520 | 3 425 | 9 519 | 16 056 |
| 3 | 84 419 | 1 164 | 1 239 | 3 234 | 5 928 | 71 520 | 6 627 | 2 225 | 15 648 | 9 411 | 2 773 | 12 325 |
| 4 | 4 697 | 47 424 | 238 | 418 | 1 386 | 3 795 | 91 387 | 2 897 | 263 | 6 131 | 3 427 | 1 403 |
| 5 | 206 | 1 606 | 18 775 | 586 | 350 | 211 | 590 | 56 846 | 1 147 | 97 | 1 980 | 1 488 |
| 6 | 169 | 76 | 252 | 11 729 | 576 | 92 | 86 | 612 | 31 836 | 447 | 106 | 911 |
| 7 | 139 | 30 | 20 | 655 | 3 386 | 98 | 6 | 37 | 139 | 11 488 | 122 | 29 |
| 8 | 23 | 102 | 28 | 36 | 150 | 453 | 97 | 57 | 114 | 189 | 3 770 | 15 |

Table 2.6. Whiting.

Sub-Area VI. Catch in numbers ('000 fish) by year and by age.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 4 | 54 | 1 |
| 1 | 2 239 | 1 126 | 4 261 | 7 037 | 684 | 697 | 2 640 | 11 064 | 13 009 | 7 577 | 17 551 | 11 918 |
| 2 | 4 857 | 12 935 | 25 182 | 18 154 | 25 631 | 2 676 | 7 712 | 9 657 | 27 463 | 42 873 | 18 712 | 45 387 |
| 3 | 41 177 | 2 454 | 10 755 | 9 729 | 9 753 | 30 312 | 3 936 | 3 447 | 6 758 | 12 215 | 39 477 | 14 329 |
| 4 | 5 299 | 28 248 | 857 | 3 583 | 2 794 | 4 514 | 30 759 | 1 168 | 1 831 | 2 035 | 3 243 | 15 730 |
| 5 | 784 | 1 767 | 16 762 | 267 | 1 276 | 818 | 1 394 | 12 800 | 469 | 505 | 307 | 1 413 |
| 6 | 68 | 213 | 803 | 4 772 | 109 | 210 | 249 | 712 | 5 293 | 68 | 60 | 104 |
| 7 | 185 | 36 | 84 | 269 | 1 708 | 14 | 47 | 58 | 273 | 1 387 | 6 | 18 |
| 8 | 12 | 17 | 23 | 31 | 155 | 392 | 78 | 64 | 33 | 64 | 194 | 1 |

Table 2.7. Cod.

Division VIIa. Catch in numbers ('000 fish) by year and by age.

| Age | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 1 | 364 | 882 | 905 | 2 762 | 777 | 2 258 | 462 | 936 | 1 817 |
| 2 | 1 563 | 1 481 | 1 710 | 2 200 | 3 241 | 1 064 | 4 284 | 759 | 2 881 |
| 3 | 1 003 | 1 050 | 344 | 824 | 832 | 1 792 | 561 | 1 632 | 479 |
| 4 | 456 | 269 | 211 | 179 | 247 | 437 | 392 | 276 | 351 |
| 5 | 177 | 186 | 229 | 76 | 61 | 172 | 60 | 152 | 39 |
| 6 | 28 | 76 | 44 | 49 | 39 | 63 | 43 | 33 | 54 |
| 7+ | 2 | 37 | 18 | 19 | 13 | 30 | 9 | 16 | 15 |

Table 3.1. Cod.
Sub-Area IV. Fishing mortality by year and by age.

| Age | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|--|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 1 | 0.22 | 0.25 | 0.22 | 0.31 | 0.29 | 0.13 | 0.07 | 0.15 | 0.16 | 0.08 | 0.21 | 0.10 | 0.11 | 0.12 |
| 2 | 0.63 | 0.48 | 0.69 | 0.68 | 0.63 | 0.69 | 0.47 | 0.61 | 0.97 | 1.09 | 0.72 | 0.61 | 0.58 | 0.52 |
| 3 | 0.40 | 0.60 | 0.61 | 0.63 | 0.66 | 0.66 | 0.67 | 0.74 | 0.76 | 0.89 | 1.03 | 0.69 | 0.41 | 0.50 |
| 4 | 0.44 | 0.41 | 0.56 | 0.54 | 0.48 | 0.64 | 0.63 | 0.59 | 0.67 | 0.68 | 0.75 | 0.25 | 0.54 | 0.43 |
| 5 | 0.42 | 0.41 | 0.38 | 0.42 | 0.52 | 0.60 | 0.72 | 0.68 | 0.67 | 0.65 | 0.59 | 0.60 | 0.55 | 0.41 |
| 6 | 0.74 | 0.59 | 0.44 | 0.42 | 0.58 | 0.49 | 0.71 | 0.66 | 0.53 | 0.78 | 0.65 | 0.70 | 0.48 | 0.41 |
| 7 | 0.21 | 0.53 | 0.36 | 0.38 | 0.49 | 0.42 | 0.55 | 0.63 | 0.58 | 0.73 | 0.71 | 0.71 | 0.65 | 0.41 |
| 8 | 0.56 | 0.53 | 0.78 | 0.47 | 0.54 | 0.46 | 0.49 | 0.31 | 0.51 | 1.17 | 0.59 | 0.65 | 0.49 | 0.41 |
| 9 | 0.18 | 1.28 | 0.13 | 0.52 | 0.36 | 0.59 | 0.49 | 0.59 | 0.50 | 1.19 | 0.21 | 0.97 | 0.74 | 0.41 |
| 10 | 0.46 | 0.06 | 1.08 | 1.46 | 0.49 | 0.46 | 0.42 | 0.96 | 0.46 | 0.59 | 0.42 | 0.67 | 0.65 | 0.41 |
| 11 | 0.28 | 0.15 | 0.08 | 1.31 | 0.14 | 0.59 | 1.38 | 0.77 | 1.42 | 0.04 | 0.93 | 0.76 | 0.74 | 0.41 |
| 12+ | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.55 | 0.41 |
| Mean $F \geq 2$ | (Sum of F's weighted by stocks in numbers) | | | | | | | | | | | | | |
| | 0.57 | 0.51 | 0.65 | 0.64 | 0.62 | 0.67 | 0.60 | 0.64 | 0.91 | 1.03 | 0.86 | 0.58 | 0.53 | 0.51 |

Table 3.2. Haddock.
Sub-Area IV. Fishing mortalities by year and by age.

| Age | 1959 | 1960 | 1961 | 1962 | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| 0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.43 | 0.03 | 0.16 | 0.12 | 0.26 |
| 1 | 0.10 | 0.16 | 0.24 | 0.13 | 0.07 | 0.03 | 1.16 | 0.09 | 0.16 | 0.07 | 0.32 | 0.07 | 0.06 | 0.24 | 0.14 | 0.35 | 0.43 | 0.27 |
| 2 | 0.56 | 0.64 | 0.56 | 0.31 | 0.33 | 0.22 | 0.08 | 0.29 | 0.29 | 0.49 | 0.50 | 0.84 | 0.35 | 0.54 | 0.49 | 0.84 | 0.47 | 0.80 |
| 3 | 0.46 | 0.74 | 0.94 | 0.63 | 0.29 | 0.89 | 0.38 | 0.61 | 0.22 | 0.57 | 1.31 | 1.14 | 0.70 | 0.97 | 1.29 | 0.83 | 0.96 | 0.93 |
| 4 | 0.98 | 0.72 | 0.76 | 0.78 | 0.51 | 0.74 | 0.51 | 0.68 | 0.50 | 0.24 | 2.55 | 1.00 | 1.03 | 1.09 | 1.05 | 1.08 | 0.91 | 1.04 |
| 5 | 1.29 | 0.90 | 0.67 | 0.56 | 0.70 | 0.97 | 0.75 | 0.90 | 0.83 | 0.30 | 1.35 | 0.40 | 1.18 | 1.95 | 1.32 | 1.08 | 0.71 | 1.10 |
| 6 | 1.03 | 1.11 | 0.84 | 0.45 | 0.48 | 1.61 | 1.60 | 1.91 | 0.43 | 0.77 | 0.58 | 0.42 | 1.20 | 1.59 | 0.09 | 0.93 | 0.91 | 1.10 |
| 7 | 1.56 | 1.03 | 1.09 | 1.34 | 0.50 | 0.59 | 1.61 | 0.73 | 1.72 | 0.09 | 2.28 | 0.09 | 0.29 | 0.52 | 0.06 | 1.03 | 1.35 | 1.10 |
| 8 | 1.52 | 1.77 | 2.04 | 2.25 | 2.33 | 2.98 | 3.17 | 0.83 | 3.06 | 1.54 | 2.45 | 1.09 | 3.50 | 1.65 | 0.01 | 0.48 | 1.09 | 1.10 |
| 9 | 0.97 | 1.31 | 2.20 | 0.97 | 0.44 | 1.45 | 2.46 | 0.65 | 1.53 | 1.86 | 1.04 | 0.05 | 1.77 | 0.20 | 0.05 | 0.23 | 0.52 | 1.10 |
| 10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 |
| Mean $F \geq 2$ | (Sum of F's weighted by stocks in numbers) | | | | | | | | | | | | | | | | | |
| | 0.89 | 0.69 | 0.73 | 0.53 | 0.35 | 0.30 | 0.38 | 0.68 | 0.68 | 0.54 | 0.57 | 1.11 | 0.94 | 0.84 | 0.70 | 0.86 | 0.58 | 0.85 |

Table 3.3. Whiting.

Sub-Area IV. Fishing mortalities by year and by age*)

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|--|------|------|------|------|------|------|------|------|------|------|------|
| 0 | 0.04 | 0.02 | 0.07 | 0.15 | 0.84 | 0.76 | 0.40 | 0.17 | 0.10 | 0.18 | 0.18 | 0.14 |
| 1 | 0.14 | 0.47 | 0.24 | 0.20 | 1.05 | 1.13 | 0.32 | 0.50 | 0.36 | 0.84 | 0.34 | 0.45 |
| 2 | 0.28 | 0.55 | 0.60 | 0.79 | 0.67 | 1.40 | 0.66 | 0.74 | 0.80 | 0.92 | 0.71 | 0.77 |
| 3 | 0.46 | 0.78 | 0.88 | 0.97 | 0.97 | 0.91 | 0.69 | 0.73 | 0.96 | 1.00 | 1.00 | 0.85 |
| 4 | 0.62 | 0.91 | 0.81 | 1.12 | 0.92 | 0.73 | 0.73 | 0.46 | 0.90 | 0.88 | 1.10 | 0.85 |
| 5 | 0.91 | 0.87 | 0.84 | 0.88 | 0.87 | 1.00 | 0.69 | 0.90 | 0.93 | 0.65 | 0.95 | 0.85 |
| 6 | 1.00 | 1.15 | 0.49 | 1.02 | 0.95 | 0.95 | 0.60 | 0.81 | 0.97 | 0.83 | 0.80 | 0.85 |
| 7 | 0.66 | 1.72 | 1.38 | 0.25 | 1.08 | 0.30 | 0.49 | 0.47 | 1.11 | 0.91 | 1.19 | 0.85 |
| 8 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 | 0.80 |
| Mean $F \geq 3$ | (Sum of F's weighted by stocks in numbers) | | | | | | | | 0.95 | 0.97 | 1.01 | 0.85 |
| | 0.50 | 0.88 | 0.84 | 0.98 | 0.96 | 0.89 | 0.72 | 0.78 | | | | |

Table 3.4. Cod.

Division VIa. Fishing mortalities by year and by age.

| Age | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|--|------|------|------|------|------|------|------|------|------|------|
| 1 | 0.01 | 0.02 | 0.04 | 0.03 | 0.02 | 0.04 | 0.06 | 0.02 | 0.08 | 0.06 | 0.09 |
| 2 | 0.19 | 0.16 | 0.28 | 0.25 | 0.14 | 0.26 | 0.44 | 0.18 | 0.41 | 0.35 | 0.27 |
| 3 | 0.50 | 0.39 | 0.48 | 0.59 | 0.40 | 0.43 | 0.57 | 0.48 | 0.44 | 0.53 | 0.48 |
| 4 | 0.56 | 0.28 | 0.73 | 0.94 | 0.62 | 0.59 | 0.93 | 0.61 | 0.72 | 0.60 | 0.66 |
| 5 | 0.59 | 0.33 | 0.38 | 0.99 | 0.60 | 0.69 | 0.83 | 0.82 | 0.63 | 0.51 | 0.70 |
| 6 | 0.34 | 0.46 | 0.54 | 0.75 | 0.59 | 0.73 | 0.70 | 0.98 | 0.84 | 0.32 | 0.70 |
| 7 | 0.66 | 0.59 | 0.66 | 0.91 | 0.38 | 0.70 | 0.72 | 0.84 | 1.06 | 0.41 | 0.70 |
| 8+ | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 |
| Mean $F \geq 2$ | (Sum of F's weighted by stocks in numbers) | | | | | | | | 0.49 | 0.42 | 0.34 |
| | 0.30 | 0.26 | 0.47 | 0.57 | 0.38 | 0.41 | 0.54 | 0.41 | | | |

*) Scottish and Dutch discards included.

Table 3.5. Haddock.

Division VIa. Fishing mortalities by year and by age.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|--|------|------|------|------|------|------|------|------|------|------|------|-------|
| 1 | 0.00 | 0.01 | 0.02 | 0.01 | 0.00 | 0.03 | 0.03 | 0.02 | 0.08 | 0.04 | 0.05 | 0.042 |
| 2 | 0.22 | 0.08 | 0.79 | 0.31 | 0.09 | 0.01 | 0.53 | 0.43 | 0.23 | 0.34 | 0.26 | 0.21 |
| 3 | 0.52 | 0.24 | 0.43 | 0.54 | 0.44 | 0.18 | 0.69 | 1.08 | 0.61 | 0.61 | 0.50 | 0.64 |
| 4 | 0.58 | 0.63 | 0.07 | 0.25 | 0.47 | 0.57 | 0.38 | 0.76 | 0.34 | 0.52 | 0.47 | 0.51 |
| 5 | 0.30 | 0.40 | 0.56 | 0.24 | 0.34 | 0.12 | 0.16 | 0.43 | 0.81 | 0.20 | 0.31 | 0.38 |
| 6 | 0.41 | 0.17 | 0.10 | 0.83 | 0.40 | 0.14 | 0.07 | 0.25 | 0.46 | 0.89 | 0.35 | 0.23 |
| 7 | 0.14 | 0.12 | 0.06 | 0.40 | 0.61 | 0.11 | 0.01 | 0.04 | 0.08 | 0.30 | 0.65 | 0.15 |
| 8 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 | 0.15 |
| Mean $F \geq 2$ (Sum of F's weighted by stocks in numbers) | | | | | | | | | | | | |
| | 0.51 | 0.58 | 0.57 | 0.51 | 0.11 | 0.19 | 0.39 | 0.44 | 0.44 | 0.41 | 0.27 | 0.32 |

Table 3.6. Whiting.

Sub-Area VI. Fishing mortalities by year and by age.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|
| 0 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.01 |
| 1 | 0.05 | 0.02 | 0.08 | 0.04 | 0.05 | 0.04 | 0.10 | 0.17 | 0.08 | 0.13 | 0.14 | 0.10 |
| 2 | 0.63 | 0.47 | 0.86 | 0.57 | 0.19 | 0.31 | 0.67 | 0.59 | 0.77 | 0.41 | 0.51 | 0.67 |
| 3 | 0.44 | 0.77 | 0.93 | 1.03 | 0.69 | 0.36 | 1.01 | 0.74 | 1.13 | 1.00 | 0.83 | 0.96 |
| 4 | 0.77 | 0.63 | 0.69 | 0.97 | 1.01 | 0.83 | 0.77 | 1.00 | 1.22 | 1.46 | 0.82 | 1.00 |
| 5 | 0.94 | 0.65 | 1.00 | 0.48 | 1.23 | 0.98 | 0.67 | 0.88 | 1.78 | 1.60 | 0.94 | 1.11 |
| 6 | 0.47 | 0.74 | 0.70 | 0.90 | 0.37 | 0.67 | 0.96 | 0.89 | 1.23 | 2.06 | 0.87 | 1.04 |
| 7 | 1.69 | 0.50 | 0.75 | 0.54 | 1.02 | 0.07 | 0.31 | 0.62 | 1.12 | 1.49 | 1.36 | 0.71 |
| 8 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.90 | 0.90 | 0.90 | 0.90 | 0.90 |
| Mean $F \geq 5$ (Sum of F's weighted by stocks in numbers) | | | | | | | | | | | | |
| | 0.99 | 0.65 | 0.98 | 0.85 | 1.03 | 0.78 | 0.68 | 0.88 | 1.26 | 1.51 | 0.92 | 1.10 |

Table 3.7. Cod.

Division VIIa. Fishing mortalities by year and by age.

| Age | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----------------|--|------|------|------|------|------|------|------|------|
| 1 | 0.12 | 0.19 | 0.17 | 0.30 | 0.25 | 0.23 | 0.19 | 0.13 | 0.35 |
| 2 | 0.60 | 0.93 | 0.66 | 0.77 | 0.69 | 0.63 | 0.90 | 0.56 | 0.71 |
| 3 | 0.82 | 1.10 | 0.57 | 0.80 | 0.77 | 1.10 | 0.82 | 1.12 | 0.86 |
| 4 | 0.86 | 0.54 | 0.68 | 0.67 | 0.59 | 1.34 | 0.77 | 1.39 | 0.78 |
| 5 | 0.89 | 1.14 | 1.33 | 0.57 | 0.51 | 1.14 | 0.65 | 0.80 | 0.75 |
| 6 | 0.43 | 1.37 | 0.97 | 1.29 | 0.65 | 1.77 | 1.06 | 0.93 | 0.75 |
| 7+ | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 | 0.75 |
| Mean F \geq 2 | (Sum of F's weighted by stocks in numbers) | | | | | | | | |
| | 0.70 | 0.95 | 0.69 | 0.77 | 0.69 | 0.95 | 0.87 | 0.91 | 0.73 |

Table 4.1. Cod.

Sub-Area IV. Stock in numbers ('000 fish) at beginning of year.

| Age | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| 0 | 286 351 | 271 531 | 384 463 | 346 244 | 112 963 | 105 115 | 452 559 | 560 740 | 105 299 | 236 172 | 224 503 | 470 218 | 145 190 | 349 925 |
| 1 | 103 698 | 234 445 | 222 311 | 314 772 | 283 481 | 92 487 | 86 061 | 370 524 | 459 095 | 86 212 | 193 361 | 183 808 | 384 982 | 118 624 |
| 2 | 88 238 | 68 141 | 149 387 | 145 564 | 189 738 | 173 313 | 66 761 | 65 851 | 260 736 | 320 605 | 64 886 | 127 878 | 136 317 | 283 558 |
| 3 | 20 553 | 38 453 | 34 566 | 61 441 | 60 484 | 82 680 | 70 820 | 34 036 | 29 433 | 80 878 | 88 603 | 25 851 | 56 821 | 62 601 |
| 4 | 9 403 | 11 271 | 17 194 | 15 386 | 26 755 | 25 625 | 34 921 | 29 758 | 13 239 | 11 265 | 27 302 | 25 778 | 10 627 | 30 734 |
| 5 | 7 542 | 4 946 | 6 142 | 8 067 | 7 318 | 13 602 | 11 077 | 15 218 | 13 469 | 5 522 | 4 671 | 10 607 | 16 445 | 5 050 |
| 6 | 2 936 | 4 058 | 2 692 | 3 428 | 4 350 | 3 575 | 6 107 | 4 436 | 6 325 | 5 642 | 2 371 | 2 121 | 4 752 | 7 735 |
| 7 | 393 | 1 151 | 1 844 | 1 421 | 1 851 | 1 996 | 1 785 | 2 462 | 1 878 | 3 035 | 2 121 | 1 013 | 866 | 2 401 |
| 8 | 1 243 | 261 | 555 | 1 052 | 796 | 929 | 1 075 | 845 | 1 076 | 857 | 1 192 | 854 | 408 | 371 |
| 9 | 27 | 584 | 125 | 209 | 537 | 380 | 479 | 541 | 506 | 528 | 219 | 541 | 366 | 205 |
| 10 | 15 | 19 | 133 | 90 | 102 | 307 | 173 | 240 | 245 | 251 | 132 | 145 | 167 | 143 |
| 11 | 4 | 8 | 14 | 37 | 17 | 51 | 159 | 93 | 75 | 127 | 115 | 71 | 61 | 72 |
| 12 | 3 | 3 | 5 | 11 | 8 | 12 | 23 | 33 | 35 | 15 | 100 | 37 | 27 | 24 |

Table 4.2. Haddock.

Sub-Area IV. Stock in numbers ('000 fish) at beginning of year.

| Age | 1963 | 1964 | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 |
|-----|-----------|-----------|-----------|---------|-----------|-----------|-----------|-----------|
| 0 | 85 790 | 140 355 | 179 261 | 937 252 | 7 690 140 | 470 168 | 133 182 | 1 189 970 |
| 1 | 3 203 000 | 70 239 | 114 913 | 146 767 | 767 358 | 6 296 160 | 384 941 | 109 040 |
| 2 | 460 159 | 2 442 310 | 55 701 | 29 372 | 109 609 | 536 361 | 4 815 590 | 228 456 |
| 3 | 58 491 | 270 647 | 1 612 100 | 42 266 | 18 024 | 66 908 | 268 801 | 2 394 100 |
| 4 | 33 580 | 35 752 | 91 246 | 906 208 | 18 776 | 11 787 | 30 888 | 59 281 |
| 5 | 14 142 | 16 564 | 14 011 | 44 966 | 375 264 | 9 370 | 7 581 | 1 982 |
| 6 | 1 440 | 5 774 | 5 116 | 5 404 | 14 919 | 133 657 | 5 694 | 1 610 |
| 7 | 1 118 | 731 | 941 | 842 | 654 | 7 910 | 50 473 | 2 604 |
| 8 | 1 062 | 557 | 330 | 154 | 332 | 96 | 5 935 | 4 240 |
| 9 | 31 | 85 | 23 | 11 | 55 | 13 | 17 | 419 |
| 10 | 16 | 16 | 16 | 2 | 5 | 10 | 2 | 5 |

| Age | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|-----------|-----------|-----------|-----------|-----------|-----------|
| 0 | 1 844 700 | 511 223 | 1 680 470 | 2 929 840 | 585 488 | 695 238 |
| 1 | 974 268 | 1 510 310 | 273 334 | 1 338 110 | 2 050 070 | 423 193 |
| 2 | 83 590 | 754 073 | 973 501 | 194 047 | 770 628 | 1 092 160 |
| 3 | 80 925 | 48 052 | 358 777 | 488 119 | 68 415 | 395 346 |
| 4 | 628 902 | 32 771 | 14 978 | 80 615 | 174 973 | 21 463 |
| 5 | 17 787 | 184 176 | 8 994 | 4 271 | 22 507 | 57 449 |
| 6 | 1 085 | 4 461 | 21 523 | 1 969 | 1 189 | 9 062 |
| 7 | 870 | 268 | 745 | 16 178 | 633 | 393 |
| 8 | 1 951 | 532 | 130 | 574 | 4 718 | 135 |
| 9 | 1 164 | 48 | 84 | 105 | 291 | 1 300 |
| 10 | 325 | 162 | 32 | 65 | 68 | 141 |

Table 4.3. Whiting.

Sub-Area IV. Stock in numbers ('000 fish) at beginning of year.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 0 | 988 160 | 1 223 190 | 3 441 470 | 1 211 980 | 2 195 170 | 2 165 990 | 3 241 110 | 3 388 710 | 2 258 920 | 3 356 000 | 1 213 750 | 2 513 570 |
| 1 | 681 856 | 776 720 | 977 208 | 2 614 330 | 857 951 | 778 008 | 829 049 | 1 783 540 | 2 343 350 | 1 667 480 | 2 304 270 | 830 025 |
| 2 | 237 112 | 486 111 | 396 304 | 631 149 | 1 757 310 | 246 792 | 205 708 | 491 134 | 885 049 | 1 345 170 | 587 005 | 1 345 510 |
| 3 | 664 198 | 146 457 | 230 561 | 177 849 | 233 724 | 732 915 | 49 820 | 87 242 | 192 000 | 326 863 | 438 209 | 236 135 |
| 4 | 144 319 | 344 301 | 54 895 | 78 456 | 55 448 | 72 335 | 242 338 | 20 408 | 34 291 | 60 328 | 98 060 | 132 021 |
| 5 | 15 454 | 63 377 | 113 781 | 19 932 | 20 918 | 18 126 | 28 539 | 95 150 | 10 598 | 11 471 | 20 459 | 26 805 |
| 6 | 6 667 | 5 117 | 21 703 | 40 157 | 6 765 | 7 168 | 5 449 | 11 697 | 31 703 | 3 423 | 4 922 | 6 465 |
| 7 | 2 102 | 2 016 | 1 329 | 10 857 | 11 899 | 2 152 | 2 265 | 2 457 | 4 254 | 9 868 | 1 217 | 1 813 |
| 8 | 279 | 892 | 297 | 273 | 6 919 | 3 308 | 1 309 | 1 137 | 1 256 | 1 143 | 3 265 | 305 |

Table 4.4. Cod.

Division VIa. Stock in numbers ('000 fish) at beginning of year.

| Age | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|--------|
| 1 | 9 406 | 4 825 | 6 266 | 2 896 | 5 277 | 8 860 | 4 390 | 7 450 | 10 144 | 25 497 | 7 014 |
| 2 | 7 209 | 7 628 | 3 859 | 4 929 | 2 295 | 4 238 | 6 952 | 3 396 | 5 962 | 7 649 | 19 738 |
| 3 | 1 258 | 4 895 | 5 340 | 2 387 | 3 149 | 1 634 | 2 674 | 3 661 | 2 326 | 3 229 | 4 428 |
| 4 | 1 165 | 625 | 2 727 | 2 704 | 1 087 | 1 731 | 869 | 1 234 | 1 859 | 1 230 | 1 554 |
| 5 | 686 | 543 | 385 | 1 076 | 862 | 481 | 783 | 281 | 547 | 742 | 555 |
| 6 | 88 | 312 | 319 | 215 | 326 | 388 | 197 | 280 | 101 | 238 | 366 |
| 7 | 36 | 51 | 162 | 153 | 83 | 148 | 152 | 80 | 86 | 36 | 141 |
| 8 | 3 | 15 | 23 | 68 | 50 | 46 | 60 | 60 | 28 | 24 | 19 |

Table 4.5. Haddock.

Division VIa. Stock in numbers ('000 fish) at beginning of year.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|---------|--------|---------|---------|---------|---------|---------|--------|--------|---------|--------|
| 1 | 6 329 | 28 100 | 37 554 | 772 051 | 21 780 | 9 491 | 88 237 | 43 061 | 17 449 | 57 409 | 119 193 | 24 528 |
| 2 | 9 188 | 5 177 | 22 755 | 30 281 | 623 693 | 17 832 | 7 563 | 70 032 | 34 722 | 13 196 | 45 224 | 93 199 |
| 3 | 227 432 | 6 034 | 3 915 | 8 448 | 18 153 | 466 511 | 14 451 | 3 645 | 37 406 | 22 561 | 7 727 | 28 465 |
| 4 | 11 643 | 110 601 | 3 893 | 2 094 | 4 021 | 9 547 | 317 539 | 5 913 | 1 010 | 16 632 | 10 055 | 3 842 |
| 5 | 879 | 5 330 | 48 157 | 2 973 | 1 338 | 2 050 | 4 420 | 177 944 | 2 257 | 591 | 8 126 | 5 161 |
| 6 | 556 | 535 | 2 922 | 22 623 | 1 906 | 781 | 1 488 | 3 087 | 94 703 | 826 | 396 | 4 873 |
| 7 | 1 137 | 303 | 369 | 2 165 | 8 070 | 1 044 | 557 | 1 141 | 1 977 | 48 995 | 278 | 229 |
| 8 | 182 | 806 | 221 | 284 | 1 185 | 3 579 | 766 | 450 | 901 | 1 493 | 29 788 | 119 |

Table 4.6. Whiting.

Sub-Area VI. Stock in numbers ('000 fish) at beginning of year.

| Age | 1965 | 1966 | 1967 | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|---------|--------|---------|---------|---------|---------|--------|---------|---------|---------|---------|---------|
| 0 | 72 234 | 74 142 | 251 357 | 17 553 | 26 597 | 39 001 | 97 727 | 226 000 | 86 685 | 174 914 | 168 552 | 1 |
| 1 | 48 633 | 59 140 | 60 703 | 205 794 | 14 371 | 21 776 | 31 932 | 80 012 | 185 033 | 70 972 | 143 204 | 137 950 |
| 2 | 11 365 | 37 796 | 47 403 | 45 855 | 162 137 | 11 149 | 17 199 | 23 762 | 55 541 | 139 757 | 51 277 | 101 429 |
| 3 | 125 805 | 4 963 | 19 351 | 16 382 | 21 297 | 109 667 | 6 723 | 7 192 | 10 816 | 20 978 | 75 957 | 25 220 |
| 4 | 10 719 | 66 076 | 1 874 | 6 274 | 4 771 | 8 726 | 62 569 | 2 007 | 2 812 | 2 861 | 6 320 | 27 012 |
| 5 | 1 394 | 4 049 | 28 845 | 769 | 1 951 | 1 423 | 3 122 | 23 790 | 605 | 681 | 547 | 2 284 |
| 6 | 197 | 444 | 1 736 | 8 718 | 390 | 467 | 438 | 1 310 | 8 080 | 83 | 112 | 174 |
| 7 | 244 | 100 | 174 | 704 | 2 891 | 222 | 195 | 137 | 439 | 1 928 | 9 | 39 |
| 8 | 26 | 37 | 50 | 67 | 336 | 849 | 169 | 117 | 60 | 117 | 355 | 2 |

Table 4.7. Cod.

Division VIIa. Stock in numbers at beginning of year.

| Age | 1968 | 1969 | 1970 | 1971 | 1972 | 1973 | 1974 | 1975 | 1976 |
|-----|-------|-------|-------|--------|-------|--------|-------|-------|-------|
| 1 | 3 654 | 5 689 | 6 442 | 11 698 | 3 906 | 12 095 | 2 869 | 8 579 | 6 749 |
| 2 | 3 790 | 2 663 | 3 863 | 4 459 | 7 095 | 2 499 | 7 871 | 1 933 | 6 180 |
| 3 | 1 953 | 1 705 | 863 | 1 635 | 1 688 | 2 914 | 1 095 | 2 631 | 903 |
| 4 | 857 | 705 | 464 | 398 | 604 | 640 | 796 | 396 | 706 |
| 5 | 327 | 296 | 337 | 192 | 166 | 273 | 138 | 302 | 81 |
| 6 | 88 | 110 | 77 | 73 | 89 | 82 | 71 | 59 | 112 |
| 7 | 3 | 47 | 23 | 24 | 16 | 38 | 11 | 20 | 19 |

Table 5.1. Predictions of catches and landings in 1978 ('000 tons).

| Species | Area | A ($F_{78}=F_{76}$) | | B ($F_{78}=0.9 \times F_{76}$) | | C ($F_{78}=0.8 \times F_{76}$) | | All Options |
|---------|--------|--------------------------|----------|-------------------------------------|----------|-------------------------------------|----------|---|
| | | Catch | Landings | Catch | Landings | Catch | Landings | |
| Cod | IV | 240 | | 220 | | 200 | | 1.3 ¹⁾ 10.9 ¹⁾ |
| | VIa | 20.6 | | 19.1 | | 17.4 | | |
| | VIb | | | | | | | |
| | VIIa | 9.3 | | 8.6 | | 7.9 | | |
| | VIIb-k | | | | | | | |
| Haddock | IV | 126 | 112 | 118 | 106 | 109 | 97 | 2.0 ¹⁾ 8.2 ¹⁾ |
| | VIa | | 11.4 | | 10.4 | | 9.5 | |
| | VIb | | | | | | | |
| | VII | | | | | | | |
| Whiting | IV | 202 | 173 | 187 | 161 | 172 | 148 | 25.7 ¹⁾ |
| | VI | | 17.8 | | 16.6 | | 15.2 | |
| | VII | | - | | - | | - | |

1) Averages for period 1966-1975.

A. Effort in 1978 the same as in 1976 and 1977.

B. Effort in 1978 10% lower than in 1976 and 1977.

C. Effort in 1978 20% lower than in 1976 and 1977.

Table 5.2. Revised estimates of yearclass strength.

| Yearclass | Cod Div.IVa,b,c | | Haddock Div.IVa,b,c | | Whiting Div.IVa,b,c | |
|-----------|--------------------|----------------------------|------------------------|----------------------------|------------------------|----------------------------|
| | IYHS ^{a)} | VPA ^{b)} M=0.2 | IYHS ^{a)} | VPA ^{b)} M=0.2 | IYHS ^{a)} | VPA ^{b)} M=0.2 |
| 1958 | | | | 368 | | |
| 1959 | | | | 234 | | |
| 1960 | | | | 152 | | |
| 1961 | | | | 638 | | |
| 1962 | | 104 | | 3 203 | | |
| 1963 | | 234 | | 70 | | |
| 1964 | | 222 | | 115 | | 682 |
| 1965 | | 315 | | 147 | | 777 |
| 1966 | 33 | 283 | 151 | 767 | 803 | 977 |
| 1967 | 5.6 | 92 | 8 891 | 6 296 | 1 726 | 2 614 |
| 1968 | 5.9 | 86 | 425 | 385 | 18 | 858 |
| 1969 | 59 | 371 | 45 | 109 | 86 | 778 |
| 1970 | 125 | 549 | 2 114 | 974 | 296 | 829 |
| 1971 | 2.6 | 86 | 3 044 | 1 510 | 710 | 1 784 |
| 1972 | 38 | 193 | 461 | 273 | 4 272 | 2 343 |
| 1973 | 10 | 184 | 3 685 | 1 338 | 703 | 1 667 |
| 1974 | 77 | 385 | 1 663 | 2 050 | 1 292 | 2 304 |
| 1975 | 6.4 | | 312 | | 1 306 | |
| 1976 | (45) | | (375) | | (1 030) | |

a) Average number per hour fishing during the International Young Herring Surveys (cf. ICES, C.M.1976/F:20).

b) Millions of fish at age 1.

Figures in brackets are provisional.

Table 5.3. Predictive regressions of VPA estimates of yearclass size (y) on yearclass strength indices (x) from research surveys ($y = B_0 + B_1 x$).

| Data | n | B_0 | B_1 | r | p | Estimated size yearclass (1 year old) '000 000 | |
|--|---|--------|-------|------|------------|---|-------|
| | | | | | | 1975 | 1976 |
| Cod IVa,b,c(M=0.2) - IYHS (1966-1974) | 9 | 100.36 | 3.47 | 0.93 | $p < 0.01$ | 6.4 | 45 |
| Haddock IVa,b,c(M=0.2) - IYHS (1966-1973*) | 7 | 258.03 | 0.30 | 0.86 | $p < 0.01$ | 352 | 371 |
| Whiting IVa,b,c(M=0.2) - IYHS (1966-1974) | 9 | 956.44 | 0.56 | 0.71 | $p < 0.05$ | 1 306 | 1 030 |

*) Excluding 1967.

Table 5.4. Percentage change in F values needed to give MSY per recruit. Also percentage gains in yield per recruit for $F = F_{max}$.

| Species | Area | % Change in F values | % Gain in yield per recruit |
|---------|------|----------------------|-----------------------------|
| Cod | IV | -50 | +15 |
| | VIa | -45 | + 7 |
| | VIIa | -65 | +20 |
| Haddock | IV | -70 | +20 |
| | VIa | 0 | 0 |
| Whiting | IV | -70 | 0* |
| | VIa | -65 | 0* |

* Yield per recruit curve flat-topped.

Table 5.5. Haddock.

Sub-Area IV. Input data for catch prediction.
1976 catch in numbers ('000 fish).

| Age | Industrial landings | | Other landings | | Discards | | Total |
|--------------|---------------------|--------|----------------|--------|----------|--------|-----------|
| | Numbers | Weight | Numbers | Weight | Numbers | Weight | Numbers |
| 0 | 144 791 | 0.025 | 0 | 0.000 | 70 | 0.041 | 144 861 |
| 1 | 60 485 | 0.064 | 2 150 | 0.230 | 28 519 | 0.108 | 91 154 |
| 2 | 157 246 | 0.157 | 200 575 | 0.280 | 194 452 | 0.185 | 552 273 |
| 3 | 23 849 | 0.334 | 173 925 | 0.410 | 22 492 | 0.246 | 220 266 |
| 4 | 60 | 0.423 | 12 618 | 0.580 | 114 | 0.253 | 12 792 |
| 5 | 2 584 | 0.556 | 32 704 | 0.710 | 75 | 0.314 | 35 363 |
| 6 | 34 | 0.666 | 5 544 | 0.940 | 0 | 0.000 | 5 578 |
| 7 | 0 | 0.000 | 242 | 1.210 | 0 | 0.000 | 242 |
| 8 | 0 | 0.000 | 83 | 1.440 | 0 | 0.000 | 83 |
| 9 | 0 | 0.000 | 800 | 1.500 | 0 | 0.000 | 800 |
| 10 | 0 | 0.000 | 87 | 1.600 | 0 | 0.000 | 87 |
| Total | 389 050 | | 428 728 | | 245 722 | | 1 063 500 |
| Total weight | | 41 629 | 165 466 | | 44 642 | | |

Table 5.6. Whiting.

Sub-Area IV. Input data for catch prediction.
1976 catch in numbers ('000 fish).

| Age | Recommendation 4 Fisheries | | Recommendation 2 Fisheries | | Discards |
|-----|----------------------------|----------------------|----------------------------|----------------------|----------|
| | Catch '000 | Mean weight kg | Catch '000 | Mean weight kg | '000 |
| 0 | | | 293 317 | .020 | 5.000 |
| 1 | 7 950 | .187 | 245 162 | .063 | 21 536 |
| 2 | 113 425 | .228 | 433 514 | .195 | 116 251 |
| 3 | 57 773 | .269 | 54 917 | .269 | 11 574 |
| 4 | 56 787 | .322 | 6 759 | .322 | 5 929 |
| 5 | 13 423 | .380 | 272 | .380 | 411 |
| 6 | 3 341 | .468 | 42 | .468 | 19 |
| 7 | 939 | .620 | 13 | .620 | 2 |
| 8 | 154 | .765 | - | .765 | |

Table 5.7. North Sea Cod, Haddock and Whiting.
Total numbers ('000) at each length group landed quarterly by Norway in 1976a).

| <u>Cod</u> <u>Length group</u> | <u>Quarter</u> | | | | <u>Total</u> |
|-----------------------------------|----------------|--------------|--------------|---------------|---------------|
| | 1 | 2 | 3 | 4 | |
| 10-14 | | 24 | | | 24 |
| 15-19 | | 166 | 95 | 47 | 308 |
| 20-24 | 24 | 47 | 47 | 47 | 165 |
| 25-29 | 95 | 71 | 24 | 308 | 498 |
| 30-34 | 95 | 214 | 95 | 379 | 783 |
| 35-39 | 118 | 261 | 236 | 261 | 876 |
| 40-44 | 94 | 118 | 118 | 188 | 518 |
| 45-49 | 47 | 24 | 212 | 94 | 377 |
| 50-54 | | 24 | 71 | 70 | 165 |
| 55-59 | 47 | | 71 | | 118 |
| Total | 520 | 949 | 969 | 1 394 | 3 832 |
| <u>Haddock</u> | | | | | |
| 5- 9 | 5 | | 422 | 18 | 445 |
| 10-14 | 211 | 79 | 2 014 | 750 | 3 054 |
| 15-19 | 396 | 2 083 | 528 | 3 513 | 6 520 |
| 20-24 | 519 | 569 | 2 970 | 1 253 | 5 311 |
| 25-29 | 844 | 545 | 1 541 | 1 330 | 4 260 |
| 30-34 | 127 | 147 | 602 | 675 | 1 551 |
| 35-39 | 22 | 47 | 121 | 242 | 432 |
| 40-44 | 9 | 4 | 47 | 45 | 105 |
| 45-49 | 9 | | 19 | 5 | 33 |
| 50-54 | | | 6 | 1 | 7 |
| Total | 2 142 | 3 474 | 8 270 | 7 832 | 21 718 |
| <u>Whiting</u> | | | | | |
| 10-14 | 4 | | 5 | 17 | 26 |
| 15-19 | 23 | | 18 | 74 | 115 |
| 20-24 | 357 | 100 | 195 | 537 | 1 189 |
| 25-29 | 1 487 | 2 095 | 988 | 4 072 | 8 642 |
| 30-34 | 972 | 1 535 | 1 197 | 5 670 | 9 374 |
| 35-39 | 389 | 559 | 495 | 1 902 | 3 345 |
| 40-44 | 25 | 50 | 185 | 270 | 530 |
| 45-49 | | 33 | 58 | 56 | 147 |
| 50-54 | | | | 2 | 2 |
| 55-59 | | | 6 | | 6 |
| Total | 3 257 | 4 372 | 3 147 | 12 600 | 23 376 |

a) Measurements from Recommendation 2 fisheries only.

Table 6.1. Cod.
Sub-Area IV. Input data for catch prediction.

| Age | 1976 Catch '000 | F-values 1976-1977 | Mean weight kg* |
|-----|--------------------|-----------------------|--------------------|
| 1 | 12 182 | 0.115 | 0.54 |
| 2 | 105 109 | 0.52 | 0.92 |
| 3 | 22 510 | 0.50 | 2.02 |
| 4 | 9 805 | 0.43 | 3.82 |
| 5 | 1 550 | 0.41 | 5.75 |
| 6 | 2 374 | 0.41 | 7.64 |
| 7 | 737 | 0.41 | 9.11 |
| 8 | 114 | 0.41 | 10.37 |
| 9 | 63 | 0.41 | 11.24 |
| 10+ | 82 | 0.41 | 12 |

a) Assuming F_{76} is 25% below average for the period 1963-75.

| Year | Recruitment at age 1 '000 | Simulated catches '000 tons | | |
|------|---------------------------|-----------------------------|------------------------------|------------------------------|
| | | $F_{78} = F_{76}$ | $F_{78} = 0.9 \times F_{76}$ | $F_{78} = 0.8 \times F_{76}$ |
| 1977 | 256 000 | 221 | 221 | 221 |
| 1978 | 230 000 | 240 | 220 | 200 |

b) Assuming F_{76} is the same as the average for the period 1963-75.

| Year | Recruitment at age 1 '000 | Simulated catches '000 tons | | |
|------|---------------------------|-----------------------------|------------------------------|------------------------------|
| | | $F_{78} = F_{76}$ | $F_{78} = 0.9 \times F_{76}$ | $F_{78} = 0.8 \times F_{76}$ |
| 1977 | 256 000 | 195 | 195 | 195 |
| 1978 | 230 000 | 211 | 195 | 177 |

*) These values had to be adjusted by -6.26% to yield the actual catch in weight in 1976.

Table 6.2. Haddock.

Sub-Area IV. Input data for catch prediction.

| Input data | | | | Prediction runs | | | | | | | |
|------------|------------------------|------------------------|-----------------|-----------------|-----------------|------------------|------------------|--------------|---------|--------------|---------|
| Age | 1976 Catch '000s | Mean weight (kg) | F ₇₆ | Run No. | Ratio of F's | | | Predicted 77 | | Predicted 78 | |
| | | | | | F ₇₆ | :F ₇₇ | :F ₇₈ | Landings | Catches | Landings | Catches |
| 0 | 144 861 | .029 | .26 | 1 | 1 | :1 | :1 | 165 | (183) | 112 | (126) |
| 1 | 91 154 | .086 | .27 | | | | | | | | |
| 2 | 552 273 | .210 | .80 | 2 | 1 | :1.25 | :1 | 189 | (210) | 93 | (106) |
| 3 | 220 266 | .396 | .93 | | | | | | | | |
| 4 | 12 792 | .549 | 1.04 | 3 | 1 | :1.25 | :1.25 | 189 | (210) | 106 | (121) |
| 5 | 35 363 | .704 | 1.1 | | | | | | | | |
| 6 | 5 578 | .940 | 1.1 | 4 | 1 | :1.25 | :0.9 | 189 | (210) | 87 | (99) |
| 7 | 242 | 1.210 | 1.1 | | | | | | | | |
| 8 | 83 | 1.440 | 1.1 | 5 | 1 | :1 | :0.9 | 165 | (183) | 106 | (118) |
| 9 | 800 | 1.500 | 1.1 | | | | | | | | |
| 10 | 87 | 1.600 | 1.1 | 6 | 1 | :1 | :0.8 | 165 | (183) | 97 | (109) |

Nos. 1 063 500

Weight 251 737

M = 0.2

Recruitment at age 0

1977 648 000 000
1978 648 000 000

Table 6.3. Whiting.

Sub-Area IV. Input data for catch prediction.

| Age | 1976 Catch landings + discards '000 | Mean weight kg | F values M = 0.2 |
|-----|---|-------------------|---------------------|
| 0 | 298 317 | 0.020 | 0.14 |
| 1 | 274 648 | 0.066 | 0.45 |
| 2 | 663 190 | 0.200 | 0.77 |
| 3 | 124 264 | 0.269 | 0.85 |
| 4 | 69 475 | 0.322 | 0.85 |
| 5 | 14 106 | 0.380 | 0.85 |
| 6 | 3 402 | 0.468 | 0.85 |
| 7 | 954 | 0.620 | 0.85 |
| 8 | 154 | 0.765 | 0.80 |

Recruitment at age 0 in '000

| | |
|------|-----------|
| 1976 | 2 300 000 |
| 1977 | 2 300 000 |
| 1978 | 2 300 000 |

Simulated catches ('000 tons) with F values in column 4

| Year | Change in F | Catch | Landings (= catch - discards) |
|------|------------------------------|-------|-------------------------------|
| 1977 | $F_{77} = F_{76}$ | 190 | 165 |
| 1978 | $F_{78} = F_{76}$ | 202 | 173 |
| | $F_{78} = 0.9 \times F_{76}$ | 187 | 161 |
| | $F_{78} = 0.8 \times F_{76}$ | 172 | 148 |

Simulated catches ('000 tons) with F values in column 4 reduced by 25%.

| Year | Change in F | Catch | Landings (= catch - discards) |
|------|------------------------------|-------|-------------------------------|
| 1977 | $F_{77} = F_{76}$ | 221 | 191 |
| 1978 | $F_{78} = F_{76}$ | 248 | 214 |
| | $F_{78} = 0.9 \times F_{76}$ | 229 | 198 |
| | $F_{78} = 0.8 \times F_{76}$ | 208 | 179 |

Table 6.4. Cod.

Division VIa. Input data for catch prediction.

| Age | 1976 Catch '000 | F values 1976-1977 | Mean weight kg* |
|-----|--------------------|-----------------------|--------------------|
| 1 | 548 | 0.09 | 0.58 |
| 2 | 4 252 | 0.27 | 1.22 |
| 3 | 1 542 | 0.48 | 2.66 |
| 4 | 688 | 0.66 | 4.25 |
| 5 | 256 | 0.70 | 5.13 |
| 6 | 169 | 0.70 | 6.41 |
| 7 | 65 | 0.70 | 8.38 |
| 8+ | 15 | 0.70 | 9.00 |

| Year | Recruitment at age 1 '000 | Simulated catches '000 tons | | |
|------|------------------------------|-----------------------------|------------------------------|------------------------------|
| | | $F_{78} = F_{76}$ | $F_{78} = 0.9 \times F_{76}$ | $F_{78} = 0.8 \times F_{76}$ |
| 1977 | 6 613 | 21.0 | 21.0 | 21.0 |
| 1978 | 6 613 | 20.6 | 19.1 | 17.4 |

Table 6.5. Haddock.

Division VIa. Input data for catch prediction.

| Age | 1976 Catch '000 | F values 1976-1977 | Mean weight kg |
|-----|--------------------|-----------------------|-------------------|
| 1 | 922 | .042 | .230 |
| 2 | 16 187 | .21 | .28 |
| 3 | 12 425 | .64 | .41 |
| 4 | 1 414 | .51 | .58 |
| 5 | 1 500 | .38 | .71 |
| 6 | 918 | .23 | .94 |
| 7 | 29 | .15 | 1.21 |
| 8 | 15 | .15 | 1.44 |
| 9 | 3 831 | .15 | 1.50 |

| Year | Recruitment at age 1 '000 | Simulated catches '000 tons | | |
|------|------------------------------|-----------------------------|-----------|--------------------------------|
| 1977 | 25 900 | Run 1 | 1977 16.3 | $(F_{77} = F_{76})$ |
| 1978 | 31 500 | | 1978 11.4 | $(F_{78} = F_{76})$ |
| | | Run 2 | 1977 10.8 | $(F_{77} = 0.6 \times F_{76})$ |
| | | | 1978 13.6 | $(F_{78} = F_{76})$ |
| | | Run 3 | 1977 16.3 | $(F_{77} = F_{76})$ |
| | | | 1978 10.4 | $(F_{78} = 0.9 \times F_{76})$ |
| | | Run 4 | 1977 16.3 | $(F_{77} = F_{76})$ |
| | | | 1978 9.5 | $(F_{78} = 0.8 \times F_{76})$ |

*) These values had to be adjusted by +2.85% to yield the actual catch in weight in 1976.

Table 6.6. Whiting.
Sub-Area VI. Input data for catch prediction.

| Age | 1976 Catch '000 | F values M = 0.2 | Mean weight kg |
|-----|--------------------|---------------------|-------------------|
| 1 | 11 918 | 0.10 | 0.213 |
| 2 | 45 387 | 0.67 | 0.241 |
| 3 | 14 329 | 0.96 | 0.267 |
| 4 | 15 730 | 1.00 | 0.310 |
| 5 | 1 413 | 1.11 | 0.377 |
| 6 | 104 | 1.04 | 0.471 |
| 7 | 18 | 0.71 | 0.563 |
| 8 | - | 0.90 | 0.690 |


| Year | Recruitment at age 1 '000 | Simulated catches '000 tons | | |
|------|------------------------------|-----------------------------|------------------------------|------------------------------|
| | | $F_{78} = F_{76}$ | $F_{78} = 0.9 \times F_{76}$ | $F_{78} = 0.8 \times F_{76}$ |
| 1977 | 77 800 | 22.4 | 22.4 | 22.4 |
| 1978 | 77 800 | 17.8 | 16.6 | 15.2 |

Table 6.7. Cod.
Division VIIa. Input data for catch prediction.

| Age | 1976 Catch '000 | F values 1976-1977 | Mean weight kg* |
|-----|--------------------|-----------------------|--------------------|
| 1 | 1 817 | 0.35 | 0.61 |
| 2 | 2 881 | 0.71 | 1.66 |
| 3 | 479 | 0.86 | 3.33 |
| 4 | 351 | 0.78 | 5.09 |
| 5 | 39 | 0.75 | 6.19 |
| 6 | 54 | 0.75 | 6.76 |
| 7+ | 15 | 0.75 | 8.30 |

| Year | Recruitment at age 1 '000 | Simulated catches ' 000 tons | | |
|------|------------------------------|------------------------------|------------------------------|------------------------------|
| | | $F_{78} = F_{76}$ | $F_{78} = 0.9 \times F_{76}$ | $F_{78} = 0.8 \times F_{76}$ |
| 1977 | 6 866 | 9.8 | 9.8 | 9.8 |
| 1978 | 6 866 | 9.3 | 8.6 | 7.9 |

*) These values had to be adjusted by -5.88% to yield the actual catch in weight in 1976.






Figure 2. North Sea whiting.




Figure 3. North Sea haddock.




Figure 4. North Sea cod.
Relationship between 1978 TAC and F_{78} as a percentage of F_{76} .




Figure 5. North Sea cod in Divisions IVa,b,c.
Predictive regression yearclass strength
on IYHS abundance estimates.

VPA
Millions of
1 year olds




Figure 6. North Sea haddock.
Predictive regression yearclass strength
(1967, 1974 data points omitted) on IYHS
abundance estimates.




Figure 7. North Sea whiting.
Predictive regression yearclass strength on
IYHS abundance estimates.